

Contains azoxystrobin, the active ingredient used in Heritage®.



FUNGICIDE



LONG-LASTING ORNAMENTAL DISEASE PROTECTION

Artavia™ 2 SC is a broad-spectrum, long-lasting strobilurin fungicide with both preventative and curative activity. It has excellent translaminar activity, providing long-term total protection for up to 28 days. Artavia 2 SC controls most major diseases in a variety of turf areas as well as in greenhouses and on specified ornamentals.

Artavia 2 SC encourages plants to utilize resources efficiently when applied as a preventative. It increases plant growth, water efficiency and maintains green leaf area. Plus, it provides outstanding protection to new plant growth making it an excellent choice for preventative disease control.

KEY BENEFITS

- Broad-spectrum control – effective on all four fungal groups
- Approved for ornamentals
- Systemic activity provides long-lasting control
- Tank-mix compatible with many fertilizers, insecticides and fungicides
- Convenient, 2 lb liquid material
- Contains ½ lb of active ingredient per quart

KEY USES

- Athletic fields
- Commercial and residential
- Field-grown ornamentals
- Golf courses
- Greenhouses
- Lawn and landscape
- Nurseries
- Parks and recreational areas
- Shade-houses
- Tobacco

PRODUCT NOTES

EPA REGISTRATION NUMBER

92647-1-91234

ACTIVE INGREDIENT

Azoxystrobin 22.9%

FORMULATION

Suspension concentrate

FRAC NUMBER

11

SIGNAL WORD

Caution

PACKAGE SIZE

12 x 1 pt

4 x 1 gal

265 gal

RESTRICTED USE

No

Refer to label for specific use restrictions.



www.AtticusLLC.com

APPLICATION INFORMATION

TURF

Artavia 2 SC should be applied prior to disease development. Mix Artavia 2 SC with the required amount of water and apply as a dilute spray application in 2 to 4 gal. of water/1,000 sq. ft. (87 to 174 gal/acre). Repeat applications at specified intervals for as long as required. For spot treatments, use 0.4 fl. oz. of Artavia 2 SC per 1 to 2 gal. of water. Do not apply more than 9.6 qt. of product/acre/year (7.1 fl. oz. product/1,000 sq. ft./year). Apply by ground only. Use the shortest specified application interval and/or use the higher specified rate when prolonged favorable disease conditions exist.

ARTAVIA 2 SC RATE CONVERSION CHART FOR TURF

FL OZ OF ARTAVIA 2 SC PER 1,000 SQ. FT.	OUNCES OF ACTIVE INGREDIENT PER 1,000 SQ. FT.	FL OZ OF ARTAVIA 2 SC PER ACRE	PINTS OF ARTAVIA 2 SC PER ACRE
0.4	0.104	17.4	1.1
0.5	0.13	21.8	1.4
0.6	0.156	26.1	1.6
0.7	0.182	30.5	1.9
0.77	0.2	33.5	2.1
1.35	0.35	58.8	3.7

AMOUNT OF ARTAVIA 2 SC TO MIX 100 GAL. FOR TURF APPLICATIONS

ARTAVIA 2 SC USE RATE (FL OZ)	SPRAY VOLUME (GAL/1,000 SQ. FT.)		
	2.0 GAL (FL OZ)	3.0 GAL (FL OZ)	4.0 GAL (FL OZ)
0.4	20	13	10
0.5	25	17	13
0.6	30	20	15
0.7	35	23	18
0.77	38.5	25.7	19.3
1.35	67.5	45	33.75

ORNAMENTALS

Apply Artavia 2 SC prior to disease development and continue throughout the season at specified intervals following resistance management guidelines. Artavia 2 SC works best when used as part of a preventative disease management program. Using Artavia 2 SC as a rescue (late curative or eradicant) treatment will not always result in satisfactory disease control.

Artavia 2 SC can be applied as a broadcast or banded spray targeted at the foliage or crown of the plant. Use sufficient water to ensure complete coverage of the target plant. Refer to the label for specific use directions for control of certain diseases. Applications may be made by ground only.

Apply 1.9 to 7.7 fl oz/100 gal (0.95 to 3.85 fl oz/50 gal) every 7 to 28 days (or as otherwise specified for a specific plant or disease). The addition of a non-silicone based wetter-sticker at the specified use rate may enhance coverage on hard-to-wet plant foliage.

Under most conditions and for most diseases, apply 3.85 to 7.7 fl oz/100 gal (1.9 to 3.85 fl oz/50 gal) on a 7 to 14 day interval.

Under light to moderate disease pressure, use the lower rates within the specified rate range (1.9 to 3.85 fl oz/100 gal, or 0.95 to 1.9 fl oz/50 gal) on a 7 to 14 day interval or the higher rates within the specified rate range (5.75 to 7.7 fl oz/100 gal or 2.85 to 3.85 fl oz/50 gal) on a 14 to 28 day interval.

Under environmental conditions which promote severe disease development, use the higher rates within the specified rate range (5.75 to 7.7 fl oz/100 gal or 2.85 to 3.85 fl oz/50 gal) on a 7 to 14 day interval.

Refer to product label for complete application and mixing instructions.

17262 ©2017 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. Artavia™ 2 SC is not manufactured, or distributed by Syngenta, seller of Heritage®. Heritage® is a registered trademark of Syngenta Group Company. Artavia™ is a trademark of Atticus LLC.

KEY DISEASES

Anthracnose leaf blight
 Black dot
 Blue mold
 Brown leaf spot
 Cercospora leaf spot
 Circle spot
 Early; blight, leaf spot
 Eye spot
 Frogeye leaf spot
 Gray leaf spot
 Kernel smut
 Late; blight, leaf spot
 Leaf smut
 Northern corn leaf; blight, spot
 Panicle blast
 Powdery mildew
 Pythium root rot
 Rhizoctonia solani
 Rust
 Scab
 Septoria leaf spot
 Sheath; blight, spot
 Southern corn leaf; blight, spot

(Refer to product label for complete list)