

TECHNICAL INFORMATION BULLETIN

AZOXYSTROBIN | GROUP 11 | FUNGICIDE



EPA Reg. No. 91234-74

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

Addition of Septoria Spot (*Septoria citri*) as Disease Controlled on Citrus Fruit

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for **Atticus Acadia™ 2 SC** before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Apply at the rate: 12.0 - 15.5 fl. oz. product/acre

Application Instructions:

Atticus Acadia™ 2 SC applications must begin prior to disease development and continue throughout the year on 7-to-12-day intervals following the resistance management guidelines. Under conditions that favor severe disease epidemics, the higher application rates must be used. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates. A horticultural spray oil must be used to improve control of greasy spot.

Do not apply more than two sequential applications of **Atticus Acadia™ 2 SC** or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.

Do not make more than four (4) applications of **Atticus Acadia™ 2 SC** or other Group 11 fungicide per year.



Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513