CONTAINS MESOTRIONE, THE ACTIVE INGREDIENT USED IN CALLISTO®

## SUPERIOR BROADLEAF WEED CONTROL

AVALLO<sup>M</sup> 4 SC

Atticus Cavallo<sup>™</sup> 4 SC is a selective herbicide that delivers superior contact and residual control of tough broadleaf weeds including velvet leaf, cocklebur, pigweed, sunflower and nightshades. It is an excellent tank-mix partner that is safe for application on labeled crops. Atticus Cavallo 4 SC is readily taken up by leaves, shoots or the roots offering a flexible pre- or post-emergence application window.

The highly selective HPPD inhibitor mode of action controls weeds resistant to glyphosate, ALS-inhibiting and triazine herbicides to ensure maximum yield potential in crops. It works to prevent the production of carotenoid which leads to plant death in broadleaf weeds.

Make Atticus Cavallo 4 SC an integral part of any weed management program where ease-of-use and outstanding broadleaf pest control are desired.

#### **KEY BENEFITS**

Atticus

- Tough on weeds, but gentle on crops
- · Flexible application window
- Superior broadleaf weed control
- · Long-lasting residual control
- Excellent tank-mix partner

#### **KEY USES**

- Asparagus
- Cranberry
- Field corn
- Seed corn
- Sorghum
- Soybean
- Sugarcane
- Sweet corn
- · Yellow popcorn

# PRODUCT NOTES

HERBICIDE

**EPA REGISTRATION NUMBER** 91234-75

ACTIVE INGREDIENT Mesotrione 40%

**FORMULATION** Suspension Concentrate

### HRAC NUMBER 27

SIGNAL WORD Caution

## PACKAGE SIZE

2 x 2.5 gal 265 gal

#### **RESTRICTED USE**

No





Bootstrapped and ready to serve, we deliver battle-tested chemistries and an experience like no other. Proud to be 100% American-owned, our mission is to help you every step of the way.

Atticus



#### **APPLICATION INFORMATION**

Atticus Cavallo<sup>™</sup> 4 SC can be applied by ground for pre- and post-emergence weed control.

In pre-emergent applications, weeds take up the product through the soil during weed emergence. Dry weather conditions can reduce pre-emergent effectiveness – at least 1/4-inch of rainfall must occur within 7 to 10 days of application; rotary hoeing activates Atticus Cavallo 4 SC.

In post-emergent applications, vulnerable weeds take up the product through treated foliage and stop growing soon after application. Atticus Cavallo 4 SC is absorbed by soil and/or through foliage of emerged weeds. Results may be reduced if applied to weeds that are under prolonged stress due to drought, heat, lack of fertility, flooding or prolonged cool temperatures. For best results, complete and uniform coverage is essential to actively growing weeds.

#### **USE RESTRICTIONS**

- Do not apply to white popcorn or ornamental (Indian) corn.
- Do not cultivate corn within seven days before or after application as weed control may be reduced.
- Do not apply through any type of irrigation system unless specified under the specific crop section of product label.
- Do not apply with suspension fertilizers as the carrier.
- Do not apply this product post-emergence in a tank-mix with emulsifiable concentrate grass herbicides, unless specifically directed under one of the tank-mix sections of this label, or crop injury can occur.
- Do not make aerial applications unless specified in the specific crop directions of product label.

#### **KEY WEEDS**

Amaranth Cocklebur Chickweed Hemp Horseweed Kochia Lambsquarter Morning glory Nightshade Nutsedge Pigweed Ragweed Sunflower Velvetleaf Waterhemp

(Refer to product label for complete list)



#### ATTICUSLLC.COM

Refer to product label for complete application and mixing instructions.

©2025 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. Atticus Cavallo™ 4 SC is not manufactured, or distributed by Syngenta, seller of Castillo®. Castillo® is a registered trademark of a Syngenta Group Company. Cavallo™ is a trademark of Atticus LLC. 23-AG-0082-B