

MESOTRIONE 40% FROM ATTICUS

Atticus

 **CAVALLO™**
4 SC



PROTECT YOUR STONE FRUIT AND TREE NUT YIELDS WITH BROADLEAF WEED CONTROL

Atticus Cavallo™ 4 SC is a selective herbicide that delivers superior contact and residual control of tough broadleaf weeds including velvet leaf, cocklebur, pigweed, sunflower and nightshades. It is an excellent tank-mix partner that is safe for application on labeled crops. Atticus Cavallo 4 SC is readily taken up by leaves, shoots or the roots offering a flexible pre- or post-emergence application window.

The highly selective HPPD inhibitor mode of action controls weeds resistant to glyphosate, ALS-inhibiting and triazine herbicides to ensure maximum yield potential in crops. It works to prevent the production of carotenoid which leads to plant death in broadleaf weeds.

Make Atticus Cavallo 4 SC an integral part of any weed management program where ease-of-use and outstanding broadleaf pest control are desired.

KEY BENEFITS

- Tough on weeds, but gentle on crops
- Flexible application window
- Superior broadleaf weed control
- Long-lasting residual control
- Excellent tank-mix partner

KEY USES

- **Stone Fruit - Group 12-12**
(Apricot, Japanese apricot, capulin, black cherry, Nanking cherry, sweet cherry, tart cherry, Chinese jujube, nectarine, peach, plum, American plum, beach plum, Canada plum, cherry plum, Chickasaw plum, Damson plum, Japanese plum, Klamath plum, prune plum, plumcot, sloe, cultivars, varieties and/or hybrids of these)
- **Tree Nuts - Group 14-12**
(African nut-tree, almond, beech nut, Brazil nut, Brazilian pine, bunya, bur oak, butternut, Cajou nut, candlenut, cashew, chestnut, chinquapin, coconut, Coquito nut, Dika nut, ginkgo, Guiana chestnut, hazelnut (filbert), heartnut, hickory nut, Japanese horse-chestnut, macadamia nut, Mongongo nut, monkeypot, monkey puzzle nut, Okari nut, Pachira nut, peach palm nut, pecan, pequi, pili nut, pine nut, pistachio, Sapucaia nut, tropical almond, black walnut, English walnut, yellowhorn, cultivars, varieties and/or hybrids of these)

PRODUCT INFORMATION

EPA REGISTRATION NUMBER
91234-75

ACTIVE INGREDIENT
Mesotrione 40%

FORMULATION
Suspension Concentrate

HRAC NUMBER
27

SIGNAL WORD
Caution

PACKAGE SIZE
4 x 1 gal
265 gal

RESTRICTED USE
No

Atticus

AtticusLLC.com

Always read and follow label instructions.

PRECAUTIONS FOR STONE FRUIT GROUP 12-12 AND TREE NUTS GROUP 14-12

1. To avoid crop injury, apply the spray to the grove or orchard floor and to the weeds, avoiding contact with crop foliage, stems or fruit. Contact of Atticus Cavallo 4 SC with the crop may result in bleaching injury that is typically temporary. Use trunk guards to protect plants until adequate bark has developed.
2. Specified rates are based on broadcast treatment. For band applications around trees in fruit or nut plantings, reduce the broadcast rate of Atticus Cavallo 4 SC and carrier per acre in proportion to the area actually sprayed. (See Banded Applications Section.)
3. Application of Atticus Cavallo 4 SC in nectarine, plum or tree nuts grown in coarse soils may cause bleaching, especially when applied during time of heavy water use and root growth such as during bud break or rapid shoot expansion.

RESTRICTIONS FOR STONE FRUIT GROUP 12-12 AND TREE NUTS GROUP 14-12

- Apply Atticus Cavallo 4 SC only to stone fruit and nut trees that have been established for a minimum of 12 months.
- **DO NOT** apply in stressed orchards - stressed due to poor weather or other abiotic factors.
- **DO NOT** exceed a total of 12 fl. oz. per acre (0.376 lb. a.i./A) of Atticus Cavallo 4 SC per year or in a 12-month period.
- **DO NOT** exceed 6 fl.oz. per acre (0.188 lb ai/A) of Atticus Cavallo 4 SC for the first application.
- **DO NOT** exceed 3 applications per year or in a 12-month period when using reduced application rates.
- Allow at least 5 months between applications of Atticus Cavallo 4 SC at 6 fl. oz./A and at least 6 weeks between applications of 6 fl. oz./A and subsequent applications of 3 fl. oz./A. (Applications must follow one of the four programs listed in Table 6 below.)
- Stone fruit or tree nuts Pre Harvest Interval (PHI) is 30 days.
- **DO NOT** use on soils with greater than 20% gravel.
- **DO NOT** apply Atticus Cavallo 4 SC through any type of irrigation system.
- **DO NOT** apply Atticus Cavallo 4 SC by air.

KEY WEEDS

Amaranth
Cocklebur
Chickweed
Hemp
Horseweed
Kochia
Lambsquarter
Morning glory
Nightshade
Nutsedge
Pigweed
Ragweed
Sunflower
Velvetleaf
Waterhemp

(Refer to product label for complete list)

For more information contact your local distributor or Atticus representative to learn how you can fight back against weeds and protect your crops.

©2021 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. Cavallo™ 4 SC is not manufactured, or distributed by Syngenta, seller of Castillo®. Castillo® is a registered trademark of Syngenta.