

# TALARIS<sup>™</sup> 70 WSB

FUNGICIDE

CONTAINS THIOPHANATE-METHYL, THE ACTIVE INGREDIENT USED IN TOPSIN® M WSB

## EFFECTIVE BROAD-SPECTRUM AND SYSTEMIC DISEASE CONTROL

Talaris<sup>™</sup> 70 WSB effectively protects valuable crops with broad-spectrum, systemic disease control. Available in an easy-to-handle water-soluble bag, Talaris 70 WSB mixes easily with other fungicides for optimum disease control and resistance management. Talaris 70 WSB is registered for application on many crops. After application, it works systemically to protect new growth from disease development.

### KEY BENEFITS

- High-quality disease control across many crops
- Broad-spectrum, systemic disease protection
- Protects new growth
- Convenient water-soluble bags for easy tank mixing

### KEY USES

- |                |              |                   |
|----------------|--------------|-------------------|
| ▪ Almonds      | ▪ Garlic     | ▪ Pistachios      |
| ▪ Apples       | ▪ Grapes     | ▪ Plums/prunes    |
| ▪ Apricots     | ▪ Nectarines | ▪ Potatoes        |
| ▪ Beans        | ▪ Onions     | ▪ Soybeans        |
| ▪ Canola       | ▪ Peanuts    | ▪ Strawberries    |
| ▪ Cherries     | ▪ Peaches    | ▪ Sugarbeets      |
| ▪ Conifer spp. | ▪ Pears      | ▪ Triticale Wheat |
| ▪ Cucurbits    | ▪ Pecans     |                   |

### PRODUCT NOTES

#### EPA REGISTRATION NUMBER

87373-6-91234

#### ACTIVE INGREDIENT

Thiophanate-methyl 70%

#### FORMULATION

Wettable powder

#### FRAC NUMBER

1

#### SIGNAL WORD

Caution

#### PACKAGE SIZE

6 x (5 x 1 lb WSB)

#### RESTRICTED USE

No



LABEL



PRODUCT INFO



PORTFOLIO

Refer to label for specific use restrictions.



Bootstrapped and ready to serve, we deliver battle-tested chemistries and an experience like no other. Proud to be 100% American-owned, our mission is to help you every step of the way.

Atticus

## APPLICATION INFORMATION

### AERIAL APPLICATIONS TO TREE CROPS

Use a minimum of 10 gal/acre for aerial application to fruit tree crops. Increased fungicidal activity is related to coverage and timing, increased volumes are required as crop canopy density increases.

NOTE: Conifer applications require higher spray volumes, use lower volumes with mist type applicators and highest volumes with conventional types.

### ROW CROP APPLICATIONS

Use a minimum of 5 gal/acre for ground application, however most ground applications should be made with 10 to 20 gal/acre as cropping situations dictate. Increased fungicidal activity is related to coverage and timing, increased volumes are required as crop canopy density increases.

## RESISTANCE MANAGEMENT

To avoid the development of disease resistance, Talaris™ 70 WSB should be used as part of a resistance management strategy with fungicides of different modes of action. Atticus does not recommend the use of products containing thiabendazole in combination or rotation with Talaris 70 WSB. These use similar chemistry and mode of action and can contribute to development of disease tolerance.

## KEY DISEASES

Anthracnose  
Apple scab  
Belly rots  
Black pox  
Black rot  
Botrytis bunch rot  
Brown rot  
Brown rot blossom blight  
Brown spot  
Cercospora leaf spot  
Cherry leaf spot  
Downy spot  
Early leaf spot  
Eyespot  
Flyspeck  
Foot rot  
Frogeye leaf spot  
Fruit brown rot  
Fruit rot  
Gray mold  
Gummy stem blight  
Jacket rot  
Late leaf spot  
Leaf blight  
Leaf scorch  
Leaf spot  
Liver spot  
Peach scab  
Pear scab  
Penicillium clove rot  
Pod and stem blight  
Powdery mildew  
Purple seed stain  
Rust  
Scab  
Sclerotinia stem rot  
Shoot blight  
Sooty blotch  
Soybean rust  
Stem end blight  
Strawbreaker  
Target spot  
Web blotch  
White mold  
White rot

*(Refer to product label for complete list)*