



FUNGICIDE

TALARIS[™]

70 WSB



EFFECTIVE BROAD-SPECTRUM AND SYSTEMIC DISEASE CONTROL

Talaris[™] 70 WSB effectively protects valuable crops with broad-spectrum, systemic disease control. Available in an easy-to-handle water-soluble bag, Talaris 70 WSB mixes easily with other fungicides for optimum disease control and resistance management. Talaris 70 WSB is registered for application on many crops. After application, it works systemically to protect new growth from disease development.

KEY BENEFITS

- High-quality disease control across many crops
- Broad-spectrum, systemic disease protection
- Protects new growth
- Convenient water-soluble bags for easy tank mixing

KEY USES

- | | | |
|----------------|--------------|-------------------|
| • Almonds | • Garlic | • Pistachios |
| • Apples | • Grapes | • Plums/prunes |
| • Apricots | • Nectarines | • Potatoes |
| • Beans | • Onions | • Soybeans |
| • Canola | • Peanuts | • Strawberries |
| • Cherries | • Peaches | • Sugarbeets |
| • Conifer spp. | • Pears | • Triticale Wheat |
| • Cucurbits | • Pecans | |

PRODUCT NOTES

EPA REGISTRATION NUMBER

87373-6-91234

ACTIVE INGREDIENT

Thiophanate-methyl 70%

FORMULATION

Wettable powder

FRAC NUMBER

1

SIGNAL WORD

Caution

PACKAGE SIZE

6 x (5 x 1 lb WSB)

RESTRICTED USE

No

Refer to label for specific use restrictions.

APPLICATION INFORMATION

AERIAL APPLICATIONS TO TREE CROPS

Use a minimum of 10 gal/acre for aerial application to fruit tree crops. Increased fungicidal activity is related to coverage and timing, increased volumes are required as crop canopy density increases.

NOTE: Conifer applications require higher spray volumes, use lower volumes with mist type applicators and highest volumes with conventional types.

ROW CROP APPLICATIONS

Use a minimum of 5 gal/acre for ground application, however most ground applications should be made with 10 to 20 gal/acre as cropping situations dictate. Increased fungicidal activity is related to coverage and timing, increased volumes are required as crop canopy density increases.

RESISTANCE MANAGEMENT

To avoid the development of disease resistance, Talaris™ 70 WSB should be used as part of a resistance management strategy with fungicides of different modes of action. Atticus does not recommend the use of products containing thiabendazole in combination or rotation with Talaris 70 WSB. These use similar chemistry and mode of action and can contribute to development of disease tolerance.

KEY DISEASES

Anthracnose
Apple scab
Belly rots
Black pox
Black rot
Botrytis bunch rot
Brown rot
Brown rot blossom blight
Brown spot
Cercospora leaf spot
Cherry leaf spot
Downy spot
Early leaf spot
Eyespot
Flyspeck
Foot rot
Frogeye leaf spot
Fruit brown rot
Fruit rot
Gray mold
Gummy stem blight
Jacket rot
Late leaf spot
Leaf blight
Leaf scorch
Leaf spot
Liver spot
Peach scab
Pear scab
Penicillium clove rot
Pod and stem blight
Powdery mildew
Purple seed stain
Rust
Scab
Sclerotinia stem rot
Shoot blight
Sooty blotch
Soybean rust
Stem end blight
Strawbreaker
Target spot
Web blotch
White mold
White rot

(Refer to product label for complete list)

Refer to product label for complete application and mixing instructions.

17344 ©2017 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state.
Talaris™ 70 WSB is not manufactured or distributed by United Phosphorus, Inc., seller of Topsin® M WSB.
Topsin® is a trademark of Nippon Soda Company, Ltd. Talaris™ is a trademark of Atticus, LLC.