

TECHNICAL INFORMATION BULLETIN

AZOXYSTROBIN	GROUP	11	FUNGICIDE
DIFENOCONAZOLE	GROUP	3	FUNGICIDE



EPA Reg. No. 91234-205

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

For Control of *Botryosphaeria* Panicle and Shoot Blight in Walnuts

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for **Acadia™ ESQ Fungicide** before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Application rate: 12 - 14 fl. oz. product/A of **Acadia™ ESQ Fungicide** to control *Botryosphaeria* panicle and shoot blight. Make a second application 14 – 21 days later then alternate to another fungicide with a non-QoI (Group 11) mode of action.

Follow all use directions under the Tree Nuts group section on the **Acadia™ ESQ Fungicide** container label.

Do not apply more than 56 oz. product of **Acadia™ ESQ Fungicide** per crop/year.
Do not apply within 45 days of harvest (45-day PHI).



Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513