

TECHNICAL INFORMATION BULLETIN

AZOXYSTROBIN	GROUP	11	FUNGICIDE
DIFENOCONAZOLE	GROUP	3	FUNGICIDE



EPA Reg. No. 91234-205

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

For Control of Red Leaf Blotch (*Polystigma amygdalinum*) on Almonds

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for **Acadia™ ESQ Fungicide** before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Application rate: For control of Red Leaf Blotch (RLB) (*Polystigma amygdalinum*) on almonds, apply **Acadia™ ESQ Fungicide** at 14.0 fl. oz./A as part of a two-spray application program. Applications should be made prior to disease presence and occur at petal fall with a secondary application with an effective product on RLB, 2 – 3 weeks post-petal-fall in a seasonal fungicide spray program.

Follow all use directions, restrictions, and precautions under the Almonds section on the **Acadia™ ESQ Fungicide** container label.



Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513