



HERBICIDE

CONTAINS SULFENTRAZONE AND CLORANSULAM-METHYL,
THE ACTIVE INGREDIENTS USED IN AUTHORITY® FIRST DF AND SONIC®

SETTING YOUR SOYBEAN CROP UP FOR SUCCESS

Maximum yield is the first goal in soybean production so your first step should be to protect your crop from broadleaf and grass weeds with Aquesta™ FR. Aquesta FR contains sulfentrazone and cloransulam-methyl, the active ingredients found in Authority® First DF and Sonic®. This powerful product provides pre-emergent protection by infiltrating the roots and shoots of cocklebur, green foxtail, ragweed, wild mustard, and many other weeds.

Aquesta FR can be used alone or tank mixed with fertilizers, insecticides, or other herbicides for maximum flexibility. Aquesta FR is adaptable to all tillage systems and can be used as part of a spring burndown program when tank mixed with a compatible burndown herbicide. Adequate rainfall of 1/2-1" is needed for herbicide activation, and once activated, Aquesta FR helps to control many of the weeds that will rob you of the yield you work hard to achieve.

KEY BENEFITS

- Pre-emergent protection against broadleaf weeds
- Prevention of weed resistance through dual modes of action
- Can be used in no-till, minimum-till, and conventional tillage systems

KEY USES

- Soybeans

PRODUCT NOTES

EPA REGISTRATION NUMBER

91234-41

ACTIVE INGREDIENTS

Sulfentrazone 62.1%
Cloransulam-methyl 7.9%

FORMULATION

Water Dispersible Granule

HRAC NUMBERS

14
2

SIGNAL WORD

Caution

PACKAGE SIZE

2 x 10 lb

RESTRICTED USE

No



LABEL



PRODUCT INFO



PORTFOLIO



Bootstrapped and ready to serve, we deliver battle-tested chemistries and an experience like no other. Proud to be 100% American-owned, our mission is to help you every step of the way.





APPLICATION INFORMATION

- Use a standard low pressure herbicide boom sprayer equipped with suitable nozzles and screens. Apply uniformly using properly calibrated nozzles (10 to 40 psi) and screens and strainers no finer than 50 mesh.
- Use 10 to 40 gallons of spray solution per acre. Do not exceed 40 psi spray pressure unless required by the spray nozzle manufacturer.
- Continuous agitation during application is required. Avoid swath overlaps. Shut off spray booms while turning, slowing or stopping, as over application may result. Do not allow Aquesta FR spray mixtures to sit overnight as settling of product and difficulty of re-suspending may occur.
- Avoid all direct, and/or indirect spray contact with non-target plants. Do not apply near desirable vegetation. Allow adequate distance between target area and desirable plants to minimize exposure.

BEST PRACTICES:

- Pre-emergence and pre-plant incorporated applications of Aquesta FR require rainfall or irrigation to activate the herbicide.
- The amount of rainfall or irrigation required for activation following application depends on existing soil moisture, organic matter content, and soil texture.
- Observe all instructions, crop restrictions, mixing directions, application precautions, replanting directions, rotational crop guidelines, and other label information of each product when tank mixing Aquesta FR.

KEY WEEDS

Barnyardgrass
Cocklebur
Common Sunflower
Green foxtail
Kochia
Lambsquarters
Marestail
Moringglory species
Nightshade species
Palmer Amaranth
Pigweed
Ragweed
Russian Thistle
Smartweed Species
Velvetleaf
Venice Mallow
Waterhemp
Wild Mustard
Yellow foxtail

(Refer to product label for complete list)