



BRAIZE™

35 WSB



Contains pyriproxyfen, the active ingredient used in Seize® 35 WP.

Insect Growth Regulator

| | | |
|---|-------|---------------|
| ACTIVE INGREDIENT: | | By Wt. |
| *Pyriproxyfen..... | | 35.0% |
| OTHER INGREDIENTS: | | 65.0% |
| TOTAL: | | 100.0% |
| *2-[1-methyl-2-(4-phenoxyphenoxy)ethoxy]pyridine | | |
| Braize™ 35 WSB is a wettable powder formula containing 35% active ingredient. | | |
| EPA Reg. No.: 91234-63 | | |

KEEP OUT OF REACH OF CHILDREN CAUTION

See below for additional Precautionary Statements.

| FIRST AID | |
|---|--|
| If in eyes: | <ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. ▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ▪ Call a poison control center or doctor for treatment advice. |
| If inhaled: | <ul style="list-style-type: none"> ▪ Move person to fresh air. ▪ If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. ▪ Call a poison control center or doctor for further treatment advice. |
| If swallowed: | <ul style="list-style-type: none"> ▪ Call a poison control center or doctor immediately for treatment advice. ▪ Have person sip a glass of water if able to swallow. ▪ Do not induce vomiting unless told to do so by the poison control center or doctor. ▪ Do not give anything by mouth to an unconscious person. |
| If on skin or clothing: | <ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15 - 20 minutes. ▪ Call a poison control center or doctor for further treatment advice. |
| HOT LINE NUMBER | |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information. | |

**For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident,
 Call CHEMTREC Day or Night Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long-sleeved shirt and long pants, shoes plus socks.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear:

- Long-sleeved shirt,
- Long pants,
- Socks plus shoes.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by disposing of equipment washwaters or rinsate. Avoid direct application and/or spray drift to beehives.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow to come in contact with oxidizing agents. Hazardous chemical reaction will occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE), and Restricted-Entry Interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants,
- Waterproof gloves,
- Socks plus shoes.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of the user, applicator and/or applicator advisor, to the extent allowable by applicable law.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

RESISTANCE MANAGEMENT

For resistance management, **Braize 35 WSB** contains a Group 7 Insecticide. Any insect population may contain individuals naturally resistant to **Braize 35 WSB** and other Group 7 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of **Braize 35 WSB** or other Group 7 insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact your Atticus, LLC representative.

Refer to specific crop sections for **MANAGING INSECT RESISTANCE** language applicable to that crop.



SPRAY DRIFT

Aerial Applications

- Do not release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a fine or coarser droplet size (ASABE S572.1), except for when applying ultra-low volume applications.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Do not apply during temperature inversions.

Airblast Applications

- All sprays must be directed into the canopy.
- Nozzles directed out of the orchard must be turned off when treating the outer row, or when making turns between rows.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Groundboom Applications

- User must only apply with the nozzle height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
 - Applicators are required to use a fine or coarser droplet size (ASABE S572.1).
 - Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

**THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.**

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

Adjust Nozzles – Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

WATER SOLUBLE PACKAGING

MULTIPLE PACKAGING

This bag contains multiple water-soluble packets of **Braize 35 WSB**. Do not handle the packets with wet gloves or allow the packets to become wet prior to addition to spray tank. Do not break open packets. Refer to the **Tables of Equivalents** to calculate the number of packets to use. If all packets are not used, close and reseal outer container to protect remaining packet(s).

Table of Equivalents

| Rates of Braize 35 WSB Per Acre | # of Water Soluble Packets To Treat 4 Acres |
|---------------------------------|---|
| 3 oz | 3 |
| 4 oz | 4 |
| 5 oz | 5 |

PRODUCT INFORMATION

Braize 35 WSB is intended for use in Integrated Pest Management (IPM) or Insect Resistance Management (IRM) programs. **Braize 35 WSB** will not control insect adults, and it is recommended to be used in combination and/or rotation with other IPM or IRM materials. Contact your local state extension service for details.

Braize 35 WSB acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. **Braize 35 WSB** has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since **Braize 35 WSB** is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

Restriction:

- Do not apply this product through any type of irrigation system.



POME FRUIT

(Crop Group 11-10)

Braize 35 WSB acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. **Braize 35 WSB** has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since **Braize 35 WSB** is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

| Crops | Pests | Product Rates Per Acre | Application Instructions |
|---|---|------------------------|---|
| Apple Azarole Crabapple Loquat Mayhaw Medlar Quince Quince, Chinese Quince, Japanese Tejocote Cultivars, varieties, and/or hybrids of these | Codling Moth Suppression of: Obliquebanded Leafroller Pandemis Leafroller | 4 to 5 oz | Apply by ground with airblast equipment in a minimum of 50 gals/A. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage. 1st Generation Codling Moth: Apply at Biofix plus 100 degree days (this is usually close to petal fall). Make a second application 14 to 21 days later or timed to coincide with peak moth flights. For subsequent applications, if needed, apply other insecticides registered for Codling Moth control. The addition of oil has been shown to increase Codling Moth activity of Braize 35 WSB . Follow local recommendations regarding in-season use and rates of oil in your area. Obliquebanded and Pandemis Leafroller: Apply to the overwintering larvae at pink to petal fall tree growth stage. Braize 35 WSB does not directly control the larvae but inhibits normal pupation and adult emergence. |
| | Florida Red Scale Lecanium Scale San Jose Scale | 4 to 5 oz | Apply by ground with airblast equipment in a minimum of 50 gals/A. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage. Apply at delayed dormant through pink. For delayed dormant use a superior spray oil at the manufacturer's recommended rate of oil. Under heavy scale pressure, use the higher rate of Braize 35 WSB . Scale: For control in-season, make application in cover sprays when scale crawlers first emerge. The use of oil at the crawler emergence stage has been shown to improve control. Follow local recommendations regarding in-season use and rates of oil in your area. |
| | Apple Aphid Rosy Apple Aphid Spirea Aphid Spotted Tentiform Leafminer Western Tentiform Leafminer | 3 to 5 oz | Apply by ground with airblast equipment in a minimum of 50 gals/A. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage. Leafminers: Apply at peak moth flight for control of 1 st or 2 nd generation. This usually occurs at pre-pink to pink apple stage for 1 st generation and 2 nd cover for 2 nd generation. Rosy Apple Aphid: Apply at 1/2" green apple growth stage. Apple and Spirea Aphid: Apply when aphids first appear, but before aphids are colonized. TIMING IS CRITICAL FOR ACCEPTABLE APHID CONTROL. |
| Pear Pear, Asian | Pear Psylla | 4 to 5 oz | Apply by ground with airblast equipment in 75 to 400 gals/A with a superior spray oil at manufacturer's recommended rates for the stage of growth. It is essential to use adequate water volume to ensure thorough coverage. Apply a single application during the delayed dormant to pink stage of growth at 5 oz/A or make two applications from delayed dormant through petal fall stage of growth at 4 to 5 oz/A. Make application just before sustained egg lay is anticipated. DO NOT SKIP ROWS DURING APPLICATION. For in-season control of pear psylla, use an insecticide recommended for this use. |
| | Lecanium Scale San Jose Scale | 4 to 5 oz | Apply by ground with airblast equipment in 100 to 400 gals/A at delayed dormant with a superior spray oil at the manufacturer's recommended rates. Under heavy scale pressure, use the higher rate. Apply in sufficient water volume for tree size. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage. For control of scale in-season make application in cover sprays when scale crawlers first emerge. The use of oil at the crawler emergence stage has been shown to improve performance. Follow local recommendations regarding in-season use and rates of oil in your area. |
| | Codling Moth Suppression of: Obliquebanded Leafroller Pandemis Leafroller | 4 to 5 oz | Apply by ground with airblast equipment in 100 to 400 gals/A. 1st Generation Codling Moth: Apply at Biofix plus 100 degree days. Make a second application 14 to 21 days later or timed to coincide with peak moth flights. For subsequent applications, if needed, apply other insecticides registered for Codling Moth control. The addition of oil has been shown to increase Codling Moth activity of Braize 35 WSB . Follow local recommendations regarding in-season use and rates of oil in your area. For additional generations apply other insecticides registered for Codling Moth control. |

Braize 35 WSB may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.

Restrictions:

- Do not apply within 45 days of harvest.
- Do not make more than 2 **Braize 35 WSB** applications per growing season.
- Do not exceed 10 oz of **Braize 35 WSB** per acre per season.
- Do not apply earlier than 14 days after last **Braize 35 WSB** treatment.
- Regardless of formulation, do not apply more than 0.218 lb pyriproxyfen per acre per season.

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Atticus, LLC is restricting **Braize 35 WSB** to no more than two applications per growing season. **Braize 35 WSB** may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.



SMALL FRUIT VINE CLIMBING CROPS (EXCEPT GRAPE)

(Subgroup 13-07E)

Not For Use in New York

Braize 35 WSB acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. **Braize 35 WSB** has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since **Braize 35 WSB** is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

| Crops | Pests | Product Rates Per Acre | Application Instructions |
|---|---|------------------------|--|
| Berry, Schisandra* Gooseberry* Grape, Amur River* Kiwifruit, Fuzzy Kiwifruit, Hardy | Maypop* Cultivars, varieties and/or hybrids of these *Not for use in CA | Scale Whitefly | 5 oz Apply with ground or air equipment as a full coverage spray (minimum of 5 gals/A by air or 50 gals/A by ground). Thorough coverage is critical for adequate control. It is essential to use adequate water volume to ensure thorough coverage. Apply at either dormant, delayed dormant or an in-season cover spray. At dormant or delayed dormant use a spray oil at the manufacturer's recommended oil rate. In a cover spray make applications when scale crawlers first emerge. Follow local recommendations regarding in-season use and rates of oil in your area. |

Braize 35 WSB may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.

Restrictions:

- Do not apply within 30 days of harvest.
- Do not make more than 2 **Braize 35 WSB** applications per growing season.
- Do not exceed 10 oz of **Braize 35 WSB** per acre per season.
- Do not apply earlier than 14 days after last **Braize 35 WSB** treatment.
- Regardless of formulation, do not apply more than 0.218 lb pyriproxyfen per acre per season.

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Atticus, LLC is restricting **Braize 35 WSB** to no more than two applications per growing season. **Braize 35 WSB** may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

STONE FRUIT

(Crop Group 12)

Braize 35 WSB acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. **Braize 35 WSB** has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since **Braize 35 WSB** is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

| Crops | Pests | Product Rates Per Acre | Application Instructions |
|--|--|------------------------|---|
| Apricot Cherry Chickasaw Plum Damson Plum Fresh Prune Japanese Plum Nectarine Peach Plum Plumcot Prune | Lecanium Scale San Jose Scale Suppression of: Peach Twig Borer Oriental Fruit Moth | 4 to 5 oz | Apply in 100 to 400 gals/A with ground equipment. Coverage is essential for good control. Apply in sufficient water volume for tree size. Use of higher water volume will assure better coverage. Scale: Apply at dormant, delayed dormant or as an in-season cover spray: At dormant, delayed dormant, use a spray oil at the manufacturer's recommended oil rate. Under heavy scale pressure, use the higher rate of Braize 35 WSB . Peach Twig Borer: Braize 35 WSB has provided good suppression of Peach Twig Borer when applied delayed dormant through bloom sprays. Use a spray oil at the manufacturer's recommended rate for the stage of crop growth. |

In-season sprays of **Braize 35 WSB** may not protect fruit from blemishes by scale.

Braize 35 WSB may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.

Restrictions:

- Do not apply within 14 days of harvest.
- Do not make more than 3 **Braize 35 WSB** applications per growing season.
- Do not exceed 15 oz of **Braize 35 WSB** per acre per season.
- Do not apply earlier than 14 days after last **Braize 35 WSB** treatment.
- Regardless of formulation, do not apply more than 0.327 lb pyriproxyfen per acre per season.

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Atticus, LLC is restricting **Braize 35 WSB** to no more than three applications per growing season. **Braize 35 WSB** may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

TREE NUTS

(Crop Group 14)

Braize 35 WSB acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. **Braize 35 WSB** has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since **Braize 35 WSB** is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

| Crops | Pests | Product Rates Per Acre | Application Instructions |
|--------|--|------------------------|--|
| Almond | Black Scale San Jose Scale Suppression of: Peach Twig Borer | 4 to 5 oz | Apply with ground equipment in 100 to 400 gals/A. Coverage is essential for good control. Apply in sufficient water volume for tree size. Use of higher water volume will assure better coverage. Apply at either dormant, delayed dormant or an in-season cover spray. At dormant, delayed dormant use a spray oil at the manufacturer's recommended oil rate. In a cover spray make applications when scale crawlers first emerge. Follow local recommendations regarding in-season use and rates of oil in your area. Braize 35 WSB has provided good suppression of Peach Twig Borer when applied delayed dormant through bloom sprays. |

Restrictions:

- Do not apply within 21 days of harvest.
- Do not make more than 1 **Braize 35 WSB** application per growing season.
- Do not exceed 5 oz of **Braize 35 WSB** per acre per season.
- Regardless of formulation, do not apply more than 0.109 lb pyriproxyfen per acre per season.

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Atticus, LLC is restricting **Braize 35 WSB** to no more than one application per growing season. **Braize 35 WSB** may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal Integrated Pest Management (IPM) and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

| Crops | Pests | Product Rates Per Acre | Application Instructions |
|---|---|------------------------|--|
| Beechnut Brazil Butternut Cashew Chestnut Chinquapin Filbert (Hazelnut) Hickory Macadamia Pecan Pistachio Walnut | Codling Moth Dusky Veined Aphid Walnut Aphid | 5 oz | Apply with ground equipment in 100 to 400 gals/A. Apply in sufficient water volume for tree size. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage. 1st Generation Codling Moth: Apply at Biofix plus 100 degree days. Make a second application 14 to 21 days later or timed to coincide with peak moth flights. For subsequent applications, if needed, apply other insecticides registered for Codling Moth control. The addition of oil has been shown to increase Codling Moth activity of Braize 35 WSB . Follow local recommendations regarding in-season use and rates of oil in your area. |
| | Black Scale Brown Soft Scale Italian Scale Lecanium Scale San Jose Scale Walnut Scale Suppression of: Peach Twig Borer | 4 to 5 oz | Apply with ground equipment in 100 to 400 gals/A at either dormant, delayed dormant or an in-season cover spray. The addition of a superior spray oil at dormant or delayed dormant may improve scale control. However, do not use a spray oil in oil sensitive crops such as Walnut. Follow manufacturer's recommendations regarding use and rates of oil for your area. In a cover spray make applications when scale crawlers first emerge. Follow local recommendations regarding in-season use and rates of oil in your area. Under heavy scale pressure, use the higher rate. Coverage is essential for good control. Apply in sufficient water volume for tree size. Use of higher water volume will assure better coverage. |

Braize 35 WSB may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.

Restrictions:

- Do not apply within 21 days of harvest.
- Do not make more than 2 **Braize 35 WSB** applications per growing season.
- Do not exceed 10 oz of **Braize 35 WSB** per acre per season.
- Do not apply earlier than 14 days after last **Braize 35 WSB** treatment.
- Regardless of formulation do not apply more than 0.218 lb pyriproxyfen per acre per season.

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Atticus, LLC is restricting **Braize 35 WSB** to no more than two applications per growing season. **Braize 35 WSB** may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment.

PESTICIDE STORAGE: Store in a cool, dry place. Keep pesticide in original container. Keep container closed when not in use. Do not put concentrate or dilute into food or drink containers. Not for use or storage in or around the home.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable outer bag. Do not reuse or refill the outer bag. Completely empty bag into application equipment. Then dispose the empty bag in a sanitary landfill or by other procedures allowed by State and local authorities.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Braize™ 35 WSB is a trademark of Atticus, LLC

Seize® is a registered trademark of Valent U.S.A. Corporation.

20210317a

SPECIMEN