

CONTAINS CLETHODIM, THE ACTIVE INGREDIENT USED IN SELECT[®] 2 EC HERBICIDE

DEPENDABLE GRASS CONTROL SOLUTION

Ceridian[™] 2 EC is a versatile herbicide for control of actively growing grasses in a variety of crops. Keep your soybean, alfalfa, dry bean, peanut, and other crops free from grasses, giving them room to thrive. Apply this product post-emergence to actively growing grass weeds, rendering them weak with reduced vigor and growth. Symptoms of control will be observed in 1-2 weeks. For best results, apply at early growth once grasses reach recommended growth stage. Make Ceridian 2 EC your go-to product for grass control.

KEY BENEFITS

- Exceptional control of annual and perennial grasses
- Flexible application sites; can be used on a variety of crops
- Volunteer corn control in soybeans

KEY USES

- Alfalfa
- Canola
- Dry Beans/Peas
- Peanuts
- Soybeans
- Sugarbeets
- Sunflowers

PRODUCT NOTES

EPA REGISTRATION NUMBER

91234-154

ACTIVE INGREDIENT

Clethodim 26.4%

FORMULATION

Emulsifiable Concentrate

HRAC NUMBER

1

SIGNAL WORD

Warning

PACKAGE SIZE

2 x 2.5 gal
265 gal

RESTRICTED USE

No



ENGLISH LABEL



SPANISH LABEL



PORTFOLIO

APPLICATION INFORMATION

AVOID SPRAY DRIFT

Do not allow spray from ground or aerial equipment to drift onto adjacent land or crops. When drift may be a problem, do everything possible to reduce spray drift, including:

- Do not spray if wind speeds are or become excessive.
- Do not spray if wind speed is 10 mph or greater. If sensitive crops or plants are downwind, extreme caution must be used under all conditions.
- Do not spray if winds are gusty.
- Use extreme caution when conditions are favorable for drift (high temperatures, drought, low relative humidity), especially when sensitive plants are located nearby.
- Do not apply when a temperature inversion exists. If inversion conditions are suspected, consult with local weather services before making an application.

Timing of Applications

Apply this product post-emergence to actively growing grasses according to rate table recommendations. Applications made to grass plants stressed by insufficient moisture or cold temperatures or to grass plants exceeding recommended growth stages may result in unsatisfactory control. Do not apply under these conditions.

In arid regions where irrigation is used to supplement limited rainfall, this product should be applied as soon as possible after irrigation (within 7 days). A second application of this product will generally provide more effective control of perennial grass weeds than a single application. Make a second application to actively growing grass 2 to 3 weeks after emergence of new growth. Cultivation of treated grasses 7 days prior to or within 7 days after application of this product may reduce weed control. Do not apply this product if rainfall is expected within one hour since control may be reduced.

KEY WEEDS

Amazon Sprangletop
Annual Bluegrass
Annual Grasses
Barnyardgrass
Bearded Sprangletop
Bentgrass
Bermudagrass
Broadleaf Signalgrass
California Brome
Canarygrass
Cheatgrass
Crowfootgrass
Downy Brome
Fall Panicum
Field Sandbur
Foxtail
Goosegrass
Green Foxtail
Hairy Crabgrass
Hardy Ryegrass
Italian Ryegrass
Itchgrass
Johnsongrass
Junglerice
Kentucky Bluegrass
Large Crabgrass
Lovegrass
Mexican Sprangletop
Orchardgrass
Perennial Bluegrass
Perennial Grasses
Poa Annua
Quackgrass
Rabbitfootgrass
Red Rice
Red Sprangletop
Ripgut Brome
Roughstalk Bluegrass
Shattercane
Smooth Crabgrass
Southern Crabgrass
Stinkgrass
Tall Fescue
Texas Panicum
Volunteer Corn
Volunteer Oat
Volunteer Rye
Volunteer Sorghum
Volunteer Wheat
Wild Oat
Wild Proso Millet
Witchgrass
Woolly Cupgrass

(Refer to product label for complete list)