

INSECTICIDE

CONTAINS PROPARGITE, THE ACTIVE INGREDIENT USED IN OMITE® 6E

END MITES QUICKLY WITH ENDOMITE™

Endomite™ is a powerful miticide that makes an excellent addition to any grower's arsenal of products. Endomite goes to work quickly, knocking out any mite that stands in its way. It controls a wide range of mites and provides effective control at every life stage.

Endomite is highly effective under extreme mite pressure. The unique fuming action of the proven active ingredient, propargite, seeks out mites wherever they hide. Plus, Endomite has the power to penetrate webbing, making it an excellent rescue treatment.

Endomite is rainfast and provides extended residual control protecting crops throughout the growing season.

KEY BENEFITS

- Full-life cycle control on a variety of mites
- Unique fuming effect provides complete coverage
- Immediate, long-lasting control of mites but gentle on beneficial insects
- Excellent rescue treatment highly effective under high mite pressure

KEY USES

- Almonds
- Cherries
- Grapes
- Hops
- Walnuts

PRODUCT NOTES

EPA REGISTRATION NUMBER 91234-33

ACTIVE INGREDIENT Propargite 69.6%

FORMULATIONEmulsifiable concentrate

IRAC NUMBER

SIGNAL WORD
Danger

PACKAGE SIZE 2 x 2.5 gal

RESTRICTED USE Yes







PRODUCT INFO



PORTFOLIO



Bootstrapped and ready to serve, we deliver battle-tested chemistries and an experience like no other. Proud to be 100% American-owned, our mission is to help you every step of the way.





APPLICATION INFORMATION

Endomite™ is a liquid emulsifiable concentrate used in sprays for controlling mites. To ensure complete coverage and achieve the best results, applying Endomite to both the upper and lower leaf surfaces as well as fruit is recommended. Endomite is most effective in daytime temperatures averaging above 70°F. For best results apply when mites first appear at the full label rate.

INSTRUCTIONS AND RATES FOR USE ON BEARING CROPS

CROP	ALMONDS (CALIFORNIA & ARIZONA ONLY)						
Mites Controlled	Clover, European red, Strawberry spider, Two-spotted spider, Pacific spider						
Application Rate (fl oz/A)	32-64	Restricted Entry Interval (Days)	22 ¹	Pre-Harvest Interval (Days)	28		
Application Instructions	Apply high rate to areas with high mite infestation and to areas which historically have had high mite infestations.						
	Ground Applications	Apply in a minimum of 50 gallons total volume per acre.					
	Aerial Application	Apply in a minimum of 15 gallons total volume per acre. One application of Endomite may be applied by air to almonds over one ground application of Endomite or other propargite formulation.					
Restrictions	 Do NOT make more than 2 applications per season. Do NOT apply within 21 days of the last application. 						
Notes	¹REI Exception: There is no exception allowed for hand weeding or thinning.						

CROP	WALNUTS						
Mites Controlled	European red, Two-spotted spider						
Application Rate (fl oz/A)	32-68	Restricted Entry Interval (Days)	30¹	Pre-Harvest Interval (Days)	21		
Application Instructions	Apply high rate to areas with high mite infestation and to areas which historically have had high mite infestations. Use a minimum of 64 fl. oz. of Endomite per acre on large mature trees.						
	Ground Applications	Apply in a minimum of 100 gallons total volume per acre.					
	Aerial Application	Apply in a minimum of 20 gallons total volume per acre. Use 48-64 fl. oz. of Endomite when applied by air.					
Restrictions	 Do NOT apply more than 136 fl. oz. (6.4 lbs. propargite a.i.) of Endomite per acre per year Do NOT make more than 2 applications per season. Do NOT apply within 21 days of the last application. 						
Notes	¹REI Exception: In addition to early entry exceptions allowed by the Worker Protection Standard, you may enter or allow workers to enter treated areas to harvest by tree shaking after 21 days following application, as long as the workers wear long pants, long sleeved shirt and shoes plus socks. There is no exception allowed for hand weeding or thinning.						

KEY INSECTS

Clover mites
European red mites
Pacific spider mites
Strawberry spider mites
Two-spotted spider mites

(Refer to product label for complete list)

ATTICUSLLC.COM





