



HERBICIDE

CONTAINS FOMESAFEN, THE ACTIVE INGREDIENT USED IN FLEXSTAR®

YOUR PARTNER FOR POST-EMERGENCE WEED CONTROL

Forsyte™ 1.88 SL is an optimal partner for post-emergence weed control. Forsyte 1.88 SL controls waterhemp and many glyphosate and ALS resistant weeds. Plus, Forsyte 1.88 SL acts quickly and is rainfast in just 1 hour, giving you a reliable solution to control many tough weeds in soybean crops. Active ingredient fomesafen has flexible uses, controlling over 50 different weed species. Growers across the nation are turning to fomesafen as a reliable and effective weed-resistance strategy.

KEY BENEFITS

- Manages some of the toughest glyphosate resistant weeds
- Fast on-contact activity provides control of over 50 weed species
- Rainfast after 1 hour

KEY USES

- Soybeans

PRODUCT NOTES

EPA REGISTRATION NUMBER

91234-165

ACTIVE INGREDIENT

Fomesafen 22.10%

FORMULATION

Soluble Liquid

HRAC NUMBER

14

SIGNAL WORD

Warning

PACKAGE SIZE

2 x 2.5 gal

265 gal

RESTRICTED USE

No



LABEL



PRODUCT INFO



PORTFOLIO



Bootstrapped and ready to serve, we deliver battle-tested chemistries and an experience like no other. Proud to be 100% American-owned, our mission is to help you every step of the way.





APPLICATION INFORMATION

Pre-plant Surface and Pre-emergence Applications

Certain germinating broadleaf weeds, grasses and sedges can be controlled or partially controlled by soil residual activity from either pre-plant surface or pre-emergence applications of this product. Moisture is necessary to activate this product in soil for residual weed control. Dry weather following applications of this product may reduce effectiveness. When adequate moisture is not received after application of this product, weed control may be improved by overhead irrigation with at least a one-fourth inch of water.

Post-emergence Applications

This product is generally most effective when used post-emergence, working through contact action. Therefore, emerged weeds must have thorough spray coverage for effective control. Best broad-spectrum post-emergence control of susceptible broadleaf weeds is obtained when this product is applied early to actively growing weeds. This usually occurs within 14 to 28 days after planting. Refer to the weed control tables for specific directions on weed growth stages and rates. Some bronzing, crinkling or spotting of labeled crop leaves may occur following post-emergence applications, but labeled crops soon outgrow these effects and develop normally.

KEY WEEDS

Amaranth palmer
Amaranth spiny
Anoda spurred
Balloonvine
Barnyardgrass
Bindweed field
Carpetweed
Cocklebur
Copper-leaf Virginia
Crabgrass
Cucumber Volunteer
Eclipta
Foxtail
Goosegrass
Hemp
Horsenettle
Jimsonweed
Johnsongrass seedling
Ladysthumb
Lambsquarters common
Mallow Venice
Mexicanweed
Milkweed climbing
Milkweed honeyvine
Morningglory
Mustard wild
Nightshade black
Nutsedge yellow
Panicum fall
Panicum Texas
Pigweed
Poinsettia wild
Prickly sida
Purslane common
Ragweed
Rocket yellow
Sesbania hemp
Sicklepod
Signalgrass Broadleaf
Smartweed Pennsylvania
Smellmelon
Sunflower common
Trumpet creeper
Velvetleaf
Waterhemp
Witchweed

(Refer to product label for complete list)

ATTICUSLLC.COM

Refer to product label for complete application and mixing instructions.

©2025 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. Forsyte™ 1.88 SL is not manufactured, or distributed by Syngenta Crop Protection, LLC, seller of Flexstar®. Flexstar® is a registered trademark of a Syngenta Group Company. Forsyte™ is a trademark of Atticus, LLC. 23-AG-0074-B

