



Contains sulfosulfuron, the active ingredient used in Certainty®.

Sertay is a Selective Herbicide for Control of Annual and Perennial Grasses and Broadleaf Weeds in Highly Managed Turf, Ornamental and Native Grass Sites, Landscaped Areas, and Ornamental Nurseries.

ACTIVE INGREDIENT:	(% by weight)
Sulfosulfuron	75.0%
OTHER INGREDIENTS:	<u>25.0%</u>
TOTAL:	100.0%

Sertay is formulated as a water dispersible granule (WDG).

EPA Reg. No.: 91234-120

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See below for additional Precautionary Statements and Directions for Use.

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

For Chemical Emergency:
Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

Sertay™ is not manufactured, or distributed by Valent USA Corporation, seller of Certainty®.



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
KEEP OUT OF REACH OF CHILDREN
CAUTION!

CAUSES MODERATE EYE IRRITATION.

Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants and shoes plus socks and protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.240 (d) (4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Non-Target Organism Advisory

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the **SPRAY DRIFT MANAGEMENT** section of this label.

Windblown Soil Particles Advisory

Sertay has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying **Sertay** if prevailing local conditions may be expected to result in off-site movement.

Groundwater Advisory

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential loading of sulfosulfuron from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow coming into contact with oxidizing agents, hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Read the entire label before using this product.

Use only according to label instructions.

Not all products specified on this label are registered for use in California. Check the registration status of each product in California before using.

Read the **LIMITATION OF WARRANTY AND LIABILITY** before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ATTICUS, LLC DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKAGING.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material, for example Barrier Laminate, Butyl Rubber \geq 14 mils, Nitrile Rubber \geq 14 mils, Neoprene Rubber \geq 14 mils, Natural Rubber \geq 14 mils, Poly-ethylene, Polyvinyl Chloride (PVC) \geq 14 mils, and Viton \geq 14 mils.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (40 CFR Part 170) for agricultural pesticides. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

PRODUCT INFORMATION

Product Description: This product is a postemergence, systemic herbicide with limited soil residual activity. It provides postemergence control of many annual and perennial sedges, grass and broad leaf weeds on highly managed turf, sod farms and native grass sites. It is a selective herbicide that can be used over the top of many perennial warm-season and selected cool-season turfgrasses. Refer to the appropriate sections of this label for approved turf species.

Use Sites: This product may be used for weed control on highly managed turf and native grasses, landscaped areas and ornamental nurseries. This product may be applied to residential and commercial turf sites, including apartment complexes, athletic fields, cemeteries, golf course fairways, golf course roughs, golf course tees, and other golf course areas, hotel properties, nurseries, office complexes, parks, public areas, retail sites, storage facilities, school grounds, sod and turfgrass seed farms, and other highly managed turfgrass sites. This product is not for use on golf course putting greens.

When to Spray: Best results are obtained when target weeds are actively growing and not disturbed by mowing for at least 2 days before and 2 days after application.

Time to Symptoms: This product is absorbed by both the roots and the foliage of plants and rapidly inhibits the growth of susceptible weeds. Susceptible weed growth stops within 24 hours of treatment even though visual symptoms are slow to develop. Susceptible weeds usually show yellowing or browning within 2 to 3 weeks. Warm, moist conditions following application will accelerate herbicidal activity. Cold, dry conditions will delay herbicidal activity. Weeds stressed by drought are less susceptible to this product.

Rainfastness: Heavy rainfall or irrigation within 2 hours after application may wash this product off of the foliage and a repeat application may be required for adequate control.

IMPORTANT: Avoid contact of this product with the roots or foliage of susceptible non-target vegetation as injury may occur. This includes areas where this product may be washed or moved into contact with roots of desirable vegetation.

Susceptible plants may be injured if seeded or transplanted into treated areas unless otherwise directed in this label.

RESISTANCE MANAGEMENT

For resistance management, **Sertay** is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to **Sertay** and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of **Sertay** or other Group 2 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout before and after application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method including hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report lack of performance or suspected resistance, contact Atticus, LLC at (984) 465-4800.



MIXING

Thoroughly clean equipment prior to mixing spray solution.

Eliminate any risk of siphoning the contents of the spray or mixing tank back into the carrier source while mixing. Use approved anti-back-siphoning devices where required by State or local regulations.

Mixing with Water

This product mixes readily with water. Mix spray solutions of this product as follows. Fill the spray tank with about 3/4 of the desired final volume. Add the labeled amount of this product. Continue the filling process while maintaining agitation. Add nonionic surfactant near the end of the filling process.

Mixing for Hand-Held Sprayers

Hand-held sprayer applications must be made at a rate of 2 gallons of spray solution per 1000 square feet.

Using the measuring scoop provided in the product packaging, follow the instructions below to prepare the proper spray solution.

Measuring Scoop Instructions

Using the **SMALL SCOOP** (0.08-gram scoop) provided, refer to the following table for the **Number of Scoops** of product required to achieve the **Desired Application Rate** when mixed in 2 gallons of water.

Desired Application Rate (oz. of product/acre)	Number of Scoops (small scoop)	Mix Volume (gallons of water)	Spray Rate (gallons/1000 ft ²)
0.25 (0.011 lb. AI/A)	2	2.0	2
0.50 (0.023 lb. AI/A)	4	2.0	2
0.75 (0.035 lb. AI/A)	6	2.0	2
1.00 (0.047 lb. AI/A)	8	2.0	2
1.25 (0.058 lb. AI/A)	10	2.0	2

Using the **LARGE SCOOP** (0.80-gram scoop) provided, refer to the following table for the appropriate **Mix Volume** (gallons of water) required to achieve the **Desired Application Rate**.

Desired Application Rate (oz. of product/acre)	Number of Scoops (large scoop)	Mix Volume (gallons of water)	Spray Rate (gallons/1000 ft ²)
0.25 (0.011 lb. AI/A)	1	10.0	2
0.50 (0.023 lb. AI/A)	1	5.0	2
0.75 (0.035 lb. AI/A)	1	3.3	2
1.00 (0.047 lb. AI/A)	1	2.5	2
1.25 (0.058 lb. AI/A)	1	2.0	2
2.00 (0.093 lb. AI/A)	2	2.5	2

Ensure that product is measured as a level scoop and is not rounded.

Add 2 teaspoons (0.33 fl. oz.) of nonionic surfactant per gallon of water.

Tank-Mixing Procedure

For tank mixtures, add individual components to the spray tank in the following sequence: water, water dispersible granules (this product), water-soluble bags, dry flowables, emulsifiable concentrates, drift control additive, water-soluble liquids, non ionic surfactants.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Surfactants and Adjuvants

Use a non ionic surfactant at 0.25 to 0.5 percent by volume (1 to 2 quarts per 100 gallons of spray solution). Use only nonionic surfactants that contain at least 90 percent active ingredient. **DO NOT use nonionic surfactants or other additives that alter the pH of the spray solution below pH 5. Use of surfactants that contain d'Limonene, methylated seed oil, or COC (crop oil concentrate) may cause temporary turf discoloration.**

Colorants and Dyes

Colorants or marking dyes may be added to spray solutions of this product; however, they can reduce product performance. Use colorants and dyes according to the manufacturer's specifications.

APPLICATION EQUIPMENT AND TECHNIQUES

Apply spray solutions of this product with properly maintained and calibrated equipment capable of delivering desired volumes. **DO NOT** apply this product through any type of irrigation system.

GROUND BROADCAST EQUIPMENT

Apply this product uniformly with properly calibrated ground equipment in 10 to 50 gallons of water per acre. Select spray volumes that ensure thorough and uniform weed coverage. Use equipment that is capable of continuous agitation. Choose nozzles that provide optimum spray distribution and uniform coverage at a spray pressure appropriate for applying postemergence herbicides. Avoid streaking, skips, overlaps and spray drift during application.



HAND-HELD SPRAYERS

Backpack sprayers, pump-up pressure sprayers, and other, similar types of hand-held sprayers may be used to apply this product. Apply to foliage of vegetation to be controlled at a rate of 2 gallons of spray solution per 1,000 square feet. Spray coverage should be uniform and complete. Do not spray to the point of runoff.

Equipment Cleaning

Thoroughly clean application equipment with a 1% solution of ammonia (1 quart of ammonia for every 25 gallons of rinse water) promptly after using this product. Use a sufficient volume of cleaning solution to thoroughly rinse all surfaces and to flush all hoses. Rinse with water and repeat the cleaning procedure with the ammonia solution. Complete the cleaning procedure by rinsing thoroughly with clean water. If visible residue is present in the spray tank, use a 1% solution of ammonia plus 0.25% nonionic surfactant (8 fl. oz. per 25 gallons of water) as the cleaning solution.

Restriction:

DO NOT apply by air.

MANDATORY SPRAY DRIFT

Ground Boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy unless making a turf, pasture, or rangeland application, in which case applicators may apply with a nozzle height no more than 4 feet above the ground.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Boom-Less Ground Applications:

- Applicators are required to use a Medium or coarser droplet size (ASABE S572.1) for all applications.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boom-Less Ground Applications:

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

Take precautions to minimize spray drift.



WARM-SEASON TURFGRASSES

This product has been shown to be safe for use on the established warm-season turfgrasses listed in this section.

Bahiagrass	Kikuyugrass
Bermudagrass (common or hybrid)	St. Augustinegrass
Buffalograss	Seashore paspalum
Centipedegrass	Zoysiagrass

Use of this product may result in temporary chlorosis, and may affect the growth pattern or delay green-up of the desirable turf. St. Augustinegrass and seashore paspalum may be more sensitive to this product than other grasses depending on environmental conditions, cultivar differences and other influential factors. For St. Augustinegrass and seashore paspalum, test this product on a small area prior to wide-scale use to determine if this product is suitable for your management and cultural practices.

Sedge Control

For the selective control of the weeds listed in this section, apply this product at 1.25 oz. (0.058 lb ai) per acre after weeds have reached the 3- to 8-leaf stage of growth. A sequential application of 1.25 oz. (0.058 lb ai) per acre may be made 4 or more weeks after the initial treatment, if needed.

Restrictions:

- **Maximum Annual Use Rate:** 2.5 oz. (0.117 lb ai) per acre per year
- **Single Maximum Use Rate:** 1.25 oz. (0.058 lb ai) per acre
- **Maximum Number of Applications/Year:** 2
- **RTI:** 4 weeks

Kyllinga, false green

Kyllinga gracilima

Kyllinga, fragrant

Kyllinga sesquiflorus

Kyllinga, green

Kyllinga brevifolia

Nutsedge, purple

Cyperus rotundus

Nutsedge, yellow

Cyperus esculentus

Sedge, globe

Cyperus croceus

Sedges, annual

Cyperus spp.

Tall Fescue (*Festuca arundinacea*) Control

Best control of tall fescue is obtained when this product is applied at 1.25 oz. (0.058 lb ai) per acre followed by a second application of 1.25 oz. (0.058 lb ai) per acre at 21 to 28 days after the initial application. If a single application is preferred, apply this product at 2.0 oz. (0.093 lb ai) per acre.

Restrictions:

- **Maximum Annual Use Rate:** 2.5 oz. (0.117 lb ai) per acre per year
- **Single Maximum Use Rate:** 2.0 oz. (0.093 lb ai) per acre
- **Maximum Number of Applications/Year:** 2
- **RTI:** 21 days

Dallisgrass (*Paspalum dilatatum*) Suppression in Bermudagrass

The following application includes the use of MSMA. MSMA can cause injury to common and hybrid bermudagrass turf. Test the following tank-mix requirements on a small area prior to wide-scale use to determine if this application is suitable for your bermudagrass management objectives.

For suppression of dallisgrass in bermudagrass turf, apply this product, when dallisgrass is actively growing, at a rate of 1.25 oz. of product (0.058 lb ai) per acre in a tank mixture with 2 lbs. of MSMA per acre and 0.25 percent by volume non-ionic surfactant (1 quart per 100 gallons of spray solution). Reapply this same tank mixture 2 to 4 weeks after initial application. As an alternative program, apply MSMA at 2 lbs. active ingredient per acre with 0.25% by volume non-ionic surfactant as an initial treatment, wait two weeks and apply 2.0 oz. of this product (0.093 lb ai) per acre. Wait an additional two weeks and apply MSMA again at 2 lbs. active ingredient per acre with 0.25% by volume nonionic surfactant.

Virginia Buttonweed (*Diodia virginiana*) Suppression

For suppression of Virginia buttonweed apply this product at 1.25 oz. (0.058 lb ai) per acre. This application will provide suppression or partial control of Virginia buttonweed for 4 to 6 weeks. For enhanced Virginia buttonweed control, tank-mix this product with a broad-leaf herbicide labeled for Virginia buttonweed control in the desired warm-season turfgrass. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Annual Bluegrass (*Poa annua*) Control in Non-Overseeded Turf

For selective control of annual bluegrass (*Poa annua*), apply this product at 1.25 - 2.0 oz. (0.058 - 0.093 lb ai) per acre. Use the higher rate of this product for control in areas of established, dense weed infestation.

Best results are obtained when weeds are in the early stage of growth and prior to tillering.

IN DORMANT BERMUDAGRASS ONLY: tank mixtures of this product with glyphosate-isopropylammonium (Roundup PRO® (EPA Reg. No. 524-529)) or diquat dibromide + glyphosate, ammonium salt (Roundup QuikPRO® (EPA Reg. No. 524-535)) herbicides may be used to increase the spectrum of vegetation controlled. Read and follow the label directions, precautionary statements and all other label information on glyphosate-isopropylammonium (Roundup PRO (EPA Reg. No. 524-529)) or diquat dibromide + glyphosate, ammonium salt [(Roundup QuikPRO (EPA Reg. No. 524-535)) herbicides. Refer to the glyphosate-isopropylammonium (Roundup PRO (EPA Reg. No. 524-529)) or diquat dibromide + glyphosate, ammonium salt (Roundup QuikPRO (EPA Reg. No. 524-535)) product labels for approved application rates. Always apply tank mixtures according to the most restrictive precautionary statements of the products being used.



Annual Bluegrass (*Poa annua*) Control Prior to Overseeding Turf with Perennial Ryegrass

Apply this product at 2.0 oz. (0.093 lb ai) per acre to control *Poa annua* prior to overseeding warm-season turf with perennial ryegrass. Begin applications after *Poa annua* germination and 7 to 10 days prior to overseeding.

Transition of Overseeded Perennial Ryegrass (*Lolium perenne*)

Best results are obtained by applying this product at 1.25 oz. (0.058 lb ai) per acre followed by a second application of 1.25 oz. (0.058 lb ai) per acre at 21 to 28 days after the initial application when daily temperatures are expected to exceed 80° F during the treatment period. If a single application is preferred, apply this product at 2.0 oz. (0.093 lb ai) per acre.

Restrictions:

- **Maximum Annual Use Rate:** 2.5 oz. (0.117 lb ai) per acre per year
- **Single Maximum Use Rate:** 2.0 oz. (0.093 lb ai) per acre
- **Maximum Number of Applications/Year:** 2
- **RTI:** 21 days

Rescuegrass (*Bromus catharticus*) Control

For selective control of rescuegrass apply this product at 0.75 oz. (0.035 lb ai) per acre followed by a second application of 0.75 oz. (0.035 lb ai) per acre at 4 to 10 weeks after the initial treatment. For best results, apply the initial treatment of this product in the fall or early winter when rescuegrass has germinated and is visible in the dormant turfgrass. Applications must be made when rescuegrass is actively growing and at the 2- to 4-leaf stage, but prior to tillering. If a single application is preferred, apply this product at 1.5 oz. (0.07 lb ai) per acre.

Restrictions:

- **Maximum Annual Use Rate:** 1.5 oz. (0.07 lb ai) per acre per year
- **Single Maximum Use Rate:** 1.5 oz. (0.07 lb ai) per acre
- **Maximum Number of Applications/Year:** 2 when using reduced rates
- **RTI:** 4 weeks

Where atrazine can be used in warm-season turfgrass apply this product at 1.0 oz. (0.046 lb ai) per acre plus atrazine at 0.5 lbs. active ingredients per acre. This treatment will provide both postemergence and residual control of rescuegrass.

Additional Weeds Controlled

For selective control or suppression of annual or perennial weeds listed in this section, apply this product at 1.25 - 2.0 oz. (0.058 - 0.093 lb ai) per acre. Use the higher rate of this product for control in areas of established, dense weed infestation. If using an initial rate of application of 1.25 oz. (0.058 lb ai) per acre, a second application of 1.25 oz. (0.058 lb ai) per acre may be made 4 or more weeks after the initial treatment, if needed.

Restrictions:

- **Maximum Annual Use Rate:** 2.5 oz. (0.117 lb ai) per acre per year
- **Single Maximum Use Rate:** 2.0 oz. (0.093 lb ai) per acre
- **Maximum Number of Applications/Year:** 2
- **RTI:** 4 weeks

SPECIMEN

Barley, little*Hordeum pusillum***Bedstraw, catchweed***Galium aparine***Beggarweed, Florida¹****Desmodium tortuosum***Bentgrass, creeping¹****Agrostis stolonifera***Bluegrass, bulbous***Poa bulbosa***Bluegrass, roughstalk***Poa trivialis***Burweed, lawn***Soliva pterosperma***Buttercup***Ranunculus arvensis***Chamber bitter****Phyllanthus urinaria***Chickweed, common***Stellaria media***Clover, white***Trifolium repens***Crowfootgrass***Dactyloctenium aegyptium***Dandelion***Taraxacum officinale***Dichondra, Carolina****Dichondra carolinensis*¹Suppression or partial control only

* NOT REGISTERED FOR USE BY CALIFORNIA

COOL-SEASON TURFGRASSES*

*NOT REGISTERED FOR USE BY CALIFORNIA

Specifications in this section are not intended for use in residential turf.

This product is for use on creeping bentgrass, Kentucky bluegrass and Kentucky bluegrass mixtures with perennial ryegrass and/or fine fescues.

On cool-season turfgrass, apply this product only when turfgrass is actively growing and mowing is required. Applications made in the fall after temperatures have decreased and regular mowing is no longer required may have an increased potential to cause turf injury and may delay spring green-up.

Use of this product may result in temporary chlorosis and discoloration, and may temporarily affect the growth pattern of the desirable turf. These symptoms generally appear 7 to 10 days after application and are typically gone within 21 to 28 days. Turf response is more pronounced under extreme environmental conditions. Perennial ryegrass, fine fescues and creeping bentgrass are more sensitive to this product. If undesirable turf response occurs, skip or delay additional applications to allow turf to recover to a desirable quality. Over-application may result in severe turf injury, thinning or loss of turfgrass stands. Test this product on a small area prior to wide-scale use to determine if this product is suitable for your turf management and cultural practices.

Weed Control in Established Creeping Bentgrass

For use in established creeping bentgrass on golf course fairways, roughs, approaches and tees.

ROUGHSTALK BLUEGRASS (POA TRIVIALIS) CONTROL

For suppression, apply 0.25 oz. of this product (0.011 lb. ai) per acre. For control, up to two additional applications of 0.25 oz. of this product (0.011 lb. ai) per acre at 14- to 21-day intervals may be made. Make all applications between June 1 and September 15.

For severe *Poa trivialis* infestations, apply a single application of 0.75 oz. (0.035 lb. ai) per acre of this product in late summer, 21 to 28 days prior to overseeding creeping bentgrass in a renovation approach. This treatment will result in temporary chlorosis and discoloration, and will temporarily affect the growth pattern of the desirable turf.As *Poa trivialis* is controlled, bare patches may occur in the turf. Creeping bentgrass can be sodded or slit-seeded into the treated area beginning 14 days after the last application of this product in order to minimize regrowth of *Poa trivialis* and to maintain overall turf quality.**Restrictions:**

- **Maximum Annual Use Rate:** 0.75 oz. (0.035 lb. ai) per acre per year
- **Single Max Use Rate:** 0.75 oz. (0.035 lb. ai) per acre
- **Max Number of Applications/Year:** 3 when using reduced rates
- **RTI:** 14 days

WEED CONTROL IN KENTUCKY BLUEGRASS

Weed control specifications in this section are for use ONLY on established pure stands of Kentucky bluegrass.



SEDGE CONTROL

For the selective control of the weeds listed in this section, apply this product at 0.75 oz. (0.035 lb. ai) per acre after weeds have reached the 3- to 8-leaf stage of growth. A sequential application of 0.75 oz. (0.035 lb. ai) per acre may be made 4 or more weeks after the initial treatment, if needed.

Kyllinga, green

Kyllinga brevifolia

Nutsedge, purple

Cyperus rotundus

Nutsedge, yellow

Cyperus esculentus

Sedge, globe

Cyperus croceus

Sedges, annual

Cyperus spp.

Restrictions:

- **Maximum Annual Use Rate:** 1.5 oz. (0.07 lb. ai) per acre per year
- **Single Max Use Rate:** 0.75 oz. (0.035 lb. ai) per acre
- **Max Number of Applications/Year:** 2
- **RTI:** 4 weeks

ROUGHSTALK BLUEGRASS (*POA TRIVIALIS*) CONTROL

Apply this product at 0.5 oz. (0.023 lb. ai) per acre followed by a second application of 0.5 oz. (0.023 lb. ai) per acre at 21 to 28 days after the initial application. If a single application is preferred, apply this product at 0.75 oz. (0.035 lb. ai) per acre.

As *Poa trivialis* is controlled, bare patches may occur in the turf. Kentucky bluegrass can be sodded or slit-seeded into the treated area beginning 14 days after the last application of this product in order to minimize regrowth of *Poa trivialis* and to maintain overall turf quality.

Restrictions:

- **Maximum Annual Use Rate:** 1.0 oz. (0.046 lb. ai) per acre per year
- **Single Max Use Rate:** 0.75 oz. (0.035 lb. ai) per acre
- **Max Number of Applications/Year:** 2 when using reduced rates
- **RTI:** 4 weeks

TALL FESCUE CONTROL

Apply this product at 0.75 oz. (0.035 lb. ai) per acre followed by a second application of 0.75 oz. (0.035 lb. ai) per acre at 21 to 28 days after the initial application.

Restrictions:

- **Maximum Annual Use Rate:** 1.5 oz. (0.07 lb. ai) per acre per year
- **Single Max Use Rate:** 0.75 oz. (0.035 lb. ai) per acre
- **Max Number of Applications/Year:** 2
- **RTI:** 21 days

ADDITIONAL WEEDS CONTROLLED

For selective control or suppression of annual or perennial weeds listed in this section, apply this product at 0.75 oz. (0.035 lb. ai) per acre. A sequential application of 0.75 oz. (0.035 lb. ai) per acre may be made 4 or more weeks after the initial treatment, if needed.

Barley, little*Hordeum pusillum***Bedstraw, catchweed***Galium aparine***Bluegrass, annual***Poa annua***Bluegrass, bulbous***Poa bulbosa***Burweed, lawn***Soliva pterosperma***Buttercup***Ranunculus arvensis***Chamber bitter***Phyllanthus urinaria***Chamomile, mayweed***Anthemus cotula***Chickweed, common***Stellaria media***Clover, white***Trifolium repens***Dandelion***Taraxacum officinale***Garlic, wild¹***Allium vineale***Henbit¹***Lamium amplexicaule***Ivy, ground¹***Glechoma hederacea***Johnsongrass***Sorghum halepense***Mustard, tumble***Sisymbrium altissimum***Mustard, wild***Sinapis arvensis***Pennycress, field***Thlaspi arvense***Quackgrass¹***Elytrigia repens***Shepherd's-purse***Capsella bursa-pastoris***Violet, wild***Viola nephrophylla***Woodsorrell, yellow***Oxalis stricta*¹Suppression or partial control only**Restrictions:**

- **Maximum Annual Use Rate:** 1.5 oz. (0.07 lb. ai) per acre per year
- **Single Max Use Rate:** 0.75 oz. (0.035 lb. ai) per acre
- **Max Number of Applications/Year:** 2
- **RTI:** 4 weeks

WEED CONTROL IN MIXED STANDS OF KENTUCKY BLUEGRASS

This product is specified for use only on established stands of Kentucky bluegrass that include perennial ryegrass and/or fine fescues.

Avoid application of this product on stands with a high percentage of perennial ryegrass or fine fescue.

Perennial ryegrass and fine fescues are more sensitive to this product, therefore, as the percentage of these grasses increase in the mix, occurrence of undesirable turf response will also increase. If undesirable turf response occurs, skip or delay additional applications to allow turf to recover to a desirable quality.

Over-application may result in severe turf injury, thinning or loss of turfgrass stands.

SEDGE SUPPRESSION

For the selective suppression of the weeds listed below apply this product at 0.5 oz. (0.023 lb. ai) per acre after weeds have reached the 3- to 8-leaf stage of growth. A sequential application of 0.5 oz. (0.023 lb. ai) per acre may be made 28 or more days after the initial treatment, if needed.

Alternatively, for suppression of the weeds listed below, apply 0.25 oz. of this product (0.011 lb. ai) per acre. For additional control, up to three additional applications of 0.25 oz. of this product (0.011 lb. ai) per acre at 28-day intervals may be made, if needed.

Nutsedge, purple*Cyperus rotundus***Nutsedge, yellow***Cyperus esculentus***Sedge, globe***Cyperus croceus***Sedges, annual***Cyperus spp.***Restrictions:**

- **Maximum Annual Use Rate:** 1.0 oz. (0.046 lb. ai) per acre per year
- **Single Max Use Rate:** 0.5 oz. (0.023 lb. ai) per acre
- **Max Number of Applications/Year:** 2 [(4 when using reduced rate of 0.25 oz. (0.011 lb. ai) per acre)]
- **RTI:** 28 days

ROUGHSTALK BLUEGRASS (POA TRIVIALIS) CONTROL

For suppression, apply 0.25 oz. of this product (0.011 lb. ai) per acre. For control, up to two additional applications of 0.25 oz. of this product (0.011 lb. ai) per acre at 14- to 21-day intervals may be made. Make all applications between June 1 and September 15.

For severe *Poa trivialis* infestations, apply a single application of 0.5 oz. (0.023 lb. ai) per acre of this product in late summer, 21 to 28 days prior to overseeding turfgrass in a renovation approach.

As *Poa trivialis* is controlled, bare patches may occur in the turf. Kentucky bluegrass, perennial ryegrass or fine fescues can be sodded or slit-seeded into the treated area beginning 14 days after the last application of this product in order to minimize regrowth of *Poa trivialis* and to maintain overall turf quality.

Restrictions:

- **Maximum Annual Use Rate:** 0.75 oz. (0.035 lb. ai) per acre per year
- **Single Max Use Rate:** 0.25 oz. (0.011 lb. ai) per acre
- **Max Number of Applications/Year:** 3
- **RTI:** 14 days

ADDITIONAL WEEDS CONTROLLED

For selective control or suppression of annual or perennial weeds listed in this section, apply this product at 0.5 oz. (0.023 lb. ai) per acre. A sequential application of 0.5 oz. (0.023 lb. ai) per acre may be made 4 or more weeks after the initial treatment, if needed.

[Alternatively, for suppression of the weeds listed in this section, apply 0.25 oz. of this product (0.011 lb. ai) per acre. For additional control, up to three additional applications of 0.25 oz. of this product (0.011 lb. ai) per acre at 28-day intervals may be made, if needed.]

Bedstraw, catchweed¹

Galium aparine

Bluegrass, annual¹

Poa annua

Bluegrass, bulbous¹

Poa bulbosa

Chamber bitter¹

Phyllanthus urinaria

Chamomile, mayweed¹

Anthemis cotula

Chickweed, common

Stellaria media

Clover, white¹

Trifolium repens

Dandelion¹

Taraxacum officinale

Fescue, tall¹

Festuca arundinacea

Garlic, wild¹

Allium vineale

Henbit¹

Lamium amplexicaule

Ivy, ground¹

Glechoma hederacea

Johnsongrass¹

Sorghum halepense

Mustard, tumble

Sisymbrium altissimum

Mustard, wild

Sinapis arvensis

Pennycress, field

Thlaspi arvense

Quackgrass¹

Elytrigia repens

Shepherd's-purse

Capsella bursa-pastoris

Woodsorrell, yellow¹

Oxalis stricta

¹Suppression or partial control only

Restrictions:

- **Maximum Annual Use Rate:** 1.0 oz. (0.046 lb. ai) per acre per year
- **Single Max Use Rate:** 0.5 oz. (0.023 lb. ai) per acre
- **Max Number of Applications/Year:** 2 [(4 when using reduced rate of 0.25 oz. (0.011 lb. ai) per acre)]
- **RTI:** 4 weeks

ORNAMENTALS

This product is for use in woody ornamentals, perennial groundcovers and warm-season ornamental grasses.

For selective control or suppression of weeds listed in this section, apply this product at 1.25 oz. (0.058 lb ai) per acre. A second application of 1.25 oz. (0.058 lb ai) per acre may be made 4 or more weeks after the initial treatment, if needed. Best results are obtained when target weeds are actively growing and not disturbed by mowing for at least 2 days before and 2 days after application.

Restrictions:

- **Maximum Annual Use Rate:** 2.5 oz. (0.117 lb ai) per acre per year
- **Single Maximum Use Rate:** 1.25 oz. (0.058 lb ai) per acre
- **Maximum Number of Applications/Year:** 2
- **RTI:** 4 weeks

Barley, little*Hordeum pusillum***Bedstraw, catchweed***Galium aparine***Beggarweed, Florida^{1*}***Desmodium tortuosum***Bluegrass, annual***Poa annua***Bluegrass, bulbous***Poa bulbosa***Bluegrass, roughstalk***Poa trivialis***Burweed, lawn***Soliva pterosperma***Buttercup***Ranunculus arvensis***Chamber, Bitter****Phyllanthus urinaria***Chickweed, common***Stellaria media***Clover, white***Trifolium repens***Crowfootgrass***Dactyloctenium aegyptium***Dandelion***Taraxacum officinale***Dichondra, Carolina****Dichondra carolinensis***Fescue, tall***Festuca arundinacea***Garlic, wild***Allium vineale***Geranium, Carolina¹***Geranium carolinianum***Henbit¹***Lamium amplexicaule*¹Suppression or partial control only

* NOT REGISTERED FOR USE BY CALIFORNIA

Ivy, ground¹*Glechoma hederacea***Johnsongrass***Sorghum halepense***Kyllinga, false green***Kyllinga gracilima***Kyllinga, fragrant***Kyllinga sesquiflorus***Kyllinga, green***Kyllinga brevifolia***Mustard, wild***Sinapis arvensis***Nutsedge, purple***Cyperus rotundus***Nutsedge, yellow***Cyperus esculentus***Pennycress, field***Thlaspi arvense***Pennywort, lawn¹ (dollarweed)***Hydrocotyle bowlesiioides***Quackgrass¹***Elytrigia repens***Ryegrass, perennial***Lolium perenne***Sedge, globe***Cyperus croceus***Sedges, annual***Cyperus spp.***Shepherd's-purse***Capsella bursa-pastoris***Violet, wild^{1*}***Viola nephrophylla***Woodsorrell, yellow^{1*}***Oxalis stricta***Directed-Spray Applications**

This product may be applied as a post-directed spray around any established warm-season ornamental grass or established woody ornamental species in landscaped areas or field production nurseries. Avoid contact of this product with leaves of desirable plants as foliar injury, discoloration or loss of the plant may result.

Over-the-Top Applications

This product may be applied as an over-the-top application on the ornamental and groundcover species listed below:

American arborvitae

Thuja occidentalis

Azalea, dwarf

Rhododendron atlanticum

Bougainvillea²

Bougainvillea glabra

Boxwood, green velvet

Buxus 'Green Velvet'

Euonymus, wintercreeper

Euonymus fortunei

Gardenia

Gardenia jasminoides

Holly, blue

Ilex x meserveae

Holly, Chinese

Ilex cornuta

Japanese Snowball Bush *

Viburnum plicatum var. tomentosum

Jasmine, Asiatic

Trachelospermum asiaticum

Jasmine, star

Trachelospermum jasminoides

Juniper, Chinese

Juniperus chinensis

Juniper, creeping

Juniperus horizontalis

Juniper, shore

Juniperus conferta

²Single application only

*NOT REGISTERED FOR USE BY CALIFORNIA

Lilac

Syringa vulgaris

Mockorange, Japanese

Pittosporum tobira

Mondo grass

Ophiopogon japonicus

Monkey grass, big blue

Liriope muscari 'Big Blue'

Monkey grass, variegated

Liriope muscari 'Variegata'

Ninebark

Physocarpus opulifolius

Oleander

Nerium oleander

Periwinkle, greater

Vinca major

Photinia, fraser

Photinia x fraseri

Pine, mugo

Pinus mugo

Rhododendron

Rhododendron spp.

Rosemary

Rosmarinus officinalis

Spirea, goldmound

Spirea x 'Goldmound'

Viburnum, Judd *

Viburnum x juddii

Preplant Applications

This product may be applied prior to planting the ornamental species listed below. Wait 14 days after the last application of this product before planting.

Boxwood, green velvet

Buxus 'Green Velvet'

Boxwood, green mountain

Buxus 'Green Mountain'

Burning Bush, dwarf

Euonymus alatus 'Compacta'

Euonymus, wintercreeper

Euonymus fortunei

Forsythia

Forsythia x intermedia

Holly, blue

Ilex x meserveae

Hydrangea, panicled

Hydrangea paniculata

Ivy, English

Hedera helix

Jasmine, winter

Jasminum nudiflorum

Juniper, Chinese

Juniperus chinensis

Juniper, creeping

Juniperus horizontalis

Lilac, dwarf Korean

Syringa meyeri 'Palibin'

Pine, mugo

Pinus mugo

Privet, golden

Ligustrum x vicaryi

Redbud

Cercis canadensis

Rhododendron

Rhododendron spp.

Serviceberry

Amelanchier alnifolia

Viburnum, American cranberrybush

Viburnum trilobum

Viburnum, Prague

Viburnum x pragensis

Weigela

Weigela florida

NATIVE GRASSES

This product generally has been shown to be safe for use on the warm-season native grasses listed in this section.

Big bluestem

Andropogon gerardii

Blue grama

Bouteloua gracilis

Buffalograss

Bouteloua dactyloides

Bushy bluestem

Andropogon glomeratus

Indiangrass

Sorghastrum nutans

Little bluestem

Schizachyrium scoparium

Lovegrass

Eragrostis curvula

Switchgrass

Panicum virgatum

Use of this product may result in temporary chlorosis or temporarily affect the growth pattern of these native grasses. If discoloration or excessive thinning of the native grasses occurs, skip or delay additional applications to allow the native grasses to recover to a desirable quality.

Test this product on a small area prior to wide-scale use to determine if this product is suitable for your management and cultural practices.

Sedge Control

For the selective control of the weeds listed in this section, apply this product at 1.25 oz. (0.058 lb ai) per acre after weeds have reached the 3- to 8-leaf stage of growth. A sequential application of 1.25 oz. (0.058 lb ai) per acre may be made 4 or more weeks after the initial treatment, if needed.

Restrictions:

- **Maximum Annual Use Rate:** 2.5 oz. (0.117 lb ai) per acre per year
- **Single Maximum Use Rate:** 1.25 oz. (0.058 lb ai) per acre
- **Maximum Number of Applications/Year:** 2
- **RTI:** 4 weeks

Kyllinga, false green

Kyllinga gracilima

Kyllinga, fragrant

Kyllinga sesquiflorus

Kyllinga, green

Kyllinga brevifolia

Nutsedge, purple

Cyperus rotundus

Nutsedge, yellow

Cyperus esculentus

Sedge, globe

Cyperus croceus

Sedges, annual

Cyperus spp.

Tall Fescue (*Festuca arundinacea*) Control

Best control of tall fescue is obtained when this product is applied at 1.25 oz. (0.058 lb ai) per acre followed by a second application of 1.25 oz. (0.058 lb ai) per acre at 21 to 28 days after the initial application. If a single application is preferred, apply this product at 2.0 oz. (0.093 lb ai) per acre.

Restrictions:

- **Maximum Annual Use Rate:** 2.5 oz. (0.117 lb ai) per acre per year
- **Single Maximum Use Rate:** 1.25 oz. (0.058 lb ai) per acre
- **Maximum Number of Applications/Year:** 2
- **RTI:** 21 days

Additional Weeds Controlled

For selective control or suppression of annual or perennial weeds listed in this section, apply this product at 1.25 oz. (0.058 lb ai) per acre. A sequential application of 1.25 oz. (0.058 lb ai) per acre may be made 4 or more weeks after the initial treatment, if needed.

Restrictions:

- **Maximum Annual Use Rate:** 2.5 oz. (0.117 lb ai) per acre per year
- **Single Maximum Use Rate:** 1.25 oz. (0.058 lb ai) per acre
- **Maximum Number of Applications/Year:** 2
- **RTI:** 4 weeks

Barley, little*Hordeum pusillum***Bedstraw, catchweed***Galium aparine***Beggarweed, Florida^{1*}***Desmodium tortuosum***Bentgrass, creeping^{*}***Agrostis stolonifera***Bluegrass, annual¹***Poa annua***Bluegrass, bulbous***Poa bulbosa***Bluegrass, roughstalk***Poa trivialis***Burweed, lawn***Soliva pterosperma***Buttercup***Ranunculus arvensis***Buttonweed, Virginia¹***Diodia virginiana***Chamber bitter^{*}***Phyllanthus urinaria***Chickweed, common***Stellaria media***Clover, white***Trifolium repens***Crowfootgrass***Dactyloctenium aegyptium***Dallisgrass¹***Paspalum dilatatum***Dandelion***Taraxacum officinale***Dichondra, Carolina^{*}***Dichondra carolinensis***Garlic, wild***Allium vineale***Geranium, Carolina¹***Geranium carolinianum***Henbit¹***Lamium amplexicaule***Ivy, ground***Glechoma hederacea***Johnsongrass***Sorghum halepense***Mustard, wild***Sinapis arvensis***Pennycress, field***Thlaspi arvense***Quackgrass¹***Elytrigia repens***Ryegrass, perennial***Lolium perenne***Shepherd's-purse***Capsella bursa-pastoris***Violet, wild^{*}***Viola nephrophylla***Woodsorrell, yellow^{*}***Oxalis stricta*¹Suppression or partial control only^{*} NOT REGISTERED FOR USE BY CALIFORNIA

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal. **PESTICIDE STORAGE:** Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area. **PESTICIDE DISPOSAL:** Pesticide spray mixture or rinsate that cannot be used must be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER HANDLING: Bag:** Nonrefillable outer bag. **DO NOT** reuse or refill the outer bag. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **Plastic Container:** Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. All such risks shall be assumed by the user or buyer. **DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Sertay™ is a trademark of Atticus, LLC

Certainty® is a registered trademark of Valent USA Corporation.

Roundup PRO® and QuikPRO® are registered trademarks of Monsanto Technology LLC.

20251017a

