

**HERBICIDE**

CONTAINS HALOSULFURON-METHYL, THE ACTIVE INGREDIENT USED IN SANDEA®

TOUGH ON NUTSEGE, GENTLE ON CROPS

Stadia™ is a selective herbicide that controls nutsedge without the damaging effects that other herbicides leave behind. It provides outstanding pre- and post-emergence control of tough broadleaf weeds that can rob crops of vital nutrients. Stadia offers extended residual control that protects crops from troublesome weeds allowing them to reach maximum yield potential.

Stadia begins working immediately on targeted weeds by stunting growth, ultimately resulting in death of the weeds. Complete control typically occurs within 7 to 14 days depending on the weed size, species and growing conditions. Stadia offers flexible application timing and is an excellent tank-mix partner.

KEY BENEFITS

- Superior control of nutsedge and broadleaf weeds
- Extended residual activity – control broadleaf weeds all season long
- Outstanding crop safety
- Rapid dissolving formulation
- Flexible application timing and tank-mix compatible

KEY USES

- | | |
|---------------|------------------------|
| ▪ Alfalfa | ▪ Pumpkins |
| ▪ Asparagus | ▪ Squash |
| ▪ Cantaloupes | ▪ Succulent Snap Beans |
| ▪ Corn | ▪ Tomatoes |
| ▪ Cucumbers | ▪ Walnuts |
| ▪ Pistachios | ▪ Watermelons |

PRODUCT NOTES

EPA REGISTRATION NUMBER

91234-31

ACTIVE INGREDIENT

Halosulfuron-methyl 75%

FORMULATION

Water dispersible granule

HRAC NUMBER

2

SIGNAL WORD

Caution

PACKAGE SIZE

4 x 80 wt oz

24 x 10 wt oz

RESTRICTED USE

No



LABEL



PRODUCT INFO



PORTFOLIO



Bootstrapped and ready to serve, we deliver battle-tested chemistries and an experience like no other. Proud to be 100% American-owned, our mission is to help you every step of the way.

**Atticus**



APPLICATION INFORMATION

Applications of Stadia™ should be made to actively growing weeds at the early stages of development. Apply by ground to produce uniform coverage on growing weeds or soil to ensure consistent weed control. Atticus recommends using a nonionic surfactant (NIS) spray additive for improved performance.

In post-emergent weed applications, Atticus recommends applying early to control troublesome weeds. If required, apply a second application for long-lasting residual control from early post-emergent treatments to control later germination of weeds.

RECOMMENDATIONS FOR OPTIMAL RESULTS

- Post-emergence applications:
 - Apply Stadia early to small (1 to 3 inches in height) actively growing broadleaf weeds for superior control and to prevent heavy weed infestation.
 - For maximum control of nutsedge it is best to apply Stadia at the 3 to 5 leaf stage.
 - Delay overhead sprinkler irrigation for 2 to 3 days after post-emergence application.
 - Do not apply to weeds that are under drought, stress, disease or insect damage.
- Annual weeds may have multiple flushes of seedlings, or treated perennials may sometimes re-grow from underground stems or roots, depending upon rainfall and other environmental conditions. To maximize control of such weeds, apply a sequential application of this product.
- Use Rate Equivalency:
 - Stadia contains 75% of the active ingredient per pound of product. The following table lists the use rate equivalency in ounces of this product by pounds per acre of active ingredient.

OUNCES OF STADIA	POUNDS OF ACTIVE INGREDIENT PER ACRE
0.5	0.0235
0.67	0.031
1	0.047
1.33	0.062
2	0.094
2.67	0.125
5.33	0.250

KEY WEEDS

Amaranth
Cocklebur
Foxtail
Horseweed/marestail
Kochia
Mustard, wild
Nightshade, black
Nutsedge; yellow and purple
Pigweed; smooth and redroot
Pokeweed, common
Ragweed
Sunflower
Thistle
Velvetleaf

SUPPRESSION

Horsetail
Milkweed; common and honeyvine
Morningglory; ivyleaf and tall

(Refer to product label for complete list)