



## FUNGICIDE



## PREVENTATIVE POWER AND SYSTEMIC CONTROL

The preventative power of Vango™ WG gives protection before disease spreads. Vango WG is a broad-spectrum fungicide which controls certain diseases in almonds, apples, grapes, stone fruits, and other valuable crops. The active ingredient cyprodinil limits the growth of pathogens and inhibits leaf penetration. Growers can rely on Vango WG for lasting protection against a wide range of fungal diseases, including exceptional control against botrytis, several leaf spots, and powdery mildews. It can be applied as a foliar spray and can be used in block, alternating spray, or tank mix programs with other crop protection products.

### KEY BENEFITS

- Offers systemic, long-lasting control of many devastating plant diseases
- Convenient tank mix compatibility
- Preventative activity protects crops before infection

### KEY USES

- Almonds
- Apples
- Grapes
- Pistachios
- Pome Fruit
- Stone Fruit

### PRODUCT NOTES

#### EPA REGISTRATION NUMBER

91234-78

#### ACTIVE INGREDIENT

Cyprodinil 75.0%

#### FORMULATION

Water Dispersible Granule

#### FRAC NUMBER

9

#### SIGNAL WORD

Caution

#### PACKAGE SIZE

4 x 50 wt oz

#### RESTRICTED USE

No



## MIXING INSTRUCTIONS

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area. Buffer the spray solution to a pH of 5.0 - 7.0 when tank mixed with any fungicide containing iprodione.

### Vango WG Alone:

Add 1/2 of the required amount of water to the mix tank. With the agitator running, add the Vango WG to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the Vango WG has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

### Vango WG + Tank Mixtures:

Add 1/2 of the required amount of water to the mix tank. Start the agitator running before adding any tank-mix partners. In general, add tank-mix partners in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables) such as Vango WG, liquid flowables, liquids, and emulsifiable concentrates. Always allow each tank-mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all the mixture has been applied.

When using Vango WG in tank mixtures, add all products in water-soluble packaging to the tank before any other tank-mix partner, including Vango WG. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank-mix partner to the tank.

If using Vango WG in a tank mixture, it is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

*Refer to product label for complete application and mixing instructions.*

©2022 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. Vango™ WG is not manufactured, or distributed by Syngenta Crop Protection, seller of Vanguard®. Vanguard® is a registered trademark of a Syngenta Group Company. Vango™ is a trademark of Atticus, LLC.

## KEY DISEASES

Alternaria  
Alternaria fruit rot  
Anthracnose  
Botrytis  
Botrytis Bunch rot  
Botrytis fruit rot  
Botrytis leaf blight/blast  
Brown rot blossom blight  
Fruit Brown rot  
Green Fruit rot  
Jacket rot  
Neck rot  
Powdery mildew  
Purple blotch  
Scab  
Shothole

*(Refer to product label for complete list)*



[AtticusLLC.com](https://AtticusLLC.com)