

SUPPLEMENTAL LABELING

CLOFENTEZINE	GROUP	10A	INSECTICIDE
--------------	-------	-----	-------------



Supplemental Label for Use on Guava and Hops.

This label expires on May 31, 2022 and must not be distributed or used after that date.

ACTIVE INGREDIENT:	(% by weight)
Clofentezine: 3,6-bis (2-chlorophenyl)-1,2,4,5-tetrazine ⁺	42.0%
OTHER INGREDIENTS:	58.0%
TOTAL	100.0%

⁺Contains 1 pound of active ingredient per quart.

EPA Reg. No.: 91234-109

KEEP OUT OF REACH OF CHILDREN
CAUTION / PRECAUCIÓN

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read the label affixed to the container for **Zorro SC** before applying.

This labeling must be in possession of the user at the time of application.

Use of **Zorro SC** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Zorro SC**.

USE INFORMATION

Zorro SC is a specific miticide, which acts as an ovicide. **Zorro SC** is not disruptive to beneficial insects and mites. It has good residual activity on European red mites, McDaniel mites, Pacific spider mites, two-spotted spider mites, Eriophyes and Yellow spider mites when applied according to label directions.

RATE & TIMING OF APPLICATION

IF THE MITE POPULATION IS APPROACHING DAMAGING LEVELS ON THE TARGET CROPS, COMBINE ZORRO SC WITH AN APPROPRIATE ADULTICIDE

Crop	Rate/Acre/Year	Timing ¹	PHI ²
Guava	4 - 8 oz	First sign of mites	3 days
Hops	4 - 8 oz	First sign of mites	21 days

¹Do not make more than 1 application of **Zorro SC** per year.

²PHI = Pre-Harvest Interval, the **latest** timing for an application prior to harvest.

APPLICATION METHOD

Apply **Zorro SC** by ground using either dilute or concentrate spray equipment. **Do not apply by air.**

Factors which warrant the higher rates (4 to 8 ounces) of **Zorro SC** per acre include:

1. For control of Pacific spider mites.
2. For control of McDaniel spider mites
3. Earliest seasonal timings for blocks with high overwintering European red mite egg populations.
4. When mite predator populations are low.
5. To ensure adequate coverage of full size standard trees and/or trees with heavy leaf densities.

Guava

50 to 400 gallons of water per acre: Apply 4 to 8 ounces of **Zorro SC** per acre. Use sufficient volume of water to obtain complete coverage, but not less than 50 gallons per acre. If the mite population is approaching damaging levels, combine **Zorro SC** with an appropriate, registered adulticide.

Do not make more than one application per year.

Hops

For boom-sprayer applications, apply 4 to 8 ounces of **Zorro SC** with a minimum carrier volume of 20 gallons per acre. For air blast applications, apply with sufficient carrier volume to obtain complete coverage, at least 50 gallons/acre. Do not make more than one application per year.

Do not apply by mechanically pressurized handguns.

Manufactured for:

Atticus, LLC

5000 CentreGreen Way, Suite 100

Cary, NC 27513