



HERBICIDE



EASY, EFFECTIVE SOYBEAN PROTECTION.

Soybean plants and broadleaf weeds are in constant competition for moisture and nutrients. Give your crop the upper hand with FrontRunner™ from Atticus. FrontRunner contains cloransulam, which is the active ingredient found in FirstRate® and is effective on many glyphosate-tolerant weeds including common ragweed, marestalk, lambsquarters and more. Use FrontRunner PPI, preplant surface, preemergence or postemergence for excellent control of broadleaves such as morning glory, sunflower, velvet leaf, and pigweed, all with an excellent crop safety profile with no crop leaf burn or stunting.

FrontRunner is designed for maximum flexibility and convenience. It can be tank mixed to knock down a full range of glyphosate-resistant weeds. FrontRunner can also be safely used in rotation with corn, alfalfa, wheat and other crops.

KEY BENEFITS

- Flexible use; can be applied PPI, preplant surface, preemergence or postemergence
- Can be tank mixed to provide full spectrum control of glyphosate resistant weeds
- Exceptional crop safety with no leaf burn or stunting

KEY USES

- Soybeans

PRODUCT NOTES

EPA REGISTRATION NUMBER
91234-84

ACTIVE INGREDIENT
Cloransulam-methyl 84%

FORMULATION
Water Dispersible Granule

HRAC NUMBER
2

SIGNAL WORD
Caution

PACKAGE SIZE
4 x 0.75 lb

RESTRICTED USE
No

Refer to label for specific use restrictions.



Relevant. Simple. Reliable.



APPLICATION INFORMATION

Apply with ground equipment using a standard low pressure (20 to 40 psi) herbicide sprayer equipped with nozzles that provide uniform coverage. For best results, apply in a spray volume of 10 gallons or more per acre for either soil or postemergence applications. Use sufficient spray volume to provide uniform coverage. Maintain sufficient agitation during mixing and spraying to ensure a uniform spray mixture. Use screens in spray lines and nozzles no finer than 50 mesh (100 mesh is finer than 50 mesh).

RESISTANCE-MANAGEMENT

For resistance management, FrontRunner is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to FrontRunner and other Group 2 herbicides. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of FrontRunner or other Group 2 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures from a different group if such use is permitted; where information on resistance in target weeds species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Scout fields prior to application to identify the weed species present and their growth stage to determine if the intended application will be effective. Scout fields after application to verify that the treatment was effective.

KEY WEEDS

Cocklebur
Common ragweed
Giant ragweed
Jimsonweed
Lambsquarters
Mallow, Venice
Marestail
Morningglory
Palmer amaranth
Pigweed
Sicklepod
Smartweed, Pennsylvania
Sunflower
Velvetleaf
Waterhemp species

(Refer to product label for complete list)

Refer to product label for complete application and mixing instructions.

©2022 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. FrontRunner™ is not manufactured, or distributed by Corteva Agriscience, seller of FirstRate®. FirstRate® is a registered trademark of Corteva Agriscience. FrontRunner™ is trademark of Atticus, LLC.



AtticusLLC.com