

## LASTING WEED CONTROL, EASY-TO-USE LIQUID FORMULATION

Zaltus<sup>TM</sup> SC herbicide protects soybean, cotton, corn, peanut crops and more from a broad-spectrum of annual broadleaf and grass weeds, including Palmer amaranth and waterhemp. The liquid formulation is easy-to-use and stays in suspension for more uniform coverage providing pre-plant burndown weed control and long-lasting residual control for up to six weeks. Zaltus SC may be applied in both your fall and spring burndown programs, making it an ideal product that fits the way you farm. An excellent performer, Zaltus SC protects more than 40 different crops from over 130 different weeds. Achieve weed control with the flexibility and power of Zaltus SC.

### KEY BENEFITS

- Liquid formulation provides quick and easy uniform coverage
- Powerful control on tough weeds
- Flexible application timing
- Up to six weeks of residual activity

### KEY USES

- Almonds
- Corn
- Cotton
- Dry Beans/Peas
- Grapes
- Peanuts
- Soybeans

### PRODUCT NOTES

#### EPA REGISTRATION NUMBER

91234-203

#### ACTIVE INGREDIENT

Flumioxazin 41.4%

#### FORMULATION

Suspension Concentrate

#### HRAC NUMBER

14

#### SIGNAL WORD

Caution

#### PACKAGE SIZE

2 x 2.5 gal

#### RESTRICTED USE

No



LABEL



PRODUCT INFO



PORTFOLIO

## APPLICATION INFORMATION

- Only apply Zaltus SC to healthy crops labeled for post-emergence use. Do not apply Zaltus SC to crops that have been weakened by disease, drought, flooding, excessive fertilization, soil salts, previously applied pesticides, nematodes, insects or winter injury.
- For best results, apply Zaltus SC as part of a burndown program to actively growing weeds.
- Applying Zaltus SC under conditions that do not promote active weed growth will reduce herbicide effectiveness. Do not apply Zaltus SC when weeds are under stress due to drought, excessive water, extremes in temperature, disease or low humidity. Weeds under stress tend to become less susceptible to herbicidal action.
- Zaltus SC is most effective when applied under warm sunny conditions.

## RESISTANCE MANAGEMENT

- Rotate the use of Zaltus SC or other Group 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

## KEY WEEDS

Barnyardgrass  
Bluegrass, annual  
Bristly starbur  
Canola, volunteer  
Carolina geranium  
Chamomile, false  
Cheatgrass  
Chickweed  
Cockle, white  
Coffee senna  
Common ragweed  
Copperleaf, hophornbeam  
Crabgrass  
Dandelion  
Deadnettle, purple  
Downy brome  
Entireleaf  
Eveningprimrose, cutleaf  
Flixweed  
Florida beggarweed  
Foxtail, giant  
Golden crownbeard  
Goosegrass  
Grass weed species  
Groundsel, cressleaf  
Hairy indigo  
Hemp sesbania  
Henbit  
Ivyleaf  
Jimsonweed  
Kochia  
Ladysthumb  
London rocket  
Lovegrass, California  
Mallow, common  
Marestail  
Morningglories  
Mustard  
Palmer amaranth  
Panicums  
Pennsylvania  
Prickly lettuce  
Ragweed, giant  
Russian thistle  
Ryegrass, Italian  
Shepherd's-purse  
Signalgrass, broadleaf  
Smartweeds  
Smellmelon  
Spurred anoda  
Tropic croton  
Velvetleaf  
Waterhemp  
Wild buckwheat  
Wild poinsettia  
Wormwood, biennial  
Yellow rocket

*(Refer to product label for complete list)*