

Contains flumioxazin, the active ingredient used in Clipper® SC Aquatic Herbicide.

HERBICIDE



SCHOONERTM SC

RAPID, CONTACT CONTROL OF TOUGH AQUATIC WEEDS

SchoonerTM SC is a contact herbicide that protects aquatic sites from a wide variety of submersed, emergent and floating weeds. For use in lakes and ponds, Schooner SC works rapidly enough to control tough weeds like watermeal, duckweed and hydrilla in both quiescent and slow-moving water. The active ingredient, flumioxazin, quickly degrades in the water column without building up in the bottom sediment. Schooner SC is an excellent choice for combating actively growing vegetation and is best used when applied to the foliage early in the morning before photosynthesis occurs. The suspension concentrate formulation offers easy mixing and handling, providing less pH sensitivity and long-lasting suspension for up to two hours. Fight aquatic weeds fast with Schooner SC.

KEY BENEFITS

- Quick and selective control of aquatic weeds and algae
- Offers minimal irrigation restrictions
- Suspension concentrate formulation dissolves quickly in the water without building up in the bottom sediment
- Convenient formulation and packaging provides easy mixing and handling

KEY USES

Quiescent or slow-moving bodies of water including:

- Bayous
- Canals
- Drainage ditches
- Lakes
- Marshes
- Ponds (including golf course ponds)
- Reservoirs

PRODUCT NOTES

EPA REGISTRATION NUMBER
91234-204

ACTIVE INGREDIENT
Flumioxazin 41.4%

FORMULATION
Suspension Concentrate

HRAC NUMBER
14

PACKAGE SIZE
24 x 8 fl oz
4 x 1 gal

RESTRICTED USE
No

Refer to label for specific use restrictions.



Relevant. Simple. Reliable.

SURFACE APPLICATION

Apply Schooner SC as a broadcast spray at 6 to 12 fluid ounces of formulated product per acre plus an adjuvant approved for use in aquatics.

SUBSURFACE APPLICATION RATES

WATER DEPTH (FEET)	PINTS OF SCHOONER 4 SC HERBICIDE REQUIRED PER SURFACE ACRE TO ACHIEVE DESIRED WATER CONCENTRATION		
	200 ppb	300 ppb	400 ppb
1	1.1	1.6	2.1
2	2.1	3.2	4.2
3	3.2	4.8	6.4
4	4.2	6.4	8.5
5	5.3	8.0	10.6

Example: to achieve an initial concentration of 200 ppb of flumioxazin in a 4-foot deep water column, apply 4.2 pints (2.1 lbs ai/A) of Schooner SC per surface acre.

USE PRECAUTIONS AND RESTRICTIONS

- Do not apply to intertidal or estuarine areas.
- There is no post-application holding restriction against use of treated water for drinking or recreational purposes (e.g. swimming, fishing).
- In areas with dense weed vegetation only treat 1/2 the water body at one time and wait 10 to 14 days before treating the remaining area. Do not retreat the same section of water within 28 days of application.
- Treated water may not be used for irrigation purposes on food crops until at least five days after application.
- Treated water may be used for irrigation purposes on turf and landscape ornamentals as outlined in the "Irrigation Restrictions Following Application" table.
- Do not use in water utilized for crawfish farming.
- Do not re-treat the same section of water with Schooner SC more than six times per year.
- Do not exceed 400 ppb of Schooner SC during any one application.

IRRIGATION RESTRICTIONS FOLLOWING APPLICATION

APPLICATION METHOD	APPLICATION RATE	AVERAGE WATER DEPTH	TURF AND LANDSCAPE ORNAMENTALS	ORNAMENTALS GROWN FOR PRODUCTION IN GREENHOUSE AND NURSERY
Surface Spray	6 to 12 oz per surface acre	Greater than 3 ft	None	5 days
		Less than 3 ft	12 hours	5 days
Subsurface	Less than 200 ppb	N/A	1 day	5 days
	200 to 300 ppb	N/A	2 days	5 days
	300 to 400 ppb	N/A	3 days	5 days

Refer to product label for complete application and mixing instructions.

©2020 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. Schooner™ SC is not manufactured, or distributed by Nufarm Americas Inc., seller of Clipper® SC Aquatic Herbicide. Clipper® SC Aquatic Herbicide is a registered trademark of Nufarm Americas Inc. Schooner™ SC is a trademark of Atticus, LLC.

KEY WEEDS

Alligator weed
Cabomba
Coontail
Duckweed
Fanwort
Filamentous algae
Frog's-bit
Green algae
Hydrilla
Hygrophila
Mosquito fern
Naiad, southern
Pondweed, curlyleaf
Pondweed, sago
Pondweed, variable-leaf
Water fern
Water lettuce
Water pennywort
Watermeal
Watermilfoil, eurasian
Watermilfoil, variable-leaf

(Refer to product label for complete list)