

CONTAINS DIQUAT DIBROMIDE, THE ACTIVE INGREDIENT USED IN REGLONE®

PRECISE, QUICK DRY DOWN HARVEST CONTROL

Gain control of your harvest with CaponeTM Desiccant, a pre-harvest, contact herbicide used to rapidly control yield robbing vines and weeds in potatoes and more. Once thoroughly covered on actively growing green, Capone Desiccant works by absorbing into plant tissue and destroying plant cells, so the mature crop can dry out from environmental conditions more quickly and evenly. Capone Desiccant also offers growers a more efficient way to harvest through its flexible application methods and its tank-mix compatibility with many other herbicides for increased effectiveness. Take control of your harvest by maximizing yields and minimizing grade loss with Capone Desiccant.

KEY BENEFITS

- Rapid dry down and contact control for quicker desiccation
- Becomes rainfast in just thirty minutes
- Flexible application methods
- Excellent tank-mix partner

KEY USES

- Alfalfa*
- Almonds
- Canola
- Clover*
- Potatoes

*Seed crop only. Refer to label for specific use restrictions.

PRODUCT NOTES

EPA REGISTRATION NUMBER

91234-254

ACTIVE INGREDIENT

Diquat Dibromide 37.3%

FORMULATION

Soluble Liquid

HRAC NUMBER

22

SIGNAL WORD

Caution

PACKAGE SIZE

2 x 2.5 gal
265 gal

RESTRICTED USE

No



LABEL



PRODUCT INFO



PORTFOLIO

APPLICATION TIMING

Capone Desiccant must be applied to emerged weeds when they are small. Weeds 1 inch to 6 inches in height are the easiest to control. When weeds have been grazed or mowed, thus removing much of the green foliage, allow the weeds to regrow to a height of 2–4 inches before spraying. For proper application timing of harvest aid applications, refer to each crop in the product label for specifications or directions. Weeds emerging after application of Capone Desiccant will not be controlled or suppressed.

SPECIFIC USE DIRECTIONS

The following table indicates use pattern, rates, minimum spray volumes, and pre-harvest interval for specific uses.

USE PATTERN	RATE (PT/ACRE)	MINIMUM TOTAL SPRAY (VOLUME/ ACRE)	PRE-HARVEST INTERVAL	DIRECTIONS
POTATOES				
Pre-harvest desiccation broadcast	1–2 pt (0.25 – 0.5 lb ai)	Ground: 20 gal Air: 5 gal	7 days	Restrictions Do not apply through Irrigation systems. Do not graze or feed treated forage to livestock. Do not use seed from treated plants for food, feed, or oil purposes. Do not apply more than 2 pt (0.5 lb ai) per acre per application. Do not apply to drought-stressed potatoes. Do not exceed a total of 4 pt (1 lb ai) per acre. Do not make more than 2 applications per acre per year. Retreatment Interval is 5 days. Precautions Make a second application if necessary to obtain additional desiccation where vine growth is dense. For improved vine coverage, a 5-day interval is advised between applications.

KEY WEEDS

- Algae
- Bladderwort
- Coontail
- Duckweed
- Elodea
- Frog’s Bit
- Hydrilla
- Naiad
- Pennywort
- Pondweeds
- Salvinia
- Waterhyacinth
- Waterlettuce
- Watermilfoils

(Refer to product label for complete list)