

This information is for promotional purposes only. Space considerations may require information to be omitted.
 Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.



Atrevia™

1.2% SL



FOR USE ON GREENHOUSE AND OUTDOOR FOOD CROPS, ORNAMENTAL FLOWERS, TREES, SHRUBS, AND PLANTS

ACTIVE INGREDIENT:	(% by weight)
Azadirachtin.....	1.2%
OTHER INGREDIENTS:	98.8%
TOTAL:	100.0%

Contains 0.0929 lb. (42.2 grams) of azadirachtin per gallon.

EPA Reg. No.: 91234-306

KEEP OUT OF REACH OF CHILDREN

CAUTION

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)**

Read entire label. Use strictly in accordance with precautionary statements and directions for use, and with applicable state and federal regulations.

See below for additional Precautionary Statements.

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> ▪ Call a poison control center or doctor immediately for treatment advice. ▪ Have person sip a glass of water if able to swallow. ▪ Do not induce vomiting unless told to do so by the poison control center or doctor. ▪ Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15-20 minutes. ▪ Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

For Chemical Emergency:
Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before use.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Do not apply when weather conditions favor drift from treated areas. Runoff from treated area may be hazardous to aquatic organisms in neighboring areas.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any irrigation system unless the chemigation instructions on this label are followed. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Socks and shoes

For field sprays:

Keep unprotected persons out of treated areas until sprays have dried.

NON-AGRICULTURAL USE REQUIREMENTS

These requirements apply to uses of this product that are NOT within the WPS for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For other uses including golf courses and other non-agricultural uses, do not enter treated areas without protective clothing until sprays have dried.

PRODUCT MODE OF ACTION

Atrevia 1.2% SL controls target pests on contact or by ingestion. The product acts on pests by way of repellence, anti-feedance and interference with the molting process. The buyer or user is reminded that the degree of efficacy of the product is largely dependent on weather conditions, intensity of pest population, area of application, type of pest, and physical stages of pests and crops.

GENERAL INFORMATION

Read all directions before using this product.

Apply **Atrevia 1.2% SL** as directed to any food or non-food crop up to and including the day of harvest at a rate not exceeding 3½ pints per acre. Refer to the Use Site section for a complete listing of crops.

MIXING

Shake well before using. Add required amount of **Atrevia 1.2% SL** to a clean spray tank with at least one-half of the water to be sprayed. Constant agitation is required, particularly with tank mixes. Agitate the mixture thoroughly and then fill the tank with remaining water and continue agitation. Thorough mixing is necessary for uniform coverage. Non-uniform mixing can cause crop injury or can result in lowered effectiveness. For tank mixes, add other components to the tank containing the **Atrevia 1.2% SL** spray mixture and agitate thoroughly. If tank mixture is allowed to sit, agitation is necessary prior to application. Adjusting the spray mixture pH between 5.5 and 7 will provide optimal performance. Always use this product promptly after mixing with water and do not let tank mix sit for any extended period.

COMPATIBILITY: **Atrevia 1.2% SL** has been found to be compatible with most commonly used pesticides and fertilizers. To avoid problems, conduct a compatibility test before using this product in a tank mix with other pesticides or with fertilizers. To test for compatibility, mix a small amount of each product, in the appropriate proportions, in a small jar test.

A jar test can quickly determine physical compatibility. The process of conducting jar test is given below:

1. Add one pint of water to a glass jar with a lid. (Use the same water source that will go in the tank.)
2. Check spray water pH and adjust if necessary. Often, the pesticide label will give the optimal pH range for best results.
3. Add the pesticides to the jar you plan to use one at a time and shake vigorously after each addition.
4. After all products have been added, shake again, let the solution stand for 15 minutes and then shake one last time and observe the results.

Results: Jar is cool to the touch, and mixture is smooth. Then it is compatible mixture.

PHYTOTOXICITY: **Atrevia 1.2% SL** has been evaluated for phytotoxicity on a wide range of crops and ornamentals. However, since testing on all varieties of all crops and ornamentals is not feasible, test a small portion of the area to be treated for phytotoxicity before treating the entire area. Further, all possible combinations or sequences of pesticide sprays, including other fertilizers, surfactants, adjuvants and other pesticides, have not been tested, thus test for phytotoxicity of spray mixtures. Clean spray equipment used to apply **Atrevia 1.2% SL** thoroughly before use. The addition of spray adjuvants enhances control in some crops under ideal conditions. Addition of certain adjuvants may cause phytotoxicity therefore, test the addition of crop oils and other adjuvants thoroughly tested before using. Do not add crop oils to spray mixtures on ornamental crops. Captan, Bordeaux mixtures, and highly alkaline products cause unacceptable phytotoxicity and/or reduced effectiveness on target pests. Avoid tank mix combinations of **Atrevia 1.2% SL** plus compounds known to be incompatible with oil-based formulations to prevent phytotoxicity. "Waxy bloom" on certain crops and ornamental plants is reduced after a **Atrevia 1.2% SL** application.

APPLICATION INSTRUCTIONS

For optimal performance spray product as soon as possible when pests are expected or when pests first appear. For foliar applications, apply **Atrevia 1.2% SL** in sufficient spray volume and with adequate spray pressure to ensure complete and thorough coverage of all plant surfaces including both the top and bottom of leaves. Avoid excessive runoff. Best results are obtained following 2-3 applications made at 7 - 10 day intervals. When pest pressure is heavy or plant canopy is dense, use higher rates and increase spray frequency. Spraying in the morning or evening hours will provide the best results. Repeat application if rain occurs within two to three hours of spraying.

SPRAY DIRECTIONS

Apply **Atrevia 1.2% SL** as a foliar spray or a drench to soil or non-soil media to control insects. When needed, soil drenches can also be used to control soil-borne pests, including soil-borne larvae of foliar insect pests. When applying as a drench, avoid excessive leaching. **Atrevia 1.2% SL** can also be applied through sub-surface soil treatment equipment. Always follow equipment manufacturers use directions. **Atrevia 1.2% SL** may be applied using any powered or manual pesticide application equipment which includes, but is not restricted to, high volume, low volume, ultra-low volume, electrostatic, fogging and chemigation. Follow the original manufacturer's instructions when using these types of equipment.

DRENCH AND ORNAMENTAL SPRAY DIRECTIONS FOR LABELED PLANTS GROWN IN GREENHOUSES, SHADECLOTHS AND NURSERIES

When used as a soil drench, apply one pint of finished spray for each gallon of soil in the pot. For most pests apply 18-21 oz. **Atrevia 1.2% SL** per 100 gallons of water. For treatment of harder to control pests, such as Dipteran leafminers, use up to 27 ounces per 100 gallons of water. Do not exceed 57 oz of **Atrevia 1.2% SL** per acre per application.

RATES

Use **Atrevia 1.2% SL** at 1-2 pints per acre for most pest and crop conditions. Under extremely heavy pest pressure up to 3½ pints may be used. Do not use less than 5 oz. per acre of **Atrevia 1.2% SL** alone. When tank mixed with other insecticidal products, the rate of **Atrevia 1.2% SL** may be reduced by ½, but not less than 4 oz per acre. Use up to 2.6 oz per 1000 square feet for manure and compost treatments.



CHEMIGATION

General Information

Apply this product through low pressure, drip (trickle) or sprinkler (center pivot, lateral move, end tow, side roll, traveler, big gun, solid set, or hand move) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service Specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Dilute Atruvia 1.2% SL with water before introduction into the system; use the diluted mixture within 8 hours. Do not apply in irrigation water if the pH exceeds 7.0. The optimum pH for application is a range of 5.5 to 6.5. If needed, the pH of the irrigation water can be adjusted by use of a suitable buffering agent. Agitation is necessary. Apply at the rate indicated in the Application Instructions using sufficient water to achieve an even distribution.

Specific Requirements for Chemigation Systems Connected to Public Water Systems -

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Requirements for Sprinkler Chemigation -

1. The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Requirements for Drip (Trickle) Chemigation -

1. The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being filled with a system interlock.



Center pivot, motorized lateral move, or traveling gun types of equipment:

Inject into the system for one revolution or run. Shut off injection equipment after one revolution or run but continue to operate irrigation system until **Atrevia 1.2% SL** has been cleared from the last sprinkler head. Do not use end guns. The system should be run at maximum speed for a foliar application.

Wheel move, side roll, end tow, solid set, or hand move types of equipment:

Adjust equipment to inject **Atrevia 1.2% SL** over a 30–60-minute period. Shut off injection equipment. Continue to operate irrigation system until **Atrevia 1.2% SL** has been cleared from the last sprinkler head. **Atrevia 1.2% SL** can be injected at the end of the irrigation cycle or as a separate application. Do not use end guns. **Atrevia 1.2% SL** must be premixed in a supply tank with water and other appropriate tank-mix chemicals. Agitation is necessary at all times.

Attention must be exercised in irrigation waters with a pH greater than 7. If the irrigation cycle will last longer than 8 hours and the Atrevia 1.2% SL is premixed in the supply tank, the tank mix must be buffered to a pH of 6 or lower. Please contact your Company sales representative should this situation apply.

Application is to be made in sufficient water and of sufficient duration to apply the appropriate rate evenly over the entire treated area.

No field runoff can be permitted during chemigation.

USE SITES:

AGRICULTURAL USE SITES – Use **Atrevia 1.2% SL** on agricultural use sites including, but not limited to, the following:

BERRIES GROUP, such as: Blackberries, Blueberries, Currants, Elderberries, Gooseberries, Huckleberries, Loganberries, Raspberries (red and black). For Strawberries – see miscellaneous.

BULB VEGETABLES, such as: Garlic, Leeks, Onions (dry bulb, green, and Welch), Shallots.

CEREAL GRAINS and GRAINS GROUP, such as: Barley, Buckwheat, Corn, Millet (pearl and Proso), Oats, Popcorn, Rice, Rye, Sorghum (milo), Teosinte, Triticale, Wheat, Wild Rice.

CITRUS FRUITS, such as: Calamondins, Citrus Citrons, Citrus Hybrids, Grapefruits, Kumquats, Lemons, Limes, Mandarins (tangerine), Oranges (sour and sweet), Pummelos, Satsuma Mandarins, White Sapote, UniQ Fruit.

COTTON and TOBACCO

CUCURBIT VEGETABLES, such as: Chayotes, Chinese Waxgourds, Citron Melons, Cucumbers, Gherkins, Gourds (edible), Muskmelons, Pumpkins, Squash (summer and winter), and Watermelons.

FORAGE CROPS, including but not limited to: Alfalfa, Alfalfa Seed, Clover, Trefoil, Vetch.

FRUITING VEGETABLES, such as: Eggplants, Groundcherries, Pepinos, Peppers (including bell pepper, chili pepper, cooking pepper pimento, sweet pepper), Tomatillos, Tomatoes.

HERBS AND SPICES, such as: Allspice, Angelica, Anise (anise seed and star), Annatto (seed), Balm (lemon balm), Basil, Borage, Burnet, Chamomile, Caper Buds, Caraway (black), Cardamom, Cassia bark, Cassia buds, Catnip, Celery Seeds, Chervil (dried), Chives, Chinese Chives, Cinnamon, Clary, Clove buds, Coriander (cilantro or Chinese parsley - leaf), Coriander (cilantro-seed), Costmary, Culantro (leaf and seed), Cumin, Curry (leaf), Dill (dillweed and seed), Fennel (common, Florence), Fenugreek, Grains of Paradise, Horehound, Hyssop, Juniper Berry, Lavender, Lemongrass, Lovage (leaf and seed), Mace, Marigolds, Marjoram, Mustard (seed), Nasturtium, Nutmeg, Parsley (dried), Pennyroyal, Pepper (black and white), Poppy (seed), Rosemary, Rue, Saffron, Sage, Savory (summer and winter), Sweet Bay (bay leaf), Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood.

LEGUME VEGETABLES (Succulent or Dried), such as: Beans, Broad Bean, Chickpeas, Guar, Jackbeans, Lablab Beans, Lentils, Peas, Pigeon Peas, Soybeans, Sword Beans.

LEAFY AND BRASSICA (COLE), such as: Amaranth, Arugula, Broccoli, Broccoli Raab (rapini), Brussels Sprouts, Cabbage, Cauliflower, Cardoon, Cavalo Broccoli (gai ion), Chinese Cabbage (Bok Choy, Napa), Chinese Mustard Cabbage (gai choy), Chinese Celery, Celery, Celtuce, Chervil, Chrysanthemum (edible-leaved, Garland), Collards, Corn Salad, Cress (garden, upland), Dandelion, Dock (sorrel), Endive (escarole), Fennel (Florence), Kale, Kohlrabi, Lettuce (head and leaf), Mizuna, Mustard Greens, Mustard Spinach, Orach, Parsley, Purslane (garden, winter), Radicchio, Rape Greens, Rhubarb, Spinach, Spinach (New Zealand, vine), Swiss Chard, Turnip Greens.

MISCELLANEOUS, such as: Asparagus, Avocado, Banana, Coffee, Cocoa, Cranberry, Figs, Globe Artichokes, Grapes, Hops, Kiwifruit, Mango, Mushroom, Okra, Olives, Papaya, Pawpaw, Peanut, Persimmon, Pineapple, Pomegranate, Strawberry, Tea, Water Chestnut, Watercress, and all other food crops.

POME FRUITS GROUP, such as: Apples, Crabapples, Loquats, Mayhaws, Oriental Pears, Pears, Quinces. (Comice varieties such as Concorde, Seckel, Forelle and Gem): DO NOT apply more than 24 fl oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)

ROOT AND TUBER VEGETABLES GROUP, such as: Arracacha, Arrowroot, Artichokes (Jerusalem, Chinese), Beets (garden, sugar), Burdock, Canna (edible), Carrots, Cassava (bitter and sweet), Celeriac (celery root), Chayote (root), Chervil (turnip-rooted), Chicory, Chufa, Dasheen (taro), Ginger, Ginseng, Horseradish, Leren, Oriental Radish (dailon), Parsley (turnip-rooted), Parsnip, Potatoes, Radishes, Rutabagas, Salsify (oyster plant, black, Spanish), Skirret, Sweet Potatoes, Tanier, Turmeric, Turnips, Yam Bean (jicama, manioc pea), Yams (true).

STONE FRUITS GROUP, such as: Apricots, Cherries (sweet and tart), Nectarines, Peaches, Plums (Chickasaw, Damson, Japanese), Plumcot, Prunes.

TREE AND NUTS GROUP, such as: Almonds, Beechnuts, Brazil Nuts, Butternuts, Cashews, Chestnuts, Chinquapin, Filberts (hazelnut), Hickory Nuts, Macadamias (bush nut), Pecans, Pistachios, Walnuts (black and English).

TROPICAL FRUITS, such as: Acerola, Atemoya, Banana, Biriba, Canistel, Cherimoya, Custard Apple, Feijoa, Guava, Jaboticaba, Ilima, Lychee, Longan, Mango, Papaya, Passionfruit, Pulasan, Rambutan, Sapote (black, mamey), Sapodilla, Soursop, Spanish Lime, Star Apple, Starfruit, Sugar Apple, Wax Jambu.

ORNAMENTAL USE SITES – Use **Atrevia 1.2% SL** on ornamental use sites including, but not limited to, the following:

ORNAMENTAL SHRUBS AND PLANTS, such as: Amaranthus, Aster, Azalea, Ferns, Fuchsia, Caladium, Carnation, Chrysanthemum, Dahlia, Daisy, Lilies, Ivy, Ficus, Gardenia, Impatiens, Iris, Jasmine, Lilac, Marigold, Philodendron, Poinsettia, Rose, Zinnia.

ORNAMENTAL TREES, such as: Ash, Birch, Cedar, Cyprus, Dogwood, Fir, Elm, Juniper, Maple, Oak, Pine, Spruce.

CHRISTMAS TREES AND CHRISTMAS TREE PLANTATIONS

NON-CROP USE SITES – Use **Atrevia 1.2% SL** on non-crop use sites including, but not limited to, the following:

UNCULTIVATED AGRICULTURAL AREAS, such as: farmyards, fuel storage areas, fence rows, rights-of-way, fallow land; soil bank land, barrier strips.

GENERAL SOIL TREATMENTS, such as: Manure, Composts, Cull piles, Mulches, soil application with no mention of crops to be grown (potting soil, topsoil).

PESTS

Atrevia 1.2% SL may be used against the following pests:

Aphids (such as pea aphid, Rosy Apple Aphid), Beetles (such as Japanese beetle), Borers, (such as peachtree borers, peach twig borers), True Bugs, (such as Lygus bugs, stink bugs), Caterpillars, (such as leafrollers, cutworms, loopers, armyworms), Flies (such as walnut husk fly, leafminers and fungus gnats), Leafhoppers, Leafminers, Whiteflies, Mealy Bugs, Mites, Psyllids (such as pear psylla), Weevils, Scales (such as San Jose scale), Thrips, (such as Western flower thrips).

SPECIMEN

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

For plastic containers ≤ 5 gallons: Nonrefillable Container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

For plastic containers > 5 gallons: Nonrefillable container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Atrevia™ 1.2% SL is a trademark of Atticus, LLC.

20220720ap1

