



INSECTICIDE



Atrevia™ 1.2% SL

ORGANIC INSECTICIDE FOR INDOOR & OUTDOOR FOOD CROPS

An effective IPM tool, Atrevia™ 1.2% SL is suitable for application to more than 200 types of indoor and outdoor food crops. Atrevia 1.2% SL is OMRI-listed and contains azadirachtin, a plant-based active ingredient derived from the neem tree. An insect growth regulator, antifeedant, and repellent, Atrevia 1.2% SL works against a wide variety of insects, repelling and controlling target pests on contact or by ingestion. It can be applied right up to the time of harvest and rotated with other synthetic and botanical insecticides. Increase the quality and quantity of crops grown traditionally, organically, and in greenhouses with the versatility of Atrevia 1.2% SL.

KEY BENEFITS

- OMRI-listed for organic production
- Broad-spectrum pest control
- Zero-day pre-harvest interval
- Excellent IPM partner with multiple modes of action
- Easy- to-use mix and spray

KEY USES

Indoor and outdoor food crops including:

- Broccoli
- Grapes
- Lettuce
- Mushrooms
- Oranges
- Pears



PRODUCT NOTES

EPA REGISTRATION NUMBER
91234-306

ACTIVE INGREDIENT
Azadirachtin 1.2%

FORMULATION
Soluble Liquid

SIGNAL WORD
Caution

PACKAGE SIZE
2 x 2.5 gal

RESTRICTED USE
No

Refer to label for specific use restrictions.



Relevant. Simple. Reliable.

APPLICATION INFORMATION

For optimal performance spray product as soon as possible when pests are expected or when pests first appear. For foliar applications, apply Atrevia 1.2% SL in sufficient spray volume and with adequate spray pressure to ensure complete and thorough coverage of all plant surfaces including both the top and bottom of leaves. Avoid excessive runoff. Best results are obtained following 2 - 3 applications made at 7 - 10 day intervals. When pest pressure is heavy or plant canopy is dense, use higher rates and increase spray frequency. Spraying in the morning or evening hours will provide the best results. Repeat application if rain occurs within 2 - 3 hours of spraying.

MIXING

Shake well before using. Add required amount of Atrevia 1.2% SL to a clean spray tank with at least one-half of the water to be sprayed. Constant agitation is required, particularly with tank mixes. Agitate the mixture thoroughly and then fill the tank with remaining water and continue agitation. Thorough mixing is necessary for uniform coverage. Non-uniform mixing can cause crop injury or can result in lowered effectiveness.

For tank mixes, add other components to the tank containing the Atrevia 1.2% SL spray mixture and agitate thoroughly. If tank mixture is allowed to sit, agitation is necessary prior to application. Adjusting the spray mixture pH between 5.5 and 7 will provide optimal performance. Always use this product promptly after mixing with water and do not let tank mix sit for any extended period.

KEY INSECTS

Aphids
Armyworms
Beetles
Caterpillars
Crickets
Cutworms
Flies
Fungus gnats
Grasshoppers
Grubs
Leaf perforators
Leafhoppers
Leafminers
Leafrollers
Locusts
Loopers
Mealy bugs
Mites
Moths
Mushroom flies
Nematodes
Psyllids
Sawflies
Scales
Thrips
True bugs
Webworms
Weevils
Whiteflies

(Refer to product label for complete list)

Refer to product label for complete application and mixing instructions.

©2023 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. Atrevia™ is a trademark of Atticus, LLC. 23-AG-0100-B