



PROTECT CROP YIELDS WITH PYRETHROID- POWERED CHEMISTRY

Cortes™ MAXX Insecticide's chemistry contains the active ingredient zeta-cypermethrin, a synthetic pyrethroid, known for its effectiveness against a wide range of pests and its fast-acting properties. A versatile weapon, Cortes MAXX Insecticide is approved for use on over 200 crops, including corn, soybeans, and alfalfa. It controls over 100 yield-robbing insects, such as aphids, beetles, cutworms, grasshoppers, and weevils, and it can be applied before, during, or after planting.

A potent tool, Cortes MAXX Insecticide works on contact and ingestion, disrupting the nervous system of pests, causing paralysis and death. Its long-lasting effects provide farmers with extended protection, and it is an excellent tank-mix partner. By offering Cortes MAXX Insecticide and its powerful pyrethroid chemistry, you can help farmers tackle infestations, achieve healthier plants, and increase their yields.

KEY BENEFITS

- Quick knockdown pyrethroid formulation
- Controls over 100 pests on 200 crops
- Long-lasting residual control of many pests
- Flexible application timing
- Excellent tank-mix partner

KEY USES

- Alfalfa
- Corn
- Cotton
- Peanuts
- Rice
- Soybeans
- Wheat



PRODUCT NOTES

EPA REGISTRATION NUMBER
91234-273

ACTIVE INGREDIENT
Zeta-Cypermethrin 9.15%

FORMULATION
Emulsifiable Concentrate

IRAC NUMBER
3A

SIGNAL WORD
Warning

PACKAGE SIZE
4 x 1 gal
120 gal

RESTRICTED USE
Yes

Refer to label for specific use restrictions.



APPLICATION INFORMATION

Use lower rates under light to moderate infestation. Use higher rates found in the product label for heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray, and field scouting. Do not exceed maximum labeled rate.

USE GUIDELINES			
CROP	APPLICATION RATE (FL OZ/A)	PHI (DAYS)	MAXIMUM YEARLY TOTAL (FL OZ/A)
Alfalfa grown for seed	2.24 - 4.0	3 (cutting or grazing) 7 (harvesting seed)	24
Field Corn	1.28 - 4.0	7	16
Sweet Corn*	2.24 - 4.0	3	24
Soybeans**	1.28 - 4.0	21	24

*Apply at minimum 3 to 5 day intervals. Apply in water using a minimum of 20 gal per acre of finished spray with ground equipment, and a minimum of 2 gal per acre of finished spray by air.

**Do not make applications less than 7 days apart or within 21 days of harvest.

CHEMIGATION USE DIRECTIONS

Apply Cortes MAXX Insecticide only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply Cortes MAXX Insecticide through any other type of irrigation system. Do not connect any irrigation system (including greenhouse systems) used for pesticide application to a public water system.

Cortes MAXX Insecticide should be applied continuously for the duration of the water application. Cortes MAXX Insecticide should be diluted in sufficient volume to ensure accurate application over the area to be treated. Use the appropriate amount of water to carry the product to the target pest. Agitation is not required when a suitable diluent is used.

Refer to product label for complete application and mixing instructions.

Cortes™ MAXX Insecticide is an EPA Restricted Use Pesticide.

©2024 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. Cortes™ MAXX Insecticide is not manufactured, or distributed by FMC Corporation, seller of Mustang® Maxx. Mustang® is a registered trademark of FMC Corporation. Cortes™ is a trademark of Atticus, LLC. 23-AG-0118-B

KEY INSECTS

Aphids
Armyworm
Bean leaf beetle
Beet armyworm
Beetles
Bollworms
Borers
Caterpillars
Chinch bug
Cloverworms
Colorado potato
Corn silkfly
Cowpea curculio
Cutworms
Earworms
Egyptian alfalfa weevil
European corn
Fall armyworm
Flea beetle
Fleahoppers
Grasshoppers
Green cloverworm
Green peach aphid
Hop vine borer
Hornworm
Imported
Japanese beetle
Kudzu bug
Leaf miner
Leaf skeletonizer
Leafhoppers
Lesser cornstalk
Loopers
Maggots
Meadow spittlebug
Mealy bugs
Mexican bean
Midges
Painted lady
Plant bug
Rootworms
Saltmarsh
Seedcorn maggot
Silver-spotted skipper
Southern corn leaf
Soybean aphid
Spittlebug
Stink bug
Tarnished plant bug
Three-cornered
Thrip
Tobacco budworm
True armyworm
Velvetbean
Webworm
Weevils
Western bean
Whiteflies
Woolly bear
Yellow striped armyworm

(Refer to product label for complete list)



www.AtticusLLC.com