



## END MITES QUICKLY WITH ENDOMITE™

Endomite™ is a powerful miticide that makes an excellent addition to any grower's arsenal of products. Endomite goes to work quickly, knocking out any mite that stands in its way. It controls a wide range of mites and provides effective control at every life stage.

Endomite is highly effective under extreme mite pressure. The unique fuming action of the proven active ingredient, propargite, seeks out mites wherever they hide. Plus, Endomite has the power to penetrate webbing making it an excellent rescue treatment.

Endomite is rainfast and provides extended residual control protecting crops throughout the growing season.

### KEY BENEFITS

- Full-life cycle control on a variety of mites
- Unique fuming effect provides complete coverage
- Immediate, long-lasting control of mites but gentle on beneficial insects
- Excellent rescue treatment – highly effective under high mite pressure

### KEY USES

- Almonds
- Cherries
- Grapes
- Hops
- Walnuts

### PRODUCT NOTES

#### EPA REGISTRATION NUMBER

91234-33

#### ACTIVE INGREDIENT

Propargite 69.6%

#### FORMULATION

Emulsifiable concentrate

#### IRAC NUMBER

12C

#### SIGNAL WORD

Danger

#### PACKAGE SIZE

2 x 2.5 gal

#### RESTRICTED USE

Yes

Refer to label for specific use restrictions.



## APPLICATION INFORMATION

Endomite™ is a liquid emulsifiable concentrate used in sprays for controlling mites. To ensure complete coverage and achieve the best results, applying Endomite to both the upper and lower leaf surfaces as well as fruit is recommended. Endomite is most effective in daytime temperatures averaging above 70°F. For best results apply when mites first appear at the full label rate.

## INSTRUCTIONS AND RATES FOR USE ON BEARING CROPS

| CROP                       | ALMONDS (CALIFORNIA & ARIZONA ONLY)   |   |                 |                             |    |
|----------------------------|---|---|-----------------|-----------------------------|----|
| Mites Controlled           | Clover, European red, Strawberry spider, Two-spotted spider, Pacific spider   |   |                 |                             |    |
| Application Rate (fl oz/A) | 32-64   | Restricted Entry Interval (Days)  | 22 <sup>1</sup> | Pre-Harvest Interval (Days) | 28 |
| Application Instructions   | Apply high rate to areas with high mite infestation and to areas which historically have had high mite infestations.  |   |                 |                             |    |
|                            | Ground Applications   | Apply in a minimum of 50 gallons total volume per acre.   |                 |                             |    |
|                            | Aerial Application  | Apply in a minimum of 15 gallons total volume per acre. One application of Endomite may be applied by air to almonds over one ground application of Endomite or other propargite formulation. |                 |                             |    |
| Restrictions               | <ul style="list-style-type: none"><li>▪ Do NOT make more than 2 applications per season.</li><li>▪ Do NOT apply within 21 days of the last application.</li></ul> |   |                 |                             |    |
| Notes                      | <sup>1</sup> REI Exception: There is no exception allowed for hand weeding or thinning.   |   |                 |                             |    |

| CROP                       | WALNUTS   |  |                 |                             |    |
|----------------------------|---|--|-----------------|-----------------------------|----|
| Mites Controlled           | European red, Two-spotted spider  |  |                 |                             |    |
| Application Rate (fl oz/A) | 32-68   | Restricted Entry Interval (Days)   | 30 <sup>1</sup> | Pre-Harvest Interval (Days) | 21 |
| Application Instructions   | Apply high rate to areas with high mite infestation and to areas which historically have had high mite infestations. Use a minimum of 64 fl. oz. of Endomite per acre on large mature trees.  |  |                 |                             |    |
|                            | Ground Applications   | Apply in a minimum of 100 gallons total volume per acre.   |                 |                             |    |
|                            | Aerial Application  | Apply in a minimum of 20 gallons total volume per acre. Use 48-64 fl. oz. of Endomite when applied by air. |                 |                             |    |
| Restrictions               | <ul style="list-style-type: none"><li>▪ Do NOT apply more than 136 fl. oz. (6.4 lbs. propargite a.i.) of Endomite per acre per year</li><li>▪ Do NOT make more than 2 applications per season.</li><li>▪ Do NOT apply within 21 days of the last application.</li></ul>   |  |                 |                             |    |
| Notes                      | <sup>1</sup> REI Exception: In addition to early entry exceptions allowed by the Worker Protection Standard, you may enter or allow workers to enter treated areas to harvest by tree shaking after 21 days following application, as long as the workers wear long pants, long sleeved shirt and shoes plus socks. There is no exception allowed for hand weeding or thinning. |  |                 |                             |    |

Refer to product label for complete application and mixing instructions.

Endomite™ is an EPA Restricted Use Pesticide.

©2024 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. Endomite™ is not manufactured or distributed by Arysta LifeScience North American, seller of Omite® 6E. Omite® is a registered trademark of MacDermid Agricultural Solutions, Inc. Endomite™ is a trademark of Atticus, LLC. 23-AG-0115-B

## KEY INSECTS

Clover mites  
European red mites  
Pacific spider mites  
Strawberry spider mites  
Two-spotted spider mites

(Refer to product label for complete list)



www.AtticusLLC.com