



FUNGICIDE



CONQUER POWDERY MILDEW & MORE WITH EASE

Rellix™ 40 WSP fungicide is a sterol inhibitor that offers high systemic action, used to control a broad range of fungal diseases in various crops, including powdery mildew, rust, leaf spot, blights, and scab. It is known for its consistent effectiveness in managing powdery mildew across diverse crops such as grapes, apples, cherries, tomatoes, peppers, snap beans, cucurbits, and strawberries, regardless of whether they face high or low disease pressure.

The active ingredient, myclobutanil, works by disrupting the growth and reproduction of fungi, preventing their spread and minimizing additional damage. Rellix 40 WSP fungicide has highly systemic properties that enable it to move rapidly towards new growth, and it does not easily wash away or become diluted when it comes into contact with rainwater or irrigation water, ensuring it remains active and effective for a longer period. Harness the power of Rellix 40 WSP fungicide to ensure a healthier, disease-free crop season.

KEY BENEFITS

- Broad-spectrum disease control for a variety of crops
- Systemic activity for thorough protection
- Very resilient to wash-off for long-lasting control
- Packaged in convenient water-soluble packets

KEY USES

- Apples
- Cherries
- Cucurbits
- Grapes
- Peppers
- Snap Beans
- Strawberries
- Tomatoes

PRODUCT NOTES

EPA REGISTRATION NUMBER

91234-284

ACTIVE INGREDIENT

Myclobutanil 40%

FORMULATION

Wettable Powder

FRAC NUMBER

3

SIGNAL WORD

Caution

PACKAGE SIZE

12 x 5 x 4 wt oz

RESTRICTED USE

No

Refer to label for specific use restrictions.

USE DIRECTIONS FOR TREE FRUITS

Best control is achieved when Rellix 40 WSP is applied on a 7- to 10-day application schedule. Rellix 40 WSP is a systemic fungicide and does not redistribute with rainfall after application. Adjust application equipment spray nozzles to apply a uniform spray throughout the entire tree canopy.

Use the following as guidance in determining the amount of Rellix 40 WSP to be used per 100 gallons of spray or per acre. Refer to specific tree fruit use directions in the product label to determine actual use rates for the control of labeled diseases.

DETERMINATION OF USE RATES ON AN ACRE BASIS

The amount of Rellix 40 WSP required per acre varies with tree size and the volume of fruit and foliage to be treated. Use the following summary table as additional guidance for the determination of appropriate per acre use rates for Rellix 40 WSP:

TREE HEIGHT	APPLE & MAYHAW*	STONE FRUITS
FEET	RATE: FL OZ/ACRE	
10 OR LESS THAN	2.5 - 5	2.5 - 4
15	3.75 - 6	4 - 6
20 OR MORE THAN	5 - 10	6
*For apple scab control, always tank mix Rellix 40 WSP with a multi-site product containing the active ingredient mancozeb.		

KEY DISEASES

Anthracnose
 Asian soybean rust
 Black rot
 Blights
 Blossom blight
 Brown rot
 Brown rot blossom blight
 Cane rust
 Leaf blight
 Leaf spot
 Orange rust
 Pod tip rot
 Powdery mildew
 Rusts
 Scabs
 Shothole
 Yellow rust

(Refer to product label for complete list)

Refer to product label for complete application and mixing instructions.

©2023 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. Rellix™ 40 WSP is not manufactured, or distributed by Corteva Agriscience United States, seller of Rally® 40 WSP Fungicide. Rally® 40 WSP Fungicide is a registered trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow. Rellix™ is a trademark of Atticus, LLC. 23-AG-0133-B