

# TECHNICAL INFORMATION BULLETIN

## RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

ZETA-CYPERMETHRIN | GROUP 3A | INSECTICIDE



EPA Reg. No. 91234-273

### FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

#### FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

This product bulletin contains new or additional directions for use which are recommended by Atticus, LLC under FIFRA Section 2(ee)

### DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for **Cortes™ MAXX Insecticide** before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

CROP	PEST	RATE OF APPLICATION
Fruiting Vegetables (Except Cucurbits) Leafy Vegetables (Except Brassica) Head and Stem Brassica Vegetables Leafy Brassica Greens Legume Vegetables (Except Soybeans)	Symphylans	4.0 fl oz/A (0.025 lb ai/A)

#### DIRECTIONS FOR USE:

Apply **Cortes™ MAXX Insecticide** before, during and after planting for protection from symphylans.

Placement of **Cortes™ MAXX Insecticide** is critical and will depend on seed type and seed placement, or transplant placement.

Treat soil around or in contact with developing plant roots with **Cortes™ MAXX Insecticide**.

Whatever application method is used, it must result in proper placement of **Cortes™ MAXX Insecticide**.

Successful application methods include (but may not be limited to) the following:

#### **Direct-seeded crops:**

- **Cortes™ MAXX Insecticide** should be applied in a narrow band (3-6 inches) and incorporated throughout the seed line.
- Incorporation should result in treated soil around and below (1-3 inches) the seed.
- Mechanical and sprinkler incorporation have been successful, but mechanical incorporation has been more consistent, where more than one seed line per bed are planted, multiple bands or bed top applications will be necessary.

#### **Transplanted Crops:**

- Apply **Cortes™ MAXX Insecticide** to the transplant row prior to transplanting in as narrow a band as practical but not less than 4-6 inches) and mechanically incorporate to a depth of 4-6 inches.
- **Cortes™ MAXX Insecticide** may also be applied as soil drench at the time of transplanting.
- Apply **Cortes™ MAXX Insecticide** to the transplant row with two shanks four inches apart and six inches deep.
- **Cortes™ MAXX Insecticide** may also be applied with a spray blade so a barrier of treated soil immediately below the transplant is obtained.

#### **RESTRICTIONS:**

- Do not apply more than 4.0 fl oz/A (0.025 lb ai/A) per application.
- Do not apply more than 24.0 fl oz/A (0.15 lb ai/A) per season.
- Do not make applications of **Cortes™ MAXX Insecticide** less than 7 days apart on the listed crops except for legume vegetables where applications can be made 5 days apart.
- Do not apply within 1 day of harvest (1-Day PHI) for Fruiting Vegetables, Leafy Vegetables (except Brassica), Head and Stem Brassica Vegetables, Leafy Brassica Greens, and Legume Vegetables (succulent shelled or edible-podded peas or beans).
- Do not apply within 21 days of harvest (21-Day PHI) for Legume Vegetables (dried shelled peas and beans).

Use information in accordance with Section 2(ee) of FIFRA

Manufactured for:  
**Atticus, LLC**  
940 NW Cary Parkway, Suite 200  
Cary, NC 27513

**Cortes™ MAXX Insecticide** is a trademark of Atticus, LLC