

# TECHNICAL INFORMATION BULLETIN

## RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

IMIDACLOPRID	GROUP	4A	INSECTICIDE
BETA-CYFLUTHRIN	GROUP	3A	INSECTICIDE

# *Cryptonyx*™ 360

EPA Reg. No. 91234-292

## FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF  
MI, OR, WA, WI**

### **For Control of Spotted Wing Drosophila on Stone Fruit**

This product bulletin contains new or additional directions for use which are recommended by Atticus, LLC under FIFRA Section 2(ee)

## **DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for **Cryptonyx™ 360** before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

### **STONE FRUIT**

**Crops of Crop Group 12: Apricot, Cherry (sweet and tart), Nectarine, Peach, Plum, (Chickasaw, Damson, and Japanese), Plumcot, Prune (fresh)**

<b>Pest Controlled</b>	<b>Product Rate Fluid Ounces/Acre</b>	<b>Application Method</b>
Spotted Wing Drosophila (adult)	2.4 – 2.8	Apply as a foliar spray, targeting adult growth stage. Reapply as necessary but no closer than a 14 day interval.

### **Restrictions**

Pre-Harvest Interval (PHI): 7 days

Maximum Cryptonyx 360 allowed per 14-day interval: **2.8 fluid ounces/A**

Maximum Cryptonyx 360 allowed per crop season: **5.6 fluid ounces/A**



Manufactured for:  
**Atticus, LLC**

940 NW Cary Parkway, Suite 200, Cary, NC 27513