



HERBICIDE

CONTAINS METSULFURON-METHYL, THE ACTIVE INGREDIENT USED IN ALLY® XP

POST-EMERGENCE CONTROL OF 30+ BROADLEAF WEEDS IN CEREAL CROPS & FALLOW

Azion™ XP post-emergence herbicide is approved for use on wheat, barley, fallow, pasture, and rangeland. With its active ingredient, metsulfuron-methyl, Azion XP effectively targets and controls over 30 key broadleaf weeds, including chickweed, Russian thistle, wild mustard, and smartweed. This herbicide demonstrates its maximum efficacy when applied to young, actively growing weeds, as it is absorbed through their foliage, inhibiting growth, and causing chlorosis in their leaves.

Additionally, Azion XP herbicide can be combined with other approved herbicides, allowing farmers to enhance weed control. It also allows for rotation to sorghum within four to ten months, depending on soil conditions. By harnessing the potent power of Azion XP, farmers can successfully combat vigorous weeds and optimize their crop potential.

KEY BENEFITS

- Delivers superb cereal crop and fallow protection
- Broad-spectrum broadleaf weed control
- Excellent tank-mix partner
- Allows rotation to sorghum in 4-10 months

KEY USES

- | | |
|-----------------------|-------------|
| • Barley | • Sorghum |
| • Fallow | • Triticale |
| • Pasture & Rangeland | • Wheat |

PRODUCT NOTES

EPA REGISTRATION NUMBER

91234-342

ACTIVE INGREDIENT

Metsulfuron-methyl 60.0%

FORMULATION

Water Dispersible Granule

HRAC NUMBER

2

SIGNAL WORD

Caution

PACKAGE SIZE

8 x 8 wt oz

RESTRICTED USE

No



ENGLISH LABEL



SPANISH LABEL



PORTFOLIO



Bootstrapped and ready to serve, we deliver battle-tested chemistries and an experience like no other. Proud to be 100% American-owned, our mission is to help you every step of the way.



USE HIGHLIGHTS

- May be applied by ground or by air.
- Use rates are 1/10 oz per acre in wheat, barley, fallow, or triticale.
- Use rates are 1/10 to 4/10 oz per acre as broadcast treatment in pasture or rangeland. Spot treatments allow up to 3/4 oz per acre.
- No grazing restrictions on wheat, barley, pasture, or rangeland.
- Applied one time per year, Azion XP can be used in wheat and barley as follows:
 - In dryland crops: Apply from 2-leaf stage, but before boot, except on durum and wampum varieties.
 - In durum and wampum varieties: Apply only with 2,4-D at tillering stage but before boot.
 - In irrigated crops: Apply at tillering stage but before boot.
 - Apply as a harvest aid treatment with surfactant (or with 2,4-D + surfactant, or with glyphosate containing herbicides) during dough stages up to 10 days before harvest. This aids in dry down of many broadleaved weeds, thereby aiding grain harvest.
- Apply one time per year to pasture or rangeland for annual weed and selective perennial weed and brush control in several varieties of pasture grasses (also see the section on Application Timing in the product label). For spot applications, use 1 oz per 100 gal of water. Do not exceed 1 2/3 oz of Azion XP per acre.
- Apply as a fallow treatment in the spring or fall when the majority of weeds have emerged and are actively growing. Do not apply during boot or early heading, as crop injury may result.

KEY WEEDS

Annual marshelder
 Black-eyed Susan
 Blue mustard
 Bitter sneezeweed
 Buckbrush
 Bur buttercup
 Burclover
 Carolina geranium
 Coast fiddleneck
 Common; broomweed, chickweed, mullein, purslane, yarrow, groundsel
 Conical catchfly
 Cowcockle
 Curly dock
 Dandelion
 Dogfennel
 False chamomile
 Field pennycress
 Filaree
 Flixweed
 Henbit
 Horsemint
 Kochia
 Marestail
 Mayweed chamomile
 Miners lettuce
 Musk thistle
 Pensacola bahiagrass
 Pigweed
 Plantain
 Plains coreopsis
 Prickly lettuce
 Purple mustard
 Purple scabious
 Russian thistle
 Serecia lespedeza
 Shepherd's purse
 Smartweed
 Snow speedwell
 Small seed false flax
 Tansy mustard
 Treacle mustard
 Tumble hill mustard
 Volunteer sunflower
 Waterpod lambsquarters
 Western snowberry
 Wild; carrot, garlic, mustard
 Woolly croton

(Refer to product label for complete list)