

TECHNICAL INFORMATION BULLETIN

INDOXACARB GROUP 22A INSECTICIDE



EPA Reg. No. 91234-318

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ARIZONA AND CALIFORNIA

For the Suppression of Leaf Footed Plant Bugs, Green Stink Bugs, *Carpophilus truncatus*, and Feeding Damage Protection in Nutmeat in Tree Nuts

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for **Armort™ 30 DG** before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Crops	Pest	Rate per Acre	Application Directions	REI
Tree Nuts; Group 14-12 almond; pistachio; walnut, black; walnut, English; cultivars, varieties, and/or hybrids of these	Leaf Footed Plant bugs, Green Stink Bugs, and <i>Carpophilus truncatus</i>	6 oz. (0.11 lb. a.i.)	Apply as required by Scouting, usually from late dormancy until Mid-June Timing and frequency of applications should be based on insect population and threshold.	12 hrs

Use Restrictions:

- DO NOT apply more than 3 applications of **Armort 30 DG** per crop season.
- DO NOT apply more than 18 oz/A of **Armort 30 DG** or 0.33 lb ai/A of indoxacarb-containing products per calendar year.
- The minimum interval between treatments is 7 days.

NOTE: Spray Volume: Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of trees or plants and density of foliage.

Ground Application: Apply in a minimum of 50 gal/A of water by conventional ground equipment. For best results, apply 100-150 gal/A of water.

Aerial Application: Apply in a minimum of 10 gal/A of water. Applying by air can result in reduced canopy coverage, which may also reduce efficacy. For best results, use an adjuvant to help increase coverage, penetration and thus performance.

Advisory: This product is highly toxic to bees and other pollinating insects exposed to direct treatment or to residue in/on blooming crops or weeds. DO NOT apply this product or allow it to drift to blooming crops or weeds while bees are foraging in the treatment area.

Recommendations:

- DO NOT apply **Armort 30 DG** later than June 15 due to MRLs constrain with EU. It is the responsibility of the grower and processor to check for residue before shipping overseas.



Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513