

ReCon™ 4 F contains the active ingredient metalaxyl and uses the same mode of action as Ridomil®, which contains mefenoxam.



## FUNGICIDE



## DEPENDABLE TURF & ORNAMENTAL PROTECTION FROM SOIL-BORNE DISEASE

ReCon™ 4 F, a systemic fungicide powered by the active ingredient metalaxyl, effectively controls soil-borne diseases in both newly seeded and established turf, as well as ornamentals. This powerful tool targets diseases caused by members of the Oomycete class of fungi, including *Pythium* and *Phytophthora* spp. ReCon 4 F protects plants by preventing pathogens from infecting roots, stems, and leaves.

With the systemic activity of ReCon 4 F, both turf and ornamentals benefit from root-to-leaf protection, promoting robust and healthy growth. Its low-odor liquid formulation easily integrates into most fungicide and fertilizer tank-mixes. Elevate your plant's health with ReCon 4 F – a key tool for proactive disease management, ensuring robust growth and vitality.

### KEY BENEFITS

- Robust preventative protection against Oomycete diseases
- Rapid uptake and translocation
- Excellent tank-mix partner
- Approved for multiple uses and many plant species

### KEY USES

- Golf Courses
- Greenhouse & Nurseries
- Landscapes
- Lawns
- Ornamentals
- Sod Farms



### PRODUCT NOTES

**EPA REGISTRATION NUMBER**  
91234-64

**ACTIVE INGREDIENT**  
Metalaxyl 44.08%

**FORMULATION**  
Suspension Concentrate

**FRAC NUMBER**  
4

**SIGNAL WORD**  
Caution

**PACKAGE SIZE**  
4 x 1 gal

**RESTRICTED USE**  
No

Refer to label for specific use restrictions.



Relevant. Simple. Reliable.



## TURF APPLICATION INFORMATION

For established turf, apply as a preventative treatment at 0.5 to 1 fl. oz. of ReCon 4 F in 3 to 5 gallons of water per 1,000 sq. ft. Retreat at 10 to 21 day intervals. During periods of prolonged conditions favorable for disease development, use 1 fl. oz. of ReCon 4 F on a 14 day schedule.

For newly seeded areas, apply 0.5 to 1.0 fl. oz. product in 5 to 10 gallons of water per 1,000 sq. ft. immediately after seeding. Retreat at 7 to 14 day intervals if conditions remain favorable for disease. **Note:** For long-term control of Pythium in areas when using seed treated with the active ingredient contained in ReCon 4 F, make application of ReCon 4 F 7 to 10 days after seeding.

## RESISTANCE MANAGEMENT

To minimize the potential for resistance.

1. Make no more than three applications of ReCon 4 F per season.
2. Apply an alternate EPA-registered fungicide for Pythium control at least once during the season.

## ORNAMENTALS APPLICATION INFORMATION

Use ReCon 4 F on container, bench, or bed-grown ornamentals in greenhouses or outdoor nurseries, and for use on ornamentals grown for indoor and outdoor landscaping, for control of damping-off, and root and stem rot diseases caused by Pythium and Phytophthora. ReCon 4 F may be applied through irrigation systems as a soil drench, as a soil surface spray, or incorporated into a soil mix for subsequent seeding or transplanting of ornamentals. ReCon 4 F may be applied as a foliar spray on azaleas.

Within a rate range given for a specific group of ornamentals, use the lower specified rate for the shortest interval listed and the higher specified rate for the longest interval. For drench applications, use enough of the specified ReCon 4 F water solution to wet the root zone of plants. In general, 1 pt. product per sq. ft. of this solution is sufficient for ornamentals growing in containers with 4 inches of growth media. Containers with growth media depth greater than 4 inches generally require 1 1/2 to 2 pts. product per sq. ft. of the solution. If soil surface applications are made, irrigate with at least 1/2 inch of water if rainfall does not occur within 7 days.

*Refer to product label for complete application and mixing instructions.*

©2024 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. ReCon™ 4 F is not manufactured or distributed by Syngenta Group Company, seller of Ridomil®. Ridomil® is a registered trademark of Syngenta Group Company. ReCon™ is a trademark of Atticus, LLC. 24-EC-0002-A

## KEY DISEASES

Downy mildew  
Phytophthora root and crown rot  
Phytophthora shoot blight  
Pythium blight  
Pythium damping-off  
Yellow tuft  
Yellow turf downy mildew

*(Refer to product label for complete list)*



[www.AtticusLLC.com](http://www.AtticusLLC.com)