

# BESTERO™ 50 WG



Contains Flonicamid, the active ingredient used in Beleaf® 50SG.

<b>ACTIVE INGREDIENT:</b>	(% by weight)
*Flonicamid. ....	50.0%
<b>OTHER INGREDIENTS:</b> .....	50.0%
<b>TOTAL:</b> .....	100.0%
* Contains 0.5 pound of active ingredient per pound of formulated product.	
EPA Reg. No.: 91234-300	

## KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

See below for additional Precautionary Statements.

FIRST AID	
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>▪ Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li><li>▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>▪ Call a poison control center or doctor for treatment advice.</li></ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"><li>▪ Immediately call a poison control center or doctor for treatment advice.</li><li>▪ Have person sip a glass of water if able to swallow.</li><li>▪ Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>▪ Do not give anything by mouth to an unconscious person.</li></ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"><li>▪ Take off contaminated clothing.</li><li>▪ Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>▪ Call a poison control center or doctor for treatment advice.</li></ul>
<b>Note to Physician:</b> Treatment is otherwise controlled by removal of exposure followed by symptomatic and supportive care.	
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

**For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident,  
Call CHEMTREC Day or Night Within USA and Canada:  
1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

Bestero™ 50 WG is not manufactured, or distributed by FMC Corporation, seller of Beleaf® 50SG.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (& DOMESTIC ANIMALS) WARNING

Causes substantial but temporary eye injury. Do not get in eyes, on skin or on clothing. Harmful if swallowed or absorbed through the skin. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants;
- Shoes plus socks;
- Protective eye wear (goggles, face shield or safety glasses); and
- Chemical-resistant and/or waterproof gloves.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

**Users should:**

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

**Pollinator Advisory:** This product is moderately toxic to bees and other pollinating insects exposed to direct treatment, or to residues in/on blooming crops or weeds. Protect pollinating insects by following label directions intended to minimize drift and to reduce risk to these organisms.

### PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow coming in contact with oxidizing agent. Hazardous chemical reaction may occur.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### Use Restrictions

- **DO NOT** apply this product in a way that it will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- **DO NOT** store **Bestero 50 WG** spray mixtures overnight.
- **DO NOT** use this product in home gardens.
- **DO NOT** use liquid fertilizer as a carrier for **Bestero 50 WG**.
- **DO NOT APPLY BESTERO 50 WG THROUGH ANY IRRIGATION SYSTEM PHYSICALLY CONNECTED TO A PUBLIC WATER SYSTEM.**
- For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval (REI) and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the **Restricted-Entry Interval (REI)** of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeve shirt;
- Shoes plus socks;
- Protective eye wear (goggles, face shield or safety glasses); and
- Chemical-resistant and/or waterproof gloves.

### PRODUCT INFORMATION

**Bestero 50 WG** is a 50 percent soluble granular formulation of the insecticide flonicamid. It controls insects which may vector viruses and pathogens. Residual control of flonicamid has been shown to reduce disease transmission and symptoms, thereby increasing plant health. **Bestero 50 WG** provides control of a variety of aphid and plant bug pests and suppression of some non-aphid pests in Brassica head and stem vegetables; Brassica leafy greens; cucurbit vegetables; fruiting vegetables (except cucurbits); hop; leaf petiole vegetables; leafy greens; tuberous and corm vegetables; legume vegetables (except soybeans); root vegetables (except sugar beets); pome fruit; stone fruit; tree nuts; citrus; commercial production of greenhouse-grown cucumbers, lettuce, peppers and tomatoes; grapes and other small vine climbing fruit (except fuzzy kiwifruit); low growing berries; canola; alfalfa (forage, hay, and seed); clover (forage and hay) (West of the Rockies); celtuce; Florence fennel; kohlrabi and mint.

The rate of application is dependent upon the insect species present, the level of insect pressure, and the amount of foliage present. Begin applications before populations begin to build or at economic thresholds according to local economic guidelines. Refer to local Cooperative Extension Guideline and/or time applications for scouting results. Thorough plant coverage is essential for good performance.

**Mode of Action:** **Bestero 50 WG** is a member of the pyridinecarboxamide class of chemistry. **Bestero 50 WG** controls target pests by contact and ingestion provoking rapid and irreversible feeding cessation. Aphids and other insects could remain on the plant until they desiccate.

### Pesticide Resistance Management:

For resistance-management, **Bestero 50 WG** contains a Group 29 insecticide. Any insect population may contain individuals naturally resistant to **Bestero 50 WG** and other Group 29 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed. To reduce the potential for developing insect resistance, rotate to an insecticide with a different mode of action. Monitor treated pest populations for resistance development. Read product label before applying any insecticide and follow label directions.

To delay insecticide resistance, take the following steps:

- Rotate the use of **Bestero 50 WG** or other Group 29 insecticides within a growing season, or among growing seasons, with different groups that control the same pests. Avoid application of more than the maximum seasonal use rate or the total number of consecutive sprays of **Bestero 50 WG** per season.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
  - o Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
  - o Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
  - o When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
  - o Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
  - o The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Atticus LLC at 984-465-4800.

### Crop Rotation Restrictions

Following application of **Bestero 50 WG**, any crop listed on this label may be planted as a rotational crop.

All other crops may be planted 30 days after the last application of **Bestero 50 WG**.

### Maximum Seasonal Application Rates and Pre-Harvest Intervals

Crop	Maximum Seasonal Total (Pounds Active Ingredient)	PHI Days
Listed Brassica Head and Stem Vegetables	0.263	0
Listed Brassica Leafy Greens	0.263	0
Listed Cucurbit Vegetables	0.263	0
Listed Fruiting Vegetables (Except Cucurbits)	0.267	0

(continued)

### Maximum Seasonal Application Rates and Pre-Harvest Intervals (continued)

Crop	Maximum Seasonal Total (Pounds Active Ingredient)	PHI Days
Grapes And Other Small Vine Climbing Fruit (Except Fuzzy Kiwifruit)	0.263	7
Hops	0.263	10
Listed Leaf Petiole Vegetables	0.263	0
Listed Leafy Greens	0.263	0
Listed Tuberous and Corm Vegetables	0.263	7
Listed Legume Vegetables (succulent and dried) Except Soybeans	0.263	7
Listed Root Vegetables (Except Sugar Beets)	0.263	3
Listed Pome Fruit	0.263	21
Listed Stone Fruit	0.263	14
Listed Tree Nuts	0.263	40
Listed Citrus Fruit	0.263	0
Greenhouse Cucumbers	0.267	0
Greenhouse Tomatoes	0.267	0
Greenhouse Peppers/Eggplants	0.267	0
Greenhouse Lettuce	0.088	0
Rapeseed	0.263	7
Listed Sunflower Commodities	0.263	0
Low Growing Berries	0.263	0
Strawberry	0.263	0
Kohlrabi	0.263	0
Celtuce	0.263	0
Florence Fennel	0.263	0
Mint (Spearmint and Peppermint)	0.263	7
Alfalfa (Forage, Hay, and Seed)	0.18	14
Clover (Forage and Hay) (Grown West of the Rockies)	0.18	60

### Mixing and Loading Instructions

The spray system must be clean and free of residues from previous applications. Fill the spray tank 1/2 full with clean water. The agitation system must be operating and sufficient to provide uniform spray mixing during application and until the spray tank has been emptied. Complete filling the spray tank to the desired level.

## Tank Mixtures

**Bestero 50 WG** can be mixed with products labeled for use on the crops/sites listed on this label in accordance with the more (most) restrictive of label limitations and precautions. No label dosage rates may be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Bestero 50 WG** is generally compatible with other insecticides, fungicides, fertilizers and micronutrient products provided sufficient free water is available for dispersion of all the tank mix products. However, the physical compatibility of **Bestero 50 WG** with tank mix partners should be evaluated using a jar test before use.

The crop safety of all potential tank mixtures on all crops may not have been tested. Before applying any tank mixture not specifically recommended on this label, the safety to the target crop must be confirmed.

In general, tank mix partners should be added in the following order: products in water-soluble packaging, wettable powders or wettable granules or dry flowables, liquid flowables, liquids, then emulsifiable concentrates. Allow each tank mix partner to become completely dispersed before adding the next product.

## Spray Equipment Clean Out:

After spraying **Bestero 50 WG**, thoroughly clean the sprayer before using sprayer equipment for any other applications. In addition, users must take appropriate steps to ensure proper equipment clean out for any other products mixed with **Bestero 50 WG** as required on the other product labels. Refer to the **STORAGE AND DISPOSAL** and **ENVIRONMENTAL HAZARDS** statements regarding disposal of equipment washwaters.

## APPLICATION INFORMATION

### Mandatory Spray Drift Controls

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers. Use the largest droplet size consistent with good pest control.

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.

### Ground Boom Applications

Utilize a boom and nozzle sprayer equipped with the appropriate nozzles, spray tips and screens and adjusted to provide optimum spray distribution and coverage at the appropriate operating pressures. Utilize nozzles that produce minimal amounts of fine spray droplets. Do not exceed 30 psi spray pressure unless otherwise required by the manufacturer of drift reducing nozzles. Sprayers should be adjusted to position spray tips a minimum of 18 inches above the crop. Be aware that overlaps and slower ground speeds while starting, stopping or turning while spraying may result in higher application rates.

### Aerial Applications

Utilize a boom and nozzle sprayer equipped with the appropriate nozzles, spray tips and screens and adjusted to provide optimum spray distribution and coverage at the appropriate operating pressures. Utilize nozzles that produce minimal amounts of fine spray droplets. Do not exceed 30 psi spray pressure unless otherwise required by the manufacturer of drift reducing nozzles. Use nozzle types and arrangements that will provide optimum coverage while producing a minimal amount of fine droplets.

## Chemigation Applications

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, drip (trickle) (greenhouse cucumbers, peppers and tomatoes as well as field-grown cucurbit vegetables (crop group 9) and fruiting vegetables (crop group 8-10) only) or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect any irrigation system, including greenhouse systems, used for pesticide application to a public water system. Crop injury, lack of effectiveness or illegal residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must also contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment. **Bestero 50 WG** should be applied continuously for the duration of the water application. **Bestero 50 WG** should be diluted in sufficient volume to ensure accurate application over the area to be treated. Use the appropriate amount of water to carry the product to the target pest. Agitation generally is not required when suitable diluents are used. A diluents test should be conducted to ensure that phase separation would not occur during dilution and application. Failure to achieve a uniform dilution throughout the time of application may result in undesirable residues or less than desirable control.

**Using Water from Public Water Systems: DO NOT APPLY BESTERO 50 WG THROUGH ANY IRRIGATION SYSTEM PHYSICALLY CONNECTED TO A PUBLIC WATER SYSTEM.** Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. **Bestero 50 WG** may be applied through irrigation systems, which may be supplied by a public water system **only if** water from the water system is discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. Before beginning chemigation, always make sure that the air gap exists and that there is no blockage of the overflow of the reservoir tank.

## SPRAY DRIFT MANAGEMENT

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.



The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. Where states have more stringent regulations, they must be observed.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

1. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.
2. Do not cultivate within 10 feet of the aquatic area so as to allow growth of a vegetative filter strip.
3. Do not apply by ground equipment within 25 feet, or by air within 150 feet of lakes; reservoirs, rivers, permanent streams, marshes or natural ponds, estuarine/marine habitats, and commercial fish farm ponds. Increase the buffer zone to 450 feet when ultra low volume (ULV) applications are made.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisories below.

### SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly, or under unfavorable environmental conditions.

#### Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

#### Controlling Droplet Size - Aircraft

- Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

### BOOM HEIGHT - Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

### RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

### SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

### WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

### INSECT SPECTRA

The following tables list various insect species, which are either effectively controlled or suppressed<sup>1</sup> when appropriate application rates of **Bestero 50 WG** are made at appropriate timings. See specific crop sections for recommendations on specific pests.

<sup>1</sup>Suppression may be erratic control ranging from good to poor, or a consistent level of control below that generally considered commercially acceptable.

#### Aphid Pests

Common Name	Scientific Name
Apple Aphid	<i>Aphis pomi</i>
Black Bean Aphid	<i>Aphis fabae</i>
Black Cherry Aphid	<i>Myzus cerasi</i>
Cabbage Aphid	<i>Brevicoryne brassicae</i>
Cotton/Melon Aphid	<i>Aphis gossypii</i>
Cowpea Aphid	<i>Aphis craccivora</i>
English Grain Aphid	<i>Sitobion avenae</i>
Green Peach Aphid	<i>Myzus persicae</i>
Greenbug	<i>Schizaphis graminum</i>
Hop Aphid	<i>Phorodon humuli</i>
Leaf Curl Plum Aphid	<i>Brachycaudis helichrysi</i>
Mealy Plum Aphid	<i>Hyalopterus pruni</i>
Foxglove Aphid	<i>Aulacorthum solani</i>
Pea Aphid	<i>Acyrtosiphon pisum</i>
Potato Aphid	<i>Macrosiphum euphorbiae</i>
Red Lettuce Aphid	<i>Uroleucon pseudambrosiae</i>
Rosy Apple Aphid	<i>Dysaphis plantaginea</i>
Spirea Aphid	<i>Aphis spiraeicola</i>
Turnip Aphid	<i>Lipaphis erysimi</i>
Woolly Apple Aphid	<i>Eriosoma lanigerum</i>

(continued)

### Aphid Pests

Common Name	Scientific Name
Red Lettuce Aphid	<i>Nasonovia ribis-nigri</i>
Spotted Alfalfa Aphid	<i>Therioaphis maculata</i>
Blue Alfalfa Aphid	<i>Acyrtosiphon kondoi</i>
Strawberry aphid	<i>Chaetosiphon fragaefolii</i>
Soybean Aphid	<i>Aphis glycines</i>
Black Pecan Aphid	<i>Melanocallis caryaefoliae</i>
Yellow Pecan Aphid	<i>Monelliopsis pecanis</i>
Blackmargined Aphid	<i>Monellia caryella</i>

### Non-Aphid Insect Pests

Common Name	Scientific Name
Cotton Fleahopper	<i>Pseudatomoscelis seriatus</i>
Greenhouse Whitefly	<i>Trialeurodes vaporariorum</i>
Tarnished Plant Bug	<i>Lygus lineolaris</i>
Western Plant Bug	<i>Lygus hesperus</i>
Potato Psyllid	<i>Bactericera cockerelli</i>
Tomato Psyllid	<i>Bactericera cockerelli</i>
Asian Citrus Psyllid	<i>Diaphorina citri</i>
Vine Mealybug	<i>Planococcus ficus</i>

### Rate Conversion Chart for Bestero 50 WG

Ounces Bestero 50 WG / A	Pounds Bestero 50 WG / A	Pounds AI / A	Treated Acres / Pound Bestero 50 WG
1.4	0.088	0.044	11.4
1.7	0.106	0.053	9.4
2.0	0.125	0.062	8.0
2.4	0.15	0.075	6.7
2.8	0.175	0.088	5.7

**VEGETABLE, BRASSICA, HEAD AND STEM, GROUP 5-16 AND KOHLRABI (0 DAY PHI) INCLUDING:**

Broccoli; Brussels sprouts; cabbage; cabbage, Chinese, napa; cauliflower; kohlrabi; cultivars, varieties, and hybrids of these commodities

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids and Plant Bugs	2.0 to 2.8	0.062 to 0.088	<ul style="list-style-type: none"> <li>▪ Begin applications before populations begin to build and before damage is evident according to local pest management guidelines.</li> <li>▪ Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage.</li> <li>▪ Rapidly growing plants may need retreatment. Scout fields often and retreat as necessary to maintain populations below damaging levels.</li> <li>▪ If identification of aphid species has not been confirmed, use HIGHER RATE.</li> </ul>
Greenhouse Whitefly	2.8	0.088	<ul style="list-style-type: none"> <li>▪ Apply when adult whiteflies first appear, before populations increase to damaging levels.</li> <li>▪ Application provides SUPPRESSION only. For control, apply in combination with other effective products labeled for use on these listed crops.</li> <li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels.</li> </ul>

**SPRAY RECOMMENDATIONS**

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 10 gallons per acre when applied by ground; use a minimum of 3 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.

**USE RESTRICTIONS**

- Do not apply more than 2.8 oz/acre (0.088 lbs. ai/acre) of **Bestero 50 WG** per single application.
- Do not apply more than 3 applications at the single maximum application rate of 2.8 oz/acre (0.088 lb. ai/acre) of **Bestero 50 WG** per year.
- Do not apply more than 8.4 oz/acre (0.263 lbs. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

**BRASSICA, LEAFY GREENS SUBGROUP 4-16B (0 DAY PHI) INCLUDING:**

Arugula; broccoli, Chinese; broccoli raab; cabbage, abyssian; cabbage, Chinese, bok choy; cabbage, seakale; collards; cress, garden; cress, upland; hanover salad; kale; maca, leaves; mizuna; mustard greens; radish, leaves; rape greens; rocket, wild; shepherd's purse; turnip greens; watercress; cultivars, varieties, and hybrids of these commodities

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids and Plant Bugs	2.0 to 2.8	0.062 to 0.088	<ul style="list-style-type: none"> <li>▪ Begin applications before populations begin to build and before damage is evident according to local pest management guidelines.</li> <li>▪ Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage.</li> <li>▪ Rapidly growing plants may need retreatment. Scout fields often and retreat as necessary to maintain populations below damaging levels.</li> <li>▪ If identification of aphid species has not been confirmed, use HIGHER RATE.</li> </ul>
Greenhouse Whitefly	2.8	0.088	<ul style="list-style-type: none"> <li>▪ Apply when adult whiteflies first appear, before populations increase to damaging levels.</li> <li>▪ Application provides SUPPRESSION only. For control, apply in combination with other effective products labeled for use on these listed crops.</li> <li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels.</li> </ul>

**SPRAY RECOMMENDATIONS**

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 10 gallons per acre when applied by ground; use a minimum of 3 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.

**USE RESTRICTIONS**

- **Bestero 50 WG** may not be applied when watercress is flooded. Water must be turned off 24 hours in advance of an application, and may be turned back on 24 hours after the application.
- Do not apply more than 2.8 oz/acre (0.088 lbs. ai/acre) of **Bestero 50 WG** per single application.
- Do not apply more than 3 applications at the single maximum application rate of 2.8 oz/acre (0.088 lb. ai/acre) of **Bestero 50 WG** per year.
- Do not apply more than 8.4 oz/acre (0.263 lbs. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.



## CUCURBIT VEGETABLES CROP GROUP 9 (0 DAY PHI) INCLUDING:

Chayote (fruit); Chinese waxgourd (Chinese preserving melon); Citron melon; Cucumber; Gherkin; Gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); Muskmelon, hybrids and/or cultivars of *Cucumis melo* (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); Pumpkin; Squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); Squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash; Watermelon (includes hybrids and/or varieties of *Citrullus lanatus*)

### Ground or Aerial Spray

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids and Plant Bugs	2.0 to 2.8	0.062 to 0.088	<ul style="list-style-type: none"><li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li><li>▪ Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage.</li><li>▪ Rapidly growing plants may need retreatment. Scout fields often and retreat as necessary to maintain populations below damaging levels.</li><li>▪ If identification of aphid species has not been confirmed, use HIGHER RATE.</li></ul>
Greenhouse Whitefly	2.8	0.088	<ul style="list-style-type: none"><li>▪ Apply when adult whiteflies first appear, before populations increase to damaging levels.</li><li>▪ Application provides SUPPRESSION only. For control, apply in combination with other effective products labeled for use on these listed crops.</li><li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels.</li></ul>
<b>SPRAY RECOMMENDATIONS</b> <p>Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 10 gallons per acre when applied by ground; use a minimum of 3 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.</p> <b>USE RESTRICTIONS</b> <ul style="list-style-type: none"><li>▪ Do not apply more than 2.8 oz/acre (0.088 lbs. ai/acre) of <b>Bestero 50 WG</b> per single application.</li><li>▪ Do not apply more than 3 applications at the single maximum application rate of 2.8 oz/acre (0.088 lb. ai/acre) of <b>Bestero 50 WG</b> per year.</li><li>▪ Do not apply more than 8.4 oz/acre (0.263 lbs. ai/acre) of <b>Bestero 50 WG</b> or other flonicamid-containing products per year.</li><li>▪ Allow a minimum of 7 days between applications.</li></ul>			

## Drip Irrigation

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids Plant Bugs Whiteflies Thrips	2.8 to 4.28	0.088 to 0.133	<ul style="list-style-type: none"> <li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li> <li>▪ Use LOWER RATE for building populations and use HIGHER RATE for greater populations.</li> <li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels. Rapidly growing plants may need retreatment.</li> <li>▪ <b>Bestero 50 WG</b> must be applied to ensure the product is in the root zone to provide effective control of target pests. Most effective application is when it is applied so that the roots are at or near the site of application. The earlier flonicamid is available to a developing plant, the earlier the protection begins.</li> <li>▪ For transplanted crops, apply <b>Bestero 50 WG</b> within 21 days of transplanting.</li> <li>▪ For direct seeding crops, make applications within 45 days of emergence.</li> </ul>
<b>USE RESTRICTIONS</b> <ul style="list-style-type: none"> <li>▪ Make no more than 2 drip applications per year.</li> <li>▪ Any subsequent applications made with products that contain flonicamid must be foliar applications.</li> <li>▪ Do not apply more than two drip applications at the minimum single application rate of 2.8 oz/ acre (0.088 lbs. ai/acre) of <b>Bestero 50 WG</b> per year.</li> <li>▪ Do not apply more than one drip irrigation application at the maximum single application rate of 4.28 oz/ acre (0.133 lbs. ai/acre) of <b>Bestero 50 WG</b> per year.</li> <li>▪ Do not apply more than 8.4 oz/ acre (0.263 lbs. ai/acre) of <b>Bestero 50 WG</b> or other flonicamid-containing products per year.</li> <li>▪ Allow a minimum of 7 days between applications.</li> </ul>			

SPECIMEN

## GREENHOUSE CUCUMBERS - FOLIAR APPLICATION (0 DAY PHI)

PESTS	Rate of Application	USE DIRECTIONS
Aphids Plant Bugs	2.8 to 4.28 oz per acre or 0.065 to 0.1 oz per 1,000 sq ft or 1.85 to 2.85 gm per 1,000 sq ft	<ul style="list-style-type: none"><li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li><li>▪ Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage.</li><li>▪ Rapidly growing plants may need retreatment. Check plants often and retreat as necessary to maintain populations below damaging levels.</li><li>▪ If identification of aphid species has not been confirmed, use HIGHER RATE.</li></ul>
Greenhouse Whitefly	4.28 oz per acre or 0.1 oz per 1,000 sq ft or 2.85 gm per 1,000 sq ft	<ul style="list-style-type: none"><li>▪ Apply when adult whiteflies first appear, before populations increase to damaging levels.</li><li>▪ Application provides SUPPRESSION only. Apply in combination with other effective products for control.</li><li>▪ Check plants often and retreat as necessary to maintain populations below damaging levels.</li></ul>

### SPRAY RECOMMENDATIONS

Thorough spray coverage is essential for optimum control. Apply in sufficient water to ensure good coverage. Use a minimum of 10 gallons per acre (0.25 gallons per 1,000 sq ft). Finished spray volume should be increased under extreme pest populations or dense plant foliage but do not exceed the point of runoff.

### USE RESTRICTIONS

- Do not apply more than 4.28 oz/acre (0.1 oz per 1,000 sq ft) of **Bestero 50 WG** per single application.
- Do not make more than 2 applications at the maximum single application rate of 4.28 oz/acre (0.1 oz per 1,000 sq ft) of **Bestero 50 WG** per year.
- Do not apply more than 8.56 oz/acre (0.267 lb. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

## GREENHOUSE CUCUMBERS - APPLICATION TO SOIL OR GROWTH MEDIA (0 DAY PHI)

PESTS	Rate of Application	USE DIRECTIONS
Aphids Plant Bugs Greenhouse Whitefly	4.28 oz/acre or 2.8 gm/1,000 sq ft or 12.5 gm/1,000 plants (based on planting density of 1 plant per 4.5 sq ft or 2.4 plants per sq meter).	<ul style="list-style-type: none"><li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li><li>▪ Apply using a drench by hand, drip irrigation, or with motorized calibrated irrigation equipment to the base of the plants.</li><li>▪ Use a minimum of 25 gallons per 1,000 plants but do not exceed the holding capacity of soil or growth media.</li><li>▪ See the <b>Chemigation</b> Section of this label for additional information.</li></ul>

### RECOMMENDATIONS

Many varieties have been tested for tolerance to **Bestero 50 WG** and show good crop safety. To assure the greatest crop safety under a wide range of conditions, treating a few plants before applying to the whole greenhouse is recommended.

### USE RESTRICTIONS

- Do not apply more than 4.28 oz/acre (0.44 oz per 1,000 plants based on 1 plant per 4.5 sq ft or 9,712 plants per acre) of **Bestero 50 WG** per single application.
- Do not make more than 2 applications at the maximum single application rate of 4.28 oz/acre (0.44 oz per 1,000 plants based on 1 plant per 4.5 sq ft or 9,712 plants per acre) of **Bestero 50 WG** per year.
- Do not apply more than 8.56 oz/acre (0.267 lb. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

## FRUITING VEGETABLES CROP GROUP 8-10 (0 DAY PHI) INCLUDING:

African eggplant; bush tomato; bell pepper; cocona; currant tomato; eggplant; garden huckleberry; goji berry; groundcherry; martynia; naranjilla; okra; pea eggplant; pepino; nonbell pepper; roselle; scarlet eggplant; sunberry; tomatillo; tomato; tree tomato; cultivars, varieties, and/or hybrids of these.

### Ground or Aerial Spray

PESTS	Rate of Application		USE DIRECTIONS
	Ounces of Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids Plant Bugs Tomato Psyllid	2.8 to 4.28	0.088 to 0.133	<ul style="list-style-type: none"><li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li><li>▪ Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage.</li><li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels. Rapidly growing plants may need retreatment.</li><li>▪ If identification of aphid species has not been confirmed, use HIGHER RATE.</li></ul>
Greenhouse Whitefly	2.8 to 4.28	0.088 to 0.133	<ul style="list-style-type: none"><li>▪ Apply when adult whiteflies first appear, before populations increase to damaging levels.</li><li>▪ Application provides SUPPRESSION only. For control, apply in combination with other effective products labeled for use on these listed crops.</li><li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels.</li></ul>

#### SPRAY RECOMMENDATIONS

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 10 gallons per acre when applied by ground; use a minimum of 3 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.

#### USE RESTRICTIONS

- Do not apply more than 4.28 oz/acre (0.133 lb. ai/acre) of **Bestero 50 WG** per single application.
- Do not apply more than 3 applications at the minimum single application rate of 2.8 oz/acre (0.088 lb. ai/acre) of **Bestero 50 WG** per year.
- Do not apply more than 2 applications at the maximum single application rate of 4.28 oz/acre (0.133 lb. ai/acre) of **Bestero 50 WG** per year.
- Do not apply more than 8.56 oz/acre (0.267 lbs. ai/ acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

## Drip Irrigation

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids Plant Bugs Whiteflies Tomato Psyllid Thrips	2.8 to 4.28	0.088 to 0.133	<ul style="list-style-type: none"> <li>Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li> <li>Use LOWER RATE for building populations and use HIGHER RATE for greater populations.</li> <li>Scout fields often and retreat as necessary to maintain populations below damaging levels. Rapidly growing plants may need retreatment.</li> <li><b>Bestero 50 WG</b> must be applied to ensure the product is in the root zone to provide effective control of target pests.</li> <li>Most effective application is when it is applied so that the roots are at or near the site of application. The earlier flonicamid is available to a developing plant, the earlier the protection begins.</li> <li>For transplanted crops, apply <b>Bestero 50 WG</b> within 21 days of transplanting.</li> <li>For direct seeding crops, make applications within 45 days of emergence.</li> </ul>
<b>USE RESTRICTIONS</b> <ul style="list-style-type: none"> <li>Make no more than 2 drip applications per year.</li> <li>Any subsequent applications made with products that contain flonicamid must be foliar applications.</li> <li>Do not apply more than two drip applications at the minimum single application rate of 2.8 oz/ acre (0.088 lbs. ai/acre) of <b>Bestero 50 WG</b> per year.</li> <li>Do not apply more than one drip irrigation application at the maximum single application rate of 4.28 oz/acre (0.133 lbs. ai/acre) of <b>Bestero 50 WG</b> per year.</li> <li>Do not apply more than 8.4 oz/ acre (0.263 lbs. ai/acre) of <b>Bestero 50 WG</b> or other flonicamid-containing products per year.</li> <li>Allow a minimum of 7 days between applications.</li> </ul>			

## GREENHOUSE TOMATOES - FOLIAR APPLICATION (0 DAY PHI)

PESTS	Rate of Application	USE DIRECTIONS
Aphids Plant Bugs Tomato Psyllid	2.8 - 4.28 oz per acre or 0.065 - 0.1 oz per 1,000 sq ft or 1.85 - 2.85 gm per 1,000 sq ft	<ul style="list-style-type: none"> <li>Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li> <li>Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage.</li> <li>Rapidly growing plants may need retreatment. Check plants often and retreat as necessary to maintain populations below damaging levels.</li> <li>If identification of aphid species has not been confirmed, use HIGHER RATE.</li> </ul>
Greenhouse Whitefly	4.28 oz per acre or 0.1 oz per 1,000 sq ft or 2.85 gm per 1,000 sq ft	<ul style="list-style-type: none"> <li>Apply when adult whiteflies first appear, before populations increase to damaging levels.</li> <li>Application provides SUPPRESSION only. Apply in combination with other effective products for control.</li> <li>Check plants often and retreat as necessary to maintain populations below damaging levels.</li> </ul>

### SPRAY RECOMMENDATIONS

Thorough spray coverage is essential for optimum control. Apply in sufficient water to ensure good coverage. Use a minimum of 10 gallons per acre (0.25 gallons per 1,000 sq ft). Finished spray volume should be increased under extreme pest populations or dense plant foliage but do not exceed the point of runoff.

### USE RESTRICTIONS

- Do not apply more than 4.28 oz/acre (0.1 oz per 1,000 sq ft) of **Bestero 50 WG** per single application.
- Do not make more than 2 applications at the maximum single application rate of 4.28 oz/acre (0.1 oz per 1,000 sq ft) of **Bestero 50 WG** per year.
- Do not apply more than 8.56 oz/acre (0.267 lb. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

## GREENHOUSE PEPPERS - FOLIAR APPLICATION (0 DAY PHI)

PESTS	Rate of Application	USE DIRECTIONS
Aphids Plant Bugs	2.8 - 4.28 oz per acre or 0.065 - 0.1 oz per 1,000 sq ft or 1.85 - 2.85 gm per 1,000 sq ft	<ul style="list-style-type: none"> <li>Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li> <li>Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage.</li> <li>Rapidly growing plants may need retreatment. Check plants often and retreat as necessary to maintain populations below damaging levels.</li> <li>If identification of aphid species has not been confirmed, use HIGHER RATE.</li> </ul>
Whiteflies Thrips	4.28 oz per acre or 0.1 oz per 1,000 sq ft or 2.85 gm per 1,000 sq ft	<ul style="list-style-type: none"> <li>Apply when whiteflies or thrips first appear, before populations increase to damaging levels.</li> <li>Apply in combination with other effective products for control.</li> <li>Check plants often and retreat as necessary to maintain populations below damaging levels.</li> </ul>

### SPRAY RECOMMENDATIONS

Thorough spray coverage is essential for optimum control. Apply in sufficient water to ensure good coverage. Use a minimum of 10 gallons per acre (0.25 gallons per 1,000 sq ft). Finished spray volume should be increased under extreme pest populations or dense plant foliage but do not exceed the point of runoff.

### USE RESTRICTIONS

- Do not apply more than 4.28 oz/acre (0.1 oz per 1,000 sq ft) of **Bestero 50 WG** per single application.
- Do not make more than 2 applications of **Bestero 50 WG** at the maximum single application rate of 4.28 oz/acre (0.1 oz per 1,000 sq ft) per year.
- Do not apply more than 8.56 oz/acre (0.267 lb. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

## HOP (10 DAY PHI)

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Hop Aphid	1.7 to 2.8	0.053 to 0.088	<ul style="list-style-type: none"> <li>Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li> <li>Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage.</li> <li>Scout fields often and retreat as necessary to maintain populations below damaging levels. Rapidly growing plants may need retreatment.</li> </ul>

### SPRAY RECOMMENDATIONS

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 50 gallons per acre when applied by ground, preferably air-blast; use a minimum of 10 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.

### USE RESTRICTIONS

- Do not apply more than 2.8 oz/acre (0.088 lbs. ai/acre) of **Bestero 50 WG** per single application.
- Do not apply more than 3 applications at the single maximum application rate of 2.8 oz/acre (0.088 lb. ai/acre) of **Bestero 50 WG** per year.
- Do not apply more than 8.4 oz/acre (0.263 lbs. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.



**LEAF PETIOLE VEGETABLE SUBGROUP 22B, CELTUCE AND FLORENCE FENNEL (0 DAY PHI) INCLUDING:**

Cardoon; celery; celery, Chinese; celtuce; fennel, Florence; fuki; rhubarb; udo; zuiki; cultivars, varieties, and hybrids of these commodities

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids and Plant Bugs	2.0 to 2.8	0.062 to 0.088	<ul style="list-style-type: none"> <li>▪ Begin applications before populations begin to build and before damage is evident according to local pest management guidelines.</li> <li>▪ Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage.</li> <li>▪ Rapidly growing plants may need retreatment. Scout fields often and retreat as necessary to maintain populations below damaging levels.</li> <li>▪ If identification of aphid species has not been confirmed, use HIGHER RATE.</li> </ul>
Greenhouse Whitefly	2.8	0.088	<ul style="list-style-type: none"> <li>▪ Apply when adult whiteflies first appear, before populations increase to damaging levels.</li> <li>▪ Application provides SUPPRESSION only. For control, apply in combination with other effective products labeled for use on these listed crops.</li> <li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels.</li> </ul>

**SPRAY RECOMMENDATIONS**

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 10 gallons per acre when applied by ground; use a minimum of 3 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.

**USE RESTRICTIONS**

- Do not apply more than 2.8 oz/acre (0.088 lbs. ai/acre) of **Bestero 50 WG** per single application.
- Do not apply more than 3 applications at the single maximum application rate of 2.8 oz/acre (0.088 lb. ai/acre) of **Bestero 50 WG** per year.
- Do not apply more than 8.4 oz/acre (0.263 lbs. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

## LEAFY GREENS SUBGROUP 4-16A (0 DAY PHI) INCLUDING:

Amaranth, Chinese; amaranth, leafy; aster, Indian; blackjack; cat's whiskers; cham-chwi; cham-na-mul; chervil, fresh leaves; chipilin; chrysanthemum, garland; cilantro, fresh leaves; corn salad; cosmos; dandelion, leaves; dang-gwi, leaves; dillweed; dock; dol-nam-mul; ebolo; endive; escarole; fameflower; feather cockscomb; good king henry; huau-zontle; jute, leaves; lettuce, bitter; lettuce, head; lettuce, leaf; orach; parsley, fresh leaves; plantain, buckhorn; primrose, English; purslane, garden; purslane, winter; radicchio; spinach; spinach, Malabar; spinach, New Zealand; spinach, tanier; Swiss chard; violet, Chinese, leaves; cultivars, varieties, and hybrids of these commodities

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids and Plant Bugs	2.0 to 2.8	0.062 to 0.088	<ul style="list-style-type: none"><li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li><li>▪ Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage.</li><li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels. Rapidly growing plants may need retreatment.</li><li>▪ If identification of aphid species has not been confirmed, use HIGHER RATE.</li></ul>
Greenhouse Whitefly	2.8	0.088	<ul style="list-style-type: none"><li>▪ Apply when adult whiteflies first appear, before populations increase to damaging levels.</li><li>▪ Application provides SUPPRESSION only. For control, apply in combination with other effective products labeled for use on these listed crops.</li><li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels.</li></ul>

### SPRAY RECOMMENDATIONS

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 10 gallons per acre when applied by ground; use a minimum of 3 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.

### USE RESTRICTIONS

- Do not apply more than 2.8 oz/acre (0.088 lbs. ai per acre) of **Bestero 50 WG** per single application.
- Do not apply more than 3 applications at the single maximum application rate of 2.8 oz/acre (0.088 lb. ai/acre) of **Bestero 50 WG** per year.
- Do not apply more than 8.4 oz/acre (0.263 lbs. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

## GREENHOUSE LETTUCE - FOLIAR APPLICATION (0 DAY PHI)

PESTS	Rate of Application	USE DIRECTIONS
Aphids Plant Bugs	2.8 oz per acre or 0.065 oz per 1,000 sq ft or 1.85 gm per 1,000 sq ft	<ul style="list-style-type: none"><li>▪ Make one application before populations begin to build and before damage is evident, according to local pest management guidelines, to control aphids and plants bugs that hide within the lettuce heart.</li></ul>

### SPRAY RECOMMENDATIONS

Thorough spray coverage is essential for optimum control. Apply in sufficient water to ensure good coverage. Use a minimum of 100 gallons per acre (2.5 gallons per 1,000 sq ft). Finished spray volume should be increased under extreme pest populations or dense plant foliage but do not exceed the point of runoff.

### USE RESTRICTIONS

- Do not make more than 1 application at the maximum single application rate of 2.8 oz/acre (0.065 lbs. ai/A) of **Bestero 50 WG** per year.
- Do not apply more than 8.4 oz/acre (0.263 lb. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.

## TUBEROUS AND CORM VEGETABLES SUBGROUP 1C (7 DAY PHI) INCLUDING:

Arracacha; Arrowroot; Chinese artichoke; Jerusalem artichoke; Edible canna; Casava (bitter and sweet); Chayote (root); Chufa; Dasheen; Ginger; Leren; Potato; Sweet potato; Tanier; Turmeric; Yam Bean; Yam (true)

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids Plant Bugs and Potato Psyllid	2.0 to 2.8	0.062 to 0.088	<ul style="list-style-type: none"> <li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li> <li>▪ Use LOWER RATE for building populations; use HIGHER RATE for greater populations and/or dense foliage.</li> <li>▪ Rapidly growing plants may need retreatment. Scout fields often and retreat as necessary to maintain populations below damaging levels.</li> <li>▪ If identification of aphid species has not been confirmed, use HIGHER RATE.</li> <li>▪ <b>Bestero 50 WG</b> is recommended as a supplemental foliar control of Green Peach Aphid in long season potatoes following an at-plant program using systemic insecticides and for primary foliar control of Green Peach Aphid in short season potatoes.</li> <li>▪ Scout fields, before aphid flights begin, at intervals and in locations sufficient to provide representative information on population development. Consult local pest management guidelines for correct procedures.</li> <li>▪ Foliar application of <b>Bestero 50 WG</b> should begin when Green Peach aphid numbers reach 5 per 100 leaves, 1 winged aphid per plant or the observation of wingless aphids; consult local pest management guidelines for specific recommendations.</li> </ul>
Greenhouse Whitefly	2.8	0.088	<ul style="list-style-type: none"> <li>▪ Apply when adult whiteflies first appear, before populations increase to damaging levels.</li> <li>▪ Application provides SUPPRESSION only. For control, apply in combination with other effective products labeled for use on these listed crops.</li> <li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels.</li> </ul>
<b>SPRAY RECOMMENDATIONS</b> Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 10 gallons per acre when applied by ground; use a minimum of 3 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.			
<b>USE RESTRICTIONS</b> <ul style="list-style-type: none"> <li>▪ Do not apply more than 2.8 oz/acre (0.088 lbs. ai/acre) of <b>Bestero 50 WG</b> per single application.</li> <li>▪ Do not apply more than 3 applications at the single maximum application rate of 2.8 oz/acre (0.088 lb. ai/acre) of <b>Bestero 50 WG</b> per year.</li> <li>▪ Do not apply more than 8.4 oz/acre (0.263 lbs. ai/acre) of <b>Bestero 50 WG</b> or other flonicamid-containing products per year.</li> <li>▪ Allow a minimum of 7 days between applications.</li> </ul>			

## ROOT VEGETABLES (EXCEPT SUGAR BEET) SUBGROUP 1B (3 DAY PHI) INCLUDING:

Beet, garden; Burdock, edible; Carrot; Celeriac; Chervil, turnip-rooted; Chicory; Ginseng; Horseradish; Parsley, turnip-rooted; Parsnip; Radish; Radish, oriental; Rutabaga; Salsify; Salsify, black; Salsify, Spanish; Skirret; Turnip

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids and Plant Bugs	2.0 to 2.8	0.062 to 0.088	<ul style="list-style-type: none"><li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li><li>▪ Use LOWER RATE for building populations; use HIGHER RATE for greater populations and/or dense foliage.</li><li>▪ Rapidly growing plants may need retreatment. Scout fields before aphid flights begin, at intervals and in locations sufficient to provide representative information on population development and retreat as necessary to maintain populations below damaging levels. Consult local pest management guidelines for correct procedures.</li><li>▪ Foliar application of <b>Bestero 50 WG</b> should begin when Green Peach aphid numbers reach 5 per 100 leaves, 1 winged aphid per plant or the observation of wingless aphids; consult local pest management guidelines for specific recommendations.</li><li>▪ If identification of aphid species has not been confirmed, use HIGHER RATE.</li></ul>
Greenhouse Whitefly	2.8	0.088	<ul style="list-style-type: none"><li>▪ Apply when adult whiteflies first appear, before populations increase to damaging levels.</li><li>▪ Application provides SUPPRESSION only. For control, apply in combination with other effective products labeled for use on these listed crops.</li><li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels.</li></ul>
<b>SPRAY RECOMMENDATIONS</b> <p>Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 10 gallons per acre when applied by ground; use a minimum of 3 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.</p> <b>USE RESTRICTIONS</b> <ul style="list-style-type: none"><li>▪ Do not apply more than 2.8 oz/acre (0.088 lbs. ai/acre) of <b>Bestero 50 WG</b> per single application.</li><li>▪ Do not apply more than 3 applications at the maximum application rate of 2.8 oz/acre (0.088 lb. ai/acre) of <b>Bestero 50 WG</b> per year.</li><li>▪ Do not apply more than 8.4 oz/acre (0.263 lbs. ai/acre) of <b>Bestero 50 WG</b> or other flonicamid-containing products per year.</li><li>▪ Allow a minimum of 7 days between applications.</li></ul>			

**POME FRUIT CROP GROUP 11-10 (21 DAY PHI) INCLUDING:**

Apple; azarole; crabapple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids and Plant Bugs	2.0 to 2.8	0.062 to 0.088	<ul style="list-style-type: none"> <li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li> <li>▪ Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage.</li> <li>▪ Scout trees often during green tip through pink growth stages and post-bloom and retreat as necessary to maintain populations below damaging levels.</li> <li>▪ If identification of aphid species has not been confirmed, use HIGHER RATE.</li> </ul>

**SPRAY RECOMMENDATIONS**

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 50 gallons per acre when applied by ground, preferably air-blast; use a minimum of 10 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.

**USE RESTRICTIONS**

- Do not apply more than 2.8 oz/acre (0.088 lbs. ai/acre) of **Bestero 50 WG** per single application.
- Do not apply more than 3 applications at the maximum single application rate of 2.8 oz/acre (0.088 lb. ai/acre) of **Bestero 50 WG** per year.
- Do not apply more than 8.4 oz/acre (0.263 lbs. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

**STONE FRUIT CROP GROUP 12-12 (14 DAY PHI) INCLUDING:**

Apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids and Plant Bugs	2.0 to 2.8	0.062 to 0.088	<ul style="list-style-type: none"> <li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li> <li>▪ Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage.</li> <li>▪ Scout trees often and retreat as necessary to maintain populations below damaging levels.</li> <li>▪ If identification of aphid species has not been confirmed, use HIGHER RATE.</li> </ul>

**SPRAY RECOMMENDATIONS**

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 50 gallons per acre when applied by ground, preferably air-blast; use a minimum of 10 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.

**USE RESTRICTIONS**

- Do not apply more than 2.8 oz/acre (0.088 lbs. ai/acre) of **Bestero 50 WG** per single application.
- Do not apply more than 3 applications at the single maximum application rate of 2.8 oz/acre (0.088 lb. ai/acre) of **Bestero 50 WG** per year.
- Do not apply more than 8.4 oz/acre (0.263 lbs. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

**TREE NUTS CROP GROUP 14-12 (40 DAY PHI) INCLUDING:**

African nut-tree; Almond; Beechnut; Brazil nut; Brazilian pine; Bunya; Bur oak; Butternut; Cajou nut; Candlenut; Cashew; Chestnut; Chinquapin; Coconut; Coquito nut; Dika nut; Ginkgo; Guiana chestnut; Hazelnut; Heartnut; Hickory nut; Japanese horse-chestnut; Macadamia nut; Mongongo nut; Monkey-pot; Monkey puzzle nut; Okari nut; Pachira nut; Peach palm nut; Pecan; Pequi; Pili nut; Pine nut; Pistachio; Sapucaia nut; Tropical almond; Walnut, black; Walnut, English; Yellowhorn

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids	2.0 to 2.8	0.062 to 0.088	<ul style="list-style-type: none"> <li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li> <li>▪ Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage.</li> <li>▪ Scout trees often and retreat as necessary to maintain populations below damaging levels.</li> <li>▪ If identification of aphid species has not been confirmed, use HIGHER RATE.</li> </ul>
Plant Bugs	2.8	0.088	<ul style="list-style-type: none"> <li>▪ Apply when bugs first appear, before populations increase to damaging levels.</li> <li>▪ Application provides SUPPRESSION only. For control, apply in combination with other effective products labeled for use on these listed crops.</li> <li>▪ Scout trees often and retreat as necessary to maintain populations below damaging levels.</li> </ul>

**SPRAY RECOMMENDATIONS**

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 50 gallons per acre when applied by ground, preferably air-blast; use a minimum of 10 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.

**USE RESTRICTIONS**

- Do not apply more than 2.8 oz/acre (0.088 lbs. ai/acre) of **Bestero 50 WG** per single application.
- Do not apply more than 3 applications at the single maximum application rate of 2.8 oz/acre (0.088 lb. ai/acre) of **Bestero 50 WG** per year.
- Do not apply more than 8.4 oz/acre (0.263 lbs. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.



## CITRUS FRUIT GROUP (0 DAY PHI) INCLUDING:

Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; mount white lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin); tangor; trifoliolate orange; uniq fruit; cultivars, varieties, and/or hybrids of these

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Asian Citrus Psyllid	2.8 to 5.7	0.088 to 0.178	<ul style="list-style-type: none"><li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li><li>▪ Use the LOWER RATE for building populations and use the HIGHER RATE for greater populations and/or dense foliage.</li><li>▪ Scout trees often and retreat as necessary to maintain populations below damaging levels.</li><li>▪ For optimum performance, include an adjuvant. Only use adjuvants known to be safe on citrus crops.</li></ul>
Aphids	2.8	0.088	<ul style="list-style-type: none"><li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li><li>▪ Scout trees often and retreat as necessary to maintain populations below damaging levels.</li></ul>

### SPRAY RECOMMENDATIONS

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 50 gallons per acre when applied by ground, preferably air-blast; use a minimum of 10 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage. Spray adjuvants may improve coverage.

### USE RESTRICTIONS

- Do not apply more than 5.7 oz/acre (0.178 lb. ai/acre) of **Bestero 50 WG** per single application.
- Do not apply more than 3 applications of **Bestero 50 WG** at the minimum single application rate of 2.8 oz./acre (0.088 lb. ai/acre) per year.
- Do not apply more than 1 application of **Bestero 50 WG** at the maximum single application rate of 5.7 oz./acre (0.178 lb. ai/acre) per year.
- Do not apply more than a total of 8.4 oz/acre (0.263 lbs. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

**GRAPES AND OTHER SMALL VINE CLIMBING FRUIT (EXCEPT FUZZY KIWIFRUIT) SUBGROUP 13-07F (7 DAY PHI) INCLUDING:**

Amur river grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and hybrids of these commodities

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids Plant Bugs ( <i>Lygus</i> spp.)	2.8 to 4.2	0.088 to 0.131	<ul style="list-style-type: none"> <li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li> <li>▪ Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage.</li> <li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels. Rapidly growing plants may need retreatment.</li> <li>▪ If identification of aphid species has not been confirmed, use HIGHER RATE.</li> </ul>
Vine Mealybugs	4.2 to 5.6	0.131 to 0.175	<ul style="list-style-type: none"> <li>▪ Apply when mealybugs first appear, before populations increase to damaging levels.</li> <li>▪ Application provides SUPPRESSION only. For control, apply in combination with other effective products labeled for use on these listed crops.</li> <li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels.</li> </ul>

**SPRAY RECOMMENDATIONS**

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 50 gallons per acre when applied by ground; use a minimum of 10 gallons per acres by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage. Spray adjuvants may improve coverage but do not use binder or sticker- type surfactants.

**USE RESTRICTIONS**

- Do not apply more than 1 application of **Bestero 50 WG** at the maximum single application rate of 5.6 oz/acre (0.175 lbs. ai/acre) per year.
- Do not apply more than 8.4 oz/acre (0.263 lbs. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year. This is equivalent to:
  - o 3 applications at 2.8 oz/acre of **Bestero 50 WG** per year, or
  - o 2 applications at 4.2 oz/acre of **Bestero 50 WG** per year
- Allow a minimum of 7 days between applications.

**LOW GROWING BERRY SUBGROUP 13-07G (0 DAY PHI) INCLUDING:**

Bearberry; bilberry; blueberry, lowbush; cloudberry; cranberry; lingonberry; muntries; partridgeberry; strawberry; cultivars, varieties, and/or hybrids of these

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids and Plant Bugs ( <i>Lygus</i> spp.) Strawberry Aphid	2.8	0.088	<ul style="list-style-type: none"> <li>▪ Apply when Aphids or Lygus first appear in the field and before populations reach high levels.</li> <li>▪ <b>Bestero 50 WG</b> will stop Aphid and Lygus feeding rapidly but it may take several days to see a reduction in Aphid and Lygus numbers. Reapply when new insects are detected.</li> <li>▪ Two sequential applications of <b>Bestero 50 WG</b> result in better Aphid and Lygus control than a single application.</li> </ul>

**SPRAY RECOMMENDATIONS**

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 50 gallons per acre when applied by ground; use a minimum of 10 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage. Spray adjuvants may improve coverage but do not use binder or sticker-type surfactants. Only use adjuvants known to be safe on strawberries.

**USE RESTRICTIONS**

- Do not make more than two applications of **Bestero 50 WG** without rotating to an insecticide with a different mode of action.
- Do not apply more than 2.8 oz./acre (0.088 lbs. ai per acre) of **Bestero 50 WG** per single application.
- Do not apply more than 3 applications at the maximum single application rate of 2.8 oz/acre (0.088 lb. ai/acre) of **Bestero 50 WG** per year.
- Do not apply more than 8.4 oz./acre (0.263 lbs. ai per acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

**RAPESEED SUBGROUP 20A (7 DAY PHI) INCLUDING:**

Borage; crambe; cuphea; echium; flax seed; gold of pleasure; hare's ear mustard; lesquerella; lunaria; meadowfoam; milkweed; mustard seed; oil radish; poppy seed; rapeseed; sesame; sweet rocket; cultivars, varieties, and/or hybrids of these

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids and Plant Bugs ( <i>Lygus</i> spp.) Cabbage Aphid Green Peach Aphid Turnip Aphid	2.8	0.088	<ul style="list-style-type: none"> <li>▪ Apply when Aphids or Lygus first appear in the field and before populations reach high levels.</li> <li>▪ <b>Bestero 50 WG</b> will stop Aphid and Lygus feeding rapidly but it may take several days to see a reduction in Aphid or Lygus numbers. Reapply when new insects are detected.</li> <li>▪ Two sequential applications of <b>Bestero 50 WG</b> result in better Aphid and Lygus control than a single application.</li> </ul>

**SPRAY RECOMMENDATIONS**

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 10 gallons per acre when applied by ground; use a minimum of 3 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage. Spray adjuvants may improve coverage but do not use binder or sticker-type surfactants. Only use adjuvants known to be safe on canola.

**USE RESTRICTIONS**

- Do not make more than two applications of **Bestero 50 WG** without rotating to an insecticide with a different mode of action.
- Do not apply more than 2.8 oz./acre (0.088 lbs. ai per acre) of **Bestero 50 WG** per single application.
- Do not apply more than 3 applications at the single maximum application rate of 2.8 oz/acre (0.088 lb. ai/acre) of **Bestero 50 WG** per year.
- Do not apply more than 8.4 oz./acre (0.263 lbs. ai per acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

## SUNFLOWER SUBGROUP 20B (0 DAY PHI)

Calendula; castor oil plant; Chinese tallowtree; euphorbia; evening primrose; jojoba; Niger seed; rose hip; safflower; stokes aster; sunflower; tallowwood; tea oil plant; veronia; cultivars, varieties, and/or hybrids of these

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids and Plant Bugs	2.8	0.088	<ul style="list-style-type: none"><li>▪ Apply when insects first appear in the field and before populations reach high levels.</li><li>▪ <b>Bestero 50 WG</b> will stop feeding rapidly but it may take several days to see a reduction in insect numbers. Reapply when new insects are detected.</li><li>▪ Two sequential applications of <b>Bestero 50 WG</b> result in better control than a single application.</li></ul>

### SPRAY RECOMMENDATIONS

Thorough spray coverage of plant foliage is essential for optimum control. Use a minimum of 5 gallons per acre by ground and 3 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.

### USE RESTRICTIONS

- Do not make more than two applications of **Bestero 50 WG** without rotating to an insecticide with a different mode of action.
- Do not apply more than 2.8 oz/acre (0.088 lbs. ai/acre) of **Bestero 50 WG** per single application.
- Do not apply more than 3 applications at the single maximum application rate of 2.8 oz/acre (0.088 lb. ai/acre) of **Bestero 50 WG** per year.
- Do not apply more than 8.4 oz/acre (0.263 lbs. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

SPECIMEN

## LEGUME VEGETABLES (SUCCULENT AND DRIED) CROP GROUP 6 except soybeans (7 DAY PHI):

Includes all members of the Legume Vegetables, succulent and dried, Crop Group 6, except soybeans: Bean (*Lupinus*) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); bean (*Phaseolus*) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean); bean (*Vigna*) (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean); broad bean (fava); chickpea (garbanzo); guar; jackbean; lablab bean; lentil; pea (*Pisum*) (includes dwarf pea, edible-podded pea, English pea, field pea, garden pea, green pea, snow-pea, sugar snap pea); pigeon pea; soybean (immature seed); sword bean

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids and Plant Bugs ( <i>Lygus</i> spp.)	2.8	0.088	<ul style="list-style-type: none"><li>▪ Apply when Aphids or Lygus first appear in the field and before populations reach high levels.</li><li>▪ <b>Bestero 50 WG</b> will stop Aphid and Lygus feeding rapidly but it may take several days to see a reduction in Aphid or Lygus numbers. Reapply when new insects are detected.</li><li>▪ Two sequential applications of <b>Bestero 50 WG</b> result in better Aphid and Lygus control than a single application.</li></ul>

### SPRAY RECOMMENDATIONS

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 10 gallons per acre when applied by ground; use a minimum of 3 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage. Spray adjuvants may improve coverage but do not use binder or sticker-type surfactants. Only use adjuvants known to be safe on canola.

### USE RESTRICTIONS

- Do not make more than two applications of **Bestero 50 WG** without rotating to an insecticide with a different mode of action.
- Do not apply more than 2.8 oz./acre (0.088 lbs. ai per acre) of **Bestero 50 WG** per single application.
- Do not apply more than 3 applications at the single maximum application rate of 2.8 oz/acre (0.088 lb. ai/acre) per year.
- Do not apply more than 8.4 oz./acre (0.263 lbs. ai per acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

## ALFALFA (FORAGE, HAY AND SEED) (14 DAY PHI)

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids and Plant Bugs	2.8	0.088	<ul style="list-style-type: none"><li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li><li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels. Rapidly growing plants may need retreatment.</li></ul>

### SPRAY RECOMMENDATIONS

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 20 gallons per acre when applied by ground; use a minimum of 10 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.

### USE RESTRICTIONS

- Do not apply more than 2.8 oz/ acre (0.088 lbs. ai per acre) of **Bestero 50 WG** per single application.
- Do not apply more than 2 applications at the maximum single application rate of 2.8 oz/acre (0.088 lb. ai/acre) of **Bestero 50 WG** per year.
- Do not apply more than 5.6 oz/acre (0.18 lbs. ai/acre) of **Bestero 50 WG** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.
- Do not allow foraging of fields or harvesting for hay within 14 days of the last application.

**CLOVER (FORAGE AND HAY) GROWN WEST OF THE ROCKIES (60 DAY PHI)**

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids and Plant Bugs	2.8	0.088	<ul style="list-style-type: none"> <li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li> <li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels. Rapidly growing plants may need retreatment.</li> </ul>
<b>SPRAY RECOMMENDATIONS</b> Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 20 gallons per acre when applied by ground; use a minimum of 10 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.			
<b>USE RESTRICTIONS</b> <ul style="list-style-type: none"> <li>▪ Do not apply more than 2.8 oz/acre (0.088 lbs. ai/acre) of <b>Bestero 50 WG</b> per application.</li> <li>▪ Do not apply more than 2 applications at the maximum single application rate of 2.8 oz/acre (0.088 lb. ai/acre) of <b>Bestero 50 WG</b> per year.</li> <li>▪ Do not apply more than 5.6 oz/acre (0.18 lbs. ai/acre) of <b>Bestero 50 WG</b> or other flonicamid-containing products per year.</li> <li>▪ Allow a minimum of 7 days between applications.</li> <li>▪ Do not allow foraging of fields or harvesting for hay within 60 days of the last application.</li> </ul>			

**MINT: Spearmint, Peppermint (7 DAY PHI)**

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Bestero 50 WG/Acre	Lbs. Active Ingredient/Acre	
Aphids	2.0 to 2.8	0.062 to 0.088	<ul style="list-style-type: none"> <li>▪ Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.</li> <li>▪ Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage.</li> <li>▪ Scout fields often and retreat as necessary to maintain populations below damaging levels. Rapidly growing plants may need retreatment.</li> <li>▪ If identification of aphid species has not been confirmed, use HIGHER RATE.</li> </ul>
<b>SPRAY RECOMMENDATIONS</b> Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 20 gallons per acre when applied by ground; use a minimum of 10 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.			
<b>USE RESTRICTIONS</b> <ul style="list-style-type: none"> <li>▪ Do not apply more than 2.8 oz/acre (0.088 lbs. ai per acre) of <b>Bestero 50 WG</b> per single application.</li> <li>▪ Do not apply more than 3 applications at the maximum single application rate of 2.8 oz/acre (0.088 lbs. ai per acre) of <b>Bestero 50 WG</b> per year.</li> <li>▪ Do not apply more than 8.4 oz/acre (0.263 lbs. ai/acre) of <b>Bestero 50 WG</b> or other flonicamid-containing products per year.</li> <li>▪ Allow a minimum of 14 days between applications.</li> </ul>			



## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area. Carefully open containers. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL:** Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER HANDLING:

**Plastic Container:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

## LIMITATION OF WARRANTY AND LIABILITY

**IMPORTANT: READ BEFORE USE.** Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Bestero™ is a trademark of Atticus, LLC.

Beleaf® 50SG is a registered trademark of Ishihara Sangyo Kaisha, Ltd.

20230913ap1