



INSECTICIDE

CONTAINS FLONICAMID, THE ACTIVE INGREDIENT USED IN BELEAF® 50SG

PUT AN END TO INSECT FEEDING QUICKLY

Bestero™ 50 WG is a highly effective insecticide that delivers exceptional pest control for a variety of crops. It aids in controlling leafy and fruiting vegetables, cucurbits, brassicas, potatoes, pome, stone fruits, and more. The powerful formula contains flonicamid, an active ingredient known to eliminate pests within only a few minutes. Bestero 50 WG works by disrupting the nervous system of insects and stopping their ability to feed, either through contact or ingestion.

An excellent partner in pest management, Bestero 50 WG offers easy mixing with no known cross-resistance to other products. A versatile insecticide, it can be applied by ground, air, or chemigation. Maximize your yield potential and put an end to harmful insects in your specialty crops with Bestero 50 WG.

KEY BENEFITS

- Translaminar movement for contact and ingestion control
- Low impact on many beneficial insects
- Excellent pest management partner with no known cross-resistance
- Long-lasting residual control

KEY USES

- | | |
|---------------|----------------|
| • Cantaloupes | • Potatoes |
| • Celery | • Pumpkins |
| • Lettuce | • Strawberries |

PRODUCT NOTES

EPA REGISTRATION NUMBER

91234-300

ACTIVE INGREDIENT

Flonicamid 50%

FORMULATION

Water Dispersible Granule

IRAC NUMBER

29

SIGNAL WORD

Warning

PACKAGE SIZE

6 x 1.5 lb

RESTRICTED USE

No



LABEL



PRODUCT INFO



PORTFOLIO



Bootstrapped and ready to serve, we deliver battle-tested chemistries and an experience like no other. Proud to be 100% American-owned, our mission is to help you every step of the way.





CROP ROTATION RESTRICTIONS

Following the application of Bestero 50 WG insecticide, any crop listed on the product label may be planted as a rotational crop. All other crops may be planted 30 days after the last application of Bestero 50 WG.

MAXIMUM SEASONAL APPLICATION RATES AND PRE-HARVEST INTERVALS		
CROP (COMPLETE LIST FOUND IN THE PRODUCT LABEL)	MAXIMUM SEASONAL TOTAL (POUNDS ACTIVE INGREDIENT)	PHI DAYS
Brassica Head and Stem Vegetables	0.263	0
Brassica Leafy Greens	0.263	0
Cucurbit Vegetables	0.263	0
Fruiting Vegetables (Except Cucurbits)	0.267	0
Grapes & Other Small Vine Fruit (Except Fuzzy Kiwifruit)	0.263	7
Hops	0.263	10
Leaf Petiole Vegetables	0.263	0
Leafy Greens	0.263	0
Tuberous and Corn Vegetables	0.263	7
Legume Vegetables (Except Soybeans)	0.263	7
Root Vegetables (Except Sugar Beets)	0.263	3
Pome Fruit	0.263	21
Stone Fruit	0.263	14
Tree Nuts	0.263	40
Citrus Fruit	0.263	0
Greenhouse Cucumbers	0.267	0
Greenhouse Tomatoes	0.267	0
Greenhouse Peppers/Eggplants	0.267	0
Greenhouse Lettuce	0.088	0
Rapeseed	0.263	7
Sunflower Commodities	0.263	0
Low Growing Berries	0.263	0
Strawberry	0.263	0
Kohlrabi	0.263	0
Celtuce	0.263	0
Florence Fennel	0.263	0
Mint (Spearmint and Peppermint)	0.263	7
Alfalfa (Forage, Hay, and Seed)	0.18	14
Clover (Forage and Hay) (Grown West of the Rockies)	0.18	60

APPLICATION INFORMATION

Clean your spray system before using Bestero 50 WG. Fill the tank halfway with clean water, agitate, and then continue filling to the desired level.

Use a boom and nozzle sprayer with appropriate nozzles that produce minimal fine spray droplets. Keep the spray pressure below 30 psi and adjust spray tops to a minimum of 18 inches above the crop. For chemigation use, refer to the product label for instructions.

KEY INSECTS

Apple aphid
Asian citrus psyllid
Black bean aphid
Black cherry aphid
Black pecan aphid
Blackmargined aphid
Blue alfalfa aphid
Cabbage aphid
Cotton / Melon aphid
Cotton fleahopper
Cowpea aphid
English grain aphid
Foxglove aphid
Green peach aphid
Greenbug
Greenhouse whitefly
Hop aphid
Leaf curl plum aphid
Mealy plum aphid
Pea aphid
Potato aphid
Potato psyllid
Red lettuce aphid
Rosy apple aphid
Soybean aphid
Spirea aphid
Spotted alfalfa aphid
Strawberry aphid
Tarnished plant bug
Tomato psyllid
Turnip aphid
Vine mealybug
Western plant bug
Woolly apple aphid
Yellow pecan aphid

(Refer to product label for complete list)