TECHNICAL INFORMATION BULLETIN

FLUTRIAFOL	GROUP	3	FUNGICIDE

RONDEL™

EPA Reg. No. 91234-384

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CALIFORNIA
For the Suppression of *Fusarium* Sp. via Drip Chemigation in Celery

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for **Rondel™** before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

CROP	DISEASE	RATE	RETREATMENT INTERVAL (DAYS)
Celery	*Fusarium Yellows/Fusarium Wilt* (Suppression only) *(Fusarium oxysporum f.sp apii)*	5-7 fl oz/A	14

Directions for Use:

- **Rondel** will provide suppression of listed diseases.
- Product is to be applied through drip irrigation. Recommend initial application at transplanting. For spring and fall planting, a second application is required, approximately 30 days later. For fall planting, additional applications may be required at 14 day intervals, for a maximum of 4 applications per season.

Restrictions:

- PHI: Do not apply within 7 days of harvest.
- Do not apply more than 7.0 fl. oz. (0.114 lb ai) product/A application.
- Do not make more than 4 applications/year.
- Do not apply more than 28 fl oz of product/A per year.
- Do not apply more than 0.455 lb ai of flutriafol or flutriafol containing products/A/year.
- Restricted Entry Interval (REI) is 12 hours
- Pre harvest interval (PHI) – 0 days

Atticus

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513