TECHNICAL INFORMATION BULLETIN

FLUTRIAFOL	GROUP	3	FUNGICIDE

RONDEL™

EPA Reg. No. 91234-384

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF ARIZONA
For the Suppression of *Fusarium Sp*. via Drip Chemigation in Tomatoes

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for **Rondel™** before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

CROP	DISEASE	RATE	RETREATMENT INTERVAL (DAYS)
Tomatoes (see label for listed crops)	*Fusarium oxysporum f. sp. Lycopersici* *Fusarium falciforme*	7 fl oz/A (0.114 lbs ai/acre)	14

Application Directions:
Apply as directed through pressurized irrigation methods. Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development at 2 weeks intervals. Best results have been obtained when **Rondel™** is applied preventively, at 2 weeks after transplanting followed by subsequent applications at 2 weeks intervals.

Restrictions:

- No single application may exceed 7 fl oz of product/A
- Do not apply more than 4 applications per year
- Do not apply more than 28 fl oz of product/A per year
- Do not apply more than 0.455 lb ai of flutriafol/A per year
- Restricted Entry Interval (REI) is 12 hours
- Pre harvest interval (PHI) – 0 days

Atticus

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513