# TECHNICAL INFORMATION BULLETIN

| FLUTRIAFOL | GROUP | 3 | FUNGICIDE |
|---|---|---|---|

# RONDEL™

**EPA Reg. No. 91234-384**

**FIFRA Section 2(ee) Recommendation**

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

**FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CALIFORNIA**
**For Management of Grape Trunk Disease Complex on Grapes, Trunk Disease Complex on Tree Nuts, and Hull Rot in Almond**

## DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for **Rondel™** before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

| CROP | DISEASES | RATE (fl. oz./A) | APPLICATION DIRECTIONS | PHI (Days) |
|---|---|---|---|---|
| **Grapes** | **Vinewood Disease**<br>• *Botryosphaeria dothidea*<br>• *Eutypa lata*<br>• *Phomopsis canker*<br>• *Phaeoacremonium parasiticum*<br>• *Neofusicoccum parvum*<br>• *Neofusicoccum mediterraneum* | 5.0<br>(0.081 lb. ai) | **New Vineyards (1-3 years old)** Make initial application at the beginning of Spring Root Flush. Make second application postharvest during Fall Root Flush. All applications must be made through a pressurized irrigation system. To achieve maximum performance, use a minimum of 10 fl. oz./A for the year.<br><br>**Established Vineyards (>3 years old)** Make initial application at the beginning of Spring Root Flush. Make second application 14 days later. Make third application at the start of Veraison. Make fourth application post-harvest during Fall Root Flush. All applications must be made through a pressurized irrigation system. To achieve maximum performance, use a minimum of 20 fl. oz./A for the year. | 14 |

Atticus

Manufactured for:
**Atticus, LLC**
940 NW Cary Parkway, Suite 200
Cary, NC 27513

| | | | | |
|---|---|---|---|---|
| **Tree Nuts** (African Tree Nut, Almond, Brazil Nut, Burr Oak, Butternut, Cajou, Cashew, Castanha-DoMaranhao, Coconut, Coquito Nut, Dika Nut, Guiana Chestnut, Hazelnut, Heartnut, Hickory Nut, Japanese Horse- Chestnut, Macadamia Nut, Monogongo Nut, Monkey-Pot, Pachira Nut, Pecan, Sapucaia Nut) | **Trunk diseases**<br>• *Botryosphaeria dothidea*<br>• *Cytospora sorbicola*<br>• *Eutypa lata*<br>• *Ceratocystis destructans,*<br>• *Neofusicoccum parvum*<br>• *Neofusicoccum mediterraneum*<br>• *Calosphaeria pulchella,*<br>• *Phomopsis canker*<br>• *Ganoderma* | 7.0 (0.114 lb. ai) | Make initial application of 7 fl. oz./A at the beginning of Spring Root Flush. Reapply at 30-day intervals as needed. Make final application at Fall Root Flush (after harvest). Apply through a pressurized irrigation system. To achieve maximum performance, make 3-4 applications at 7 fl. oz./A for the year. | 14 |
| **Almond** | **Hull Rot**<br>• *Monilinia fructicola*<br>• *Aspergillus niger*<br><br>**SUPPRESSION:**<br>• *Rhizopus stolonifera* | 7.0 (0.114 lb. ai) | Apply as a foliar spray in sufficient water to obtain thorough coverage of foliage and/or fruit. Foliar application can be as early as 3-4 weeks before hull split. For best results and under high disease pressure apply early hull split and again 14-days later. Refer to university and/or extension guidelines for best application timings and IPM. | 14 |

For management of wood diseases on grapes and tree nuts, apply **Rondel™** as directed through pressurized irrigation methods such as macro-sprinkler, micro-sprinkler, or drip systems. For best results, use **Rondel™** as part of a long-term management program.

**RESTRICTIONS:**

**Grapes**

- No single application may exceed 5 fl. oz. (0.081 lb. ai) of product/A.
- Do not apply more than 6 applications per year.
- Do not apply more than 30 fl. oz. (0.488 lb. ai) of product/A per year.
- Do not apply more than 0.488 lb. ai flutriafol/A per year.
- Restricted Entry Interval (REI) is 5 days for girdling or turning of grapes. For all other activities the REI is 12 hours.
- **Preharvest Interval (PHI):** Do not apply within 14 days of harvest.

**Tree Nuts**

- No single application may exceed 7 fl. oz. (0.114 lb. ai) of product/A.
- Do not apply more than 4 applications per year.
- Do not apply more than 28 fl. oz. (0.455 lb. ai) of product/A per year.
- Do not apply more than 0.455 lb. ai flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours.
- **Preharvest Interval (PHI)**: Do not apply within 14 days of harvest.