TECHNICAL INFORMATION BULLETIN

FLUTRIAFOL	GROUP	3	FUNGICIDE

RONDEL™

EPA Reg. No. 91234-384

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

This recommendation for Rondel™ is valid until December 31, 2030 or until withdrawn, cancelled or suspended.

Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist

FOR DISTRIBUTION AND USE ONLY AZ, CO, CT, DE, ID, IL, IN, IA, KY, ME, MD, MA, MN, MO, NH, NJ, NM, NC, OH, OR, PA, SC, TN, UT, VT, VA, WA, WV, AND WI

For Control of Powdery Mildew in Pome, Hops, and Stone Fruit Via Drip Chemigation

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for **Rondel™** before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

CROP	DISEASE	RATE (fl. oz./A)	APPLICATION DIRECTIONS	Retreatment Interval (Days)
Hops	Powdery mildew *Podosphaera macularis*	7 (0.144 lb. ai)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary.	14
Pome Fruit (See label for list of pome fruit)	Powdery Mildew *Podosphaera leucotricha*	6 (0.098 lb. ai)		7
Stone Fruit (See label for list of stone fruit)	Powdery mildew *Podosphaera clandestina*	7 (0.144 lb. ai)		7
When applied as directed through drip irrigation and microsprinklers, **Rondel™** will prevent powdery mildew on the foliage of hops, pome, and stone fruit.				

Atticus

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513