

TECHNICAL INFORMATION BULLETIN

PROTHIOCONAZOLE

GROUP

3

FUNGICIDE



EPA Reg. No. 91234-345

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF MASSACHUSETTS AND NEW JERSEY

For Use in Aerial Applications to Cranberries

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for **Provixo™ 480 SC** before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

CROPS	DISEASE CONTROLLED	RATE OF PROVIXO 480 SC (FL. OZ./ACRE)	APPLICATION DIRECTIONS
Highbush Cranberry	Septoria leaf spot (<i>Septoria</i> spp.) Monilinia blight (<i>Monilinia vaccinicorymbosi</i>) Valdensinia leaf spot (<i>Valdensinia heterodoxa</i>) Leaf Rust (<i>Thekopsora minima</i>) Anthracnose (<i>Colletotrichum gloeosporioides</i>) Botrytis blight (<i>Botrytis cinerea</i>) Phomopsis canker and twig blight (<i>Phomopsis vaccinii</i>) Alternaria fruit rot (<i>Alternaria</i> spp.) White pine blister rust (<i>Cronartium ribicola</i>)	5.7 (0.178 lb. ai)	Apply Provixo 480 SC at the first sign of disease. For aerial applications: Apply Provixo 480 SC using a minimum of 2 gpa spray solution. An adjuvant may be used to improve spray coverage. Refer to the adjuvant product label for specific use directions. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals. Repeat applications as needed using a 7 to 10-day spray interval if conditions remain favorable for continued or increasing disease development.
Other Requirements:			
<ul style="list-style-type: none"> • Apply up to two (2) applications of Provixo 480 SC per year. • A maximum of 11.4 fl. oz. (0.356 lb. prothioconazole) of Provixo 480 SC may be applied per acre per year. • Do not apply within 7 days of harvest. 			



Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

TECHNICAL INFORMATION BULLETIN

CROPS	DISEASES CONTROLLED	RATE OF PROVIXO 480 SC (FL. OZ./ACRE)	APPLICATION DIRECTIONS
Cranberry	Fruit rot: <i>Coleophoma empetri</i> , <i>Glomerella cingulata</i> , <i>Phyllosticta vaccinii</i> , <i>Physalospora vaccinii</i> , <i>Allantophomopsis lycopodina</i> , <i>Allantophomopsis cytispora</i> , <i>Fusicoccum putrefaciens</i> , <i>Penicillium</i> spp., <i>Phomopsis vaccinii</i> , <i>Colletotrichum acutatum</i> , <i>Botrytis</i> spp., <i>Monilinia</i> spp. Valdensinia leaf spot (<i>Valdensinia heterodoxa</i>)	5.0 (0.156 lb. ai)	For aerial applications: Apply Provixo 480 SC using a minimum of 2 gpa spray solution. An adjuvant may be used to improve spray coverage. Refer to the adjuvant product label for specific use directions. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals. Repeat applications as needed using a 7 to 10-day spray interval if conditions remain favorable for continued or increasing disease development.
Other Requirements: <ul style="list-style-type: none"> • Apply up to two (2) applications of Provixo 480 SC per year. • A maximum of 10.0 fl. oz. (0.313 lb. prothioconazole) of Provixo 480 SC may be applied per acre per year. • Do not apply within 45 days of harvest. 			



Manufactured for:
Atticus, LLC
 940 NW Cary Parkway, Suite 200
 Cary, NC 27513