TECHNICAL INFORMATION BULLETIN

CHLORANTRANILIPROLE	GROUP	28	INSECTICIDE

Osaria™ ULTRA 70 WDG

EPA Reg. No. 91234-379

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA
For Control of Orange Tortix on Grapes

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for **Osaria™ Ultra 70 WDG** before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

CROP	INSECT	RATE PER ACRE	PHI (days)	REI (hours)
Grapes	Orange tortrix	1.3 - 2.2 oz (0.055 – 0.098 lb a.i.)	14	4

- Do not apply more than 4.6 oz **Osaria™ Ultra 70 WDG** or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.
- Do not make more than 4 applications per calendar year.
- Consult the UC IPM Pest Management Guidelines for adult emergence monitoring using pheromone traps and degree day accumulation to determine the timing of application to first or second generation. For best results, make the first application at initiation of egg hatch, to small larvae, or first signs of infestations. To manage resistance, target only one of the two generations with **Osaria™ Ultra 70 WDG** or any other group 28 containing insecticide.
- Use higher rates of **Osaria™ Ultra 70 WDG** for moderate to heavy insect pressure. Make repeat applications on 7 to 10 day schedule if monitoring indicates continued feeding activity.
- Do not apply less than 30 gal water per acre by ground.

Atticus

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513