

TECHNICAL INFORMATION BULLETIN

FLONICAMID GROUP 29 INSECTICIDE



EPA Reg. No. 91234-301

FIFRA Section 2(ee) Recommendation

This recommendation expires December 31, 2029

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

FOR DISTRIBUTION AND USE WITHIN THE STATE OF TEXAS ONLY

FOR MANAGEMENT OF TWO-SPOTTED COTTON LEAFHOPPER IN COTTON

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for **CablON™** before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

PESTS	Rate of Application		USE DIRECTIONS
	Ounces Cablon /Acre	Lbs. Active Ingredient /Acre	
Two-Spotted Cotton Leafhopper (<i>Amrasca biguttula</i>)	1.7 to 2.8	0.053 - 0.088	• Begin applications before populations begin to build or at economic thresholds according to local economic guidelines. • The low rate can be used early season for low pest densities or when tank mixing with other products labeled for target insect control on cotton. • Use the high rate for large pest populations, dense foliage, and longer residual. • Rapidly growing cotton may need retreatment. Scout fields often and retreat as necessary to maintain pest populations below damaging levels.

SPRAY RECOMMENDATIONS

Thorough spray coverage of plant foliage is essential for optimum control. Use a minimum of 5 gallons per acre by ground and 3 gallons per acre by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.

USE RESTRICTIONS

- Do not apply more than 2.8 oz/acre (0.088 lb. ai/acre) of **CablON** per single application.
- Do not apply more than 3 applications at the single maximum application rate of 2.8 oz/acre (0.088 lb. ai/acre) of **CablON** per year.
- Do not apply more than 8.4 oz/acre (0.263 lb. ai/acre) of **CablON** or other flonicamid-containing products per year.
- Allow a minimum of 7 days between applications.

Consult your local Atticus, LLC representative, extension service specialist, certified crop advisor, or state agricultural experiment station for any additional local use recommendations for your area.



Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513