

SALIXTM

2.6 F

Contains Tebuconazole, the active ingredient used in Raxil® 2.6F Seed Treatment Fungicide.



**SEED
TREATMENT**

**FOR USE ONLY BY COMMERCIAL SEED TREATERS
FOR CONTROL OF SPECIFIED DISEASES ON BARLEY, CORN, OATS, TRITICALE AND WHEAT**

ACTIVE INGREDIENT: (% by weight)
Tebuconazole, Alpha-[2-(4-Chlorophenyl)ethyl]-alpha-(1, 1-dimethyl-ethyl)-1 H-1,2,4-triazole-1-ethanol 28.3%
OTHER INGREDIENTS: 71.7%
TOTAL 100.0%
Contains 2.6 lbs. tebuconazole per gallon
EPA Reg. No.: 91234-461

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you DO NOT understand the label, find someone to explain it to you in detail.)**

See below for additional Precautionary Statements and Directions for Use.

FIRST AID

If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by the poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at **1-844-685-9173** for emergency medical treatment information.

**For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

SalixTM 2.6 F is not manufactured, or distributed by Bayer CropScience, seller of Raxil®.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Avoid breathing (dust, vapor, or spray mist).

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt, long pants or coveralls
- Chemical-resistant gloves made of waterproof material

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory

Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT mix or allow coming in contact with oxidizing agent. Hazardous Chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on agricultural establishments in hopper-box, planter-box, slurry-box or other seed treatment applications at or immediately before planting.

PRODUCT INFORMATION

Salix 2.6 F controls or suppresses certain designated seed and soilborne diseases of barley, corn, oats, triticale and wheat.

SEED TREATMENT: Apply **Salix 2.6 F** as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment. Prepare a slurry by mixing the specified dosage in 9 to 20 fl. oz. (0.18 to 0.41 lb. ai) of water per 100 lbs. of seed. Add dye** to the resulting slurry. Mix the slurry thoroughly with the seeds to provide uniform coverage.

**21 CFR Chapter I, Section 2.25 requires grain seed treated with poisonous substances in excess of recognized tolerances be suitably colored with an EPA approved dye, such as one of the dyes listed in 40 CFR Section 180.1001 (c) and (d), to prevent their subsequent inadvertent use as a food for man or feed for animals.

RESISTANCE MANAGEMENT

The component in **Salix 2.6 F** belongs to resistance group 3 (see resistance group number on front panel of label). Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotating and/or tank mixing with products having different modes of action; or limiting the total number of applications per year. Use labeled rates. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide/insecticide resistance management and/or integrated pest management recommendations for specific crops and pest populations. Atticus, LLC encourages responsible product stewardship to ensure effective long-term control of pests on this label.

BARLEY

Diseases	Rate (fl. oz./100 lbs. of seed)	Directions for Use
Barley Stripe (suppression) Covered Smut* Early Season <i>Rhizoctonia</i> Root Rot Early Season Common Root Rot Early Season <i>Fusarium</i> Foot Rot* Early Season Suppression of Powdery Mildew Early Season Suppression of Barley Leaf Rust Loose Smut*	0.1 (0.002 lb. ai)	Refer to the Seed Treatment section on this label for the specified method of treating seed.

*Not Registered for Use by California.

CORN (SWEET CORN, FIELD CORN, FIELD CORN GROWN FOR SEED, AND POPCORN)*

Diseases	Rate (fl. oz./100 lbs. of seed)	Directions for Use
Soilborne and Seedborne Fusarium	0.075 – 0.10 (0.0015 – 0.0020 lb. ai)	Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed should be sound and well cured prior to treatment. Product should be diluted with sufficient water to ensure complete seed coverage. Add dye to the resulting slurry. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with Salix 2.6 F . The length of control will vary depending on the rate used.
Soilborne and Seedborne Head smut (<i>Sphacelotheca reiliana</i>)	0.37 – 0.74 (0.0075 – 0.015 lb. ai)	

*Not Registered for Use by California.

OATS

Diseases	Rate (fl. oz./100 lbs. of seed)	Directions for Use
Early Season <i>Rhizoctonia</i> Root Rot Early Season Common Root Rot Early Season <i>Fusarium</i> Foot Rot* Early Season Suppression of Powdery Mildew Early Season Suppression of Crown Rust Loose Smut* Pythium Damping-off	0.1 (0.002 lb. ai)	Refer to the Seed Treatment section on this label for the specified method of treating seed.

*Not Registered for Use by California.

WHEAT and TRITICALE

Diseases	Rate (fl. oz./100 lbs. of seed)	Directions for Use
Early Season Septoria Disease Complex Early Season <i>Rhizoctonia</i> Root Rot Early Season Common Root Rot Early Season <i>Fusarium</i> Foot Rot* Early Season Suppression of Powdery Mildew Early Season Suppression of Wheat Leaf Rust Flag Smut* Loose Smut* Stinking Smut*	0.1 (0.002 lb. ai)	Refer to the Seed Treatment section on this label for the specified method of treating seed.

*Not Registered for Use by California.

SEED BAG LABELING REQUIREMENTS

The Federal Seed Act requires that containers containing treated seeds shall be labeled with the following statements:

- This seed has been treated with **Salix 2.6 F**, a fungicide containing tebuconazole.
- **DO NOT** use treated seed for feed, food, or oil purposes.

The U.S. Environmental Protection Agency requires the following statements on containers containing seed treated with tebuconazole:

- Store treated seed away from food and feedstuffs.
- **DO NOT** allow children, pets or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- Barley, wheat and oat, green forage may be grazed or harvested for hay 31 days after seeding.
- **DO NOT** contaminate bodies of water when disposing of planting equipment washwater.
- Dispose of seed packaging or containers in accordance with local requirements.
- Excess treated seed may be used for ethanol production if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

SPECIMEN

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

For plastic containers ≤ 5 gallons: Nonrefillable Container: DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

For plastic containers > 5 gallons: Nonrefillable container: DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Salix™ is a trademark of Atticus, LLC.

Raxil® is a registered trademark of Bayer.

20260226a

