This information is for promotional purposes only. Space considerations may require information to be omitted.
Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.

Agnela™ FL

Contains Metalaxyl, the active ingredient used in Allegiance® FL Seed Treatment Fungicide.

SEED TREATMENT

For Control of Seed Rot and Damping-Off Diseases of Certain Crops

ACTIVE INGREDIENT:	**(% by weight)**
Metalaxyl (CAS No.57837-19-1)	28.98%
OTHER INGREDIENTS:	71.02%
TOTAL	100.00%

Contains 2.65 lbs. metalaxyl per gallon

EPA Reg. No.: 91234-464

KEEP OUT OF REACH OF CHILDREN

WARNING • AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you DO NOT understand the label, find someone to explain it to you in detail.)

See below for additional Precautionary Statements and Directions for Use.

SPECIMEN

FIRST AID	
If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • **DO NOT** induce vomiting unless told to do so by the poison control center or doctor. • **DO NOT** give anything by mouth to an unconscious person.
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at **1-844-685-9173** for emergency medical treatment information.	

For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

Agnela™ FL is not manufactured, or distributed by Bayer CropScience, seller of Allegiance® FL.

Atticus

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes skin irritation. Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes. **DO NOT** get on skin or on clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Socks and chemical-resistant footwear;
- Chemical-resistant gloves (barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, and/or Viton ≥ 14 mils).
- When mixing and loading, and when cleaning equipment, wear a chemical-resistant apron.

NOTE: Persons involved in bagging treated seed, sewing or moving bags of treated seed, or cleaning up bagging areas or seed treatment equipment are pesticide handlers and must wear the PPE required on this label for pesticide handlers.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and change into clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing them. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** apply when weather conditions favor drift from treated areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

GROUNDWATER ADVISORY: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

SPECIMEN

PHYSICAL OR CHEMICAL HAZARDS

DO NOT mix or allow coming in contact with Oxidizing agents. Hazardous Chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all label directions carefully before use.

DO NOT apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 24 hours**.

Exception: Once the seeds are planted in soil, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no worker contact with the soil sub-surface or treated seed.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Socks and chemical-resistant footwear
- Chemical-resistant gloves
- Protective eyewear

PRODUCT INFORMATION

Agnela FL is a systemic fungicide seed treatment specifically for control of systemic downy mildews, *Pythium* and *Phytophthora* spp. For control of other soilborne diseases, use in combination with other seed treatment fungicides.

DO NOT use **Agnela FL** in combination with other seed treatment products unless compatibility and safety to the crop has been verified. **DO NOT** mix with any pesticide that has a prohibition against tank-mixing.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Agnela FL may be applied on its own, as water-based slurry, or in combination with other registered seed treatment insecticides and fungicides through standard slurry or mist- type commercial seed treatment equipment.

NOTE: **Agnela FL** is a systematic fungicide with a specific mode of action. **Agnela FL** could be subject to development of resistant stains of fungi. Development of resistance cannot be predicted. Therefore, the company cannot assume liability for crop damage resulting from resistant strains of fungi. Consult your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible **Agnela FL** resistant strains of fungi which may occur.

SPECIMEN

FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

USE RESTRICTIONS

- Seed treated commercially or for sale must be dyed an un-natural color using an EPA approved colorant before being sold. When using formulations that **DO NOT** contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.
- **IN CALIFORNIA:** ONLY for use in commercial seed-treatment establishments. NOT allowed for "on-farm" use in California.
- **IMPORTANT:** MAXIMUM USAGE when applying multiple metalaxyl and mefenoxam containing products to the same crop within the same season: **DO NOT** apply more than the maximum seasonal total for the active ingredient as stated on the label of the lowest seasonal total on that crop.
- **DO NOT** use other metalaxyl or mefenoxam products for control of Pythium in water-seeded rice.
- **DO NOT** apply **Agnela FL** to spinach which is destined to be grown in the greenhouse.

As stated by regulations specified in 40 CFR 153.155, all seed treated with **Agnela FL** must be colored with an EPA-approved dye or colorant to visually distinguish seed treated with a pesticide. This product contains such dye or colorant. This regulation is necessary to prevent treated seed from accidental use as a food for man or feed for animals.

- **DO NOT** use treated seed for feed, food, or oil purposes.
- Store treated seed away from feeds and foodstuffs.
- Seed treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- If excess treated seed remains after planting, dispose of it by burying away from bodies of water.
- The maximum annual application rate for metalaxyl or mefenoxam must not exceed 12.3 lb. ai per acre per calendar year or the lesser value listed in the crop section.
- **DO NOT** harvest mustard greens. Rape green seeds grown and harvested from **Agnela FL** treated seed are not for human or feed consumption. Rapeseed grown and harvested from **Agnela FL** treated seed is only for industrial uses and cannot be used for edible oil or any other human consumption or animal feed.

Atticus

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, **Agnela FL** contains a Group 4 fungicide. Any fungal population may contain individuals naturally resistant to **Agnela FL** and other Group 4 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **Agnela FL** or other Group 4 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models along is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance management and/or IPM recommendations for specific crops and pathogens.
- For information or to repost suspected resistance, contact Atticus, LLC. You can also contact your pesticide distribution or university extension specialist to report resistance.

SEED CONTAINER LABEL REQUIREMENTS

The Federal Seed Act requires that containers containing treated seeds shall be labeled with the following statements:

- This seed has been treated with metalaxyl fungicide.
- **DO NOT** use treated seed for feed, food or oil purposes.

The U.S. Environmental Protection Agency requires the following statements on containers containing seed treated with metalaxyl:

- Store treated seed away from food and feedstuffs.
- When opening this bag or loading/pouring the treated seed/seed-pieces, wear long sleeved shirt, long pants, shoes, socks, chemical-resistant gloves.
- Excess treated seed may be used for ethanol production only if
  (1) by-products are not used for livestock feed and
  (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- **DO NOT** contaminate water bodies when disposing of planting equipment wash waters.
- **DO NOT** allow children, pets or livestock to have access to treated seeds.
- **DO NOT** use other metalaxyl or mefenoxam products for control of pythium in water-seeded rice.
- **DO NOT** apply metalaxyl to spinach which is destined to be grown in the greenhouse.
- The maximum annual application rate for metalaxyl or mefenoxam must not exceed 12.3 lb. ai/A.

SPECIMEN

Groundwater Advisory: Metalaxyl has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. Therefore, contain any product spills or equipment leaks and dispose of wastes according to the disposal instructions on this label.

APPLICATION INSTRUCTIONS

BEANS (EXCEPT SOYBEANS)

Field beans, green beans, chickpeas (garbanzos), kidney beans, lima beans, lupines, navy beans, okra, pole beans, snap beans, string beans, wax beans, and lentils.

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium seed rot and damping-off Early season *Phytophthora*	1,600	0.75 (0.016 lb. ai)	0.469 (0.001 lb. ai)	0.0439
Restriction: • **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.				
[1]Based on average seeds per lb.				

BRASSICA (COLE) LEAFY VEGETABLES

Broccoli (including Chinese and raab), Brussels sprouts, cabbage (including Chinese bok choy, Chinese napa and mustard), cauliflower, cavalo broccolo, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, and rape greens

Diseases Controlled	Application Rate (Fl. Oz./100 lbs. of Seed)
Pythium damping-off	0.75 (0.016 lb. ai)
Restriction: • **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.	

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

BUCKWHEAT

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium damping-off	13,000	0.75 (0.016 lb. ai)	0.058 (0.0010 lb. ai)	0.0054

Restriction:
- **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.

[1]Based on average seeds per lb.

CANOLA*, MUSTARD SEED*, and RAPESEED*

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium damping-off	100,000	0.25 – 0.50 (0.005 – 0.01 lb. ai)	0.0025 – 0.005 (0.000052 – 0.0001 lb. ai)	0.00023 – 0.00047

Restriction:
- **DO NOT** harvest mustard greens. Rape green seeds grown and harvested from **Agnela FL** treated seed are not for human or feed consumption. Rapeseed grown and harvested from **A216.50** treated seed is only for industrial uses and cannot be used for edible oil or any other human consumption or animal feed.
- **DO NOT** apply more than 0.50 fl. oz. (0.01 lb. ai) per 100 lbs. of seed in a single application. [1]Based on average seeds per lb.

[1]Based on average seeds per lb.

*Not Registered for Use by California.

CEREAL GRAINS
Wheat, Barley, Oats, Rye and Triticale

Diseases Controlled	Crop	Application Rate			
		Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium damping-off	Wheat, Barley, Oats	13,000	0.75 (0.016 lb. ai)	0.058 (0.0012 lb. ai)	0.0054
	Rye, Triticale	18,850	0.75 (0.016 lb. ai)	0.040 (0.0008 lb. ai)	0.0037

Restriction:
- **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.

[1]Based on average seeds per lb.

SPECIMEN

COTTON

Diseases Controlled	Application Rate			
	Seeds/Lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium seed rot and damping-off	4,500	0.75 - 1.50 (0.016 – 0.032 lb. ai)	0.167 - 0.333 (0.0035 – 0.0069 lb. ai)	0.0156 - 0.0312

Restrictions:
- **DO NOT** apply more than 1.50 fl. oz. (0.032 lb. ai) per 100 lbs. of seed in a single application.
- **DO NOT** apply more than 0.2 fl. oz. (0.004 lb. ai) per acre per year by seed treatment.

[1]Based on average seeds per lb.

CORN, SWEET

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium damping-off	3,150	0.75 (0.016 lb. ai)	0.238 (0.0049 lb. ai)	0.0223
Systemic downy mildew	3,150	1.5 (0.032 lb. ai)	0.476 (0.0099 lb. ai)	0.0446

Restriction:
- **DO NOT** apply more than 1.50 fl. oz. (0.032 lb. ai) per 100 lbs. of seed in a single application.

[1]Based on average seeds per lb.

CORN, FIELD

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium damping-off	1,680	0.75 (0.016 lb. ai)	0.446 (0.0092 lb. ai)	0.0419

Restriction:
- **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.

[1]Based on average seeds per lb.

CORN, POP

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium damping-off	3,060	0.75 (0.016 lb. ai)	0.245 (0.0051 lb. ai)	0.0230

Restriction:
- **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.

[1]Based on average seeds per lb.

CUCUMBER

Diseases Controlled	Application Rate (Fl. Oz./100 lbs. of Seed)
Pythium damping-off	0.75 (0.016 lb. ai)

Restriction:
- **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.

CUCURBIT VEGETABLES

Chayote (fruit), Chinese waxgourd, citron melon, gherkin, edible gourd (includes hyotan, cucuza, Chinese okra, and hechima), Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), muskmelon (includes true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), pumpkin, summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), winter squash (includes butternut squash, calabaza, hubbard squash), Cucumis mixta, Cucumis pepo (includes acorn squash, spaghetti squash), watermelon (includes hybrids and/or varieties of Citrullus lanatus).

SPECIMEN

Diseases Controlled	Application Rate (Fl. Oz./100 lbs. of Seed)
Pythium damping-off	0.75 (0.016 lb. ai)

Restriction:
- **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.

DILL

Diseases Controlled	Application Rate (Fl. Oz./100 lbs. of Seed)
Pythium damping-off	0.75 (0.016 lb. ai)

Restriction:
- **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.

EDIBLE SOYBEANS

Diseases Controlled	Application Rate (Fl. Oz./100 lbs. of Seed)
Pythium damping-off	0.75 (0.016 lb. ai)

Restriction:
- **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.

FORAGE GRASSES

Grasses grown for hay, grazing or silage, corn (fodder or silage), sorghum (hay or silage), and small grain grown for hay, grazing or silage.

Diseases Controlled	Application Rate (Fl. Oz./100 lbs. of Seed)
Pythium damping-off	0.75 (0.016 lb. ai)
Restriction: • **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.	

FORAGE LEGUMES

Alfalfa, clover, lespedeza, beans (for forage), soybeans (for forage), soybean hay, peanuts (for forage), peanut hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch and velvet beans (for forage).

Diseases Controlled	Application Rate (Fl. Oz./100 lbs. of Seed)
Pythium damping-off Early season *Phytophthora*	0.75 – 1.5 (0.016 – 0.032 lb. ai)
Restriction: • **DO NOT** apply more than 1.5 fl. oz. (0.032 lb. ai) per 100 lbs. of seed in a single application.	

FRUITING VEGETABLES

Eggplants, ground cherry, pepino, pepper (including bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), tomatillo, and tomato

Diseases Controlled	Application Rate (Fl. Oz./100 lbs. of Seed)
Pythium damping-off	0.75 (0.016 lb. ai)
Restriction: • **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.	

LEAFY VEGETABLES

Amaranth (leafy, Chinese spinach, tampala), cardoon, celery (including Chinese), celtuce, chervil, chrysanthemum (edible-leaved and garland), corn salad, cress (garden and upland), dandelion, dock, endive, fennel (finochio), lettuce (head and leaf), orach, parsley, purslane (garden and winter), radicchio, rhubarb, and Swiss chard.

Diseases Controlled	Application Rate (Fl. Oz./100 lbs. of Seed)
Pythium damping-off	0.75 (0.016 lb. ai)
Restriction: • **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.	

SPECIMEN

MILLET

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium damping-off	80,000	0.75 (0.016 lb. ai)	0.009 (0.0002 lb. ai)	0.0009
Restriction: • **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.				
[1]Based on average seeds per lb.				

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

ONIONS
(DRY BULB AND GREEN)

Diseases Controlled	Application Rate (Fl. Oz./100 lbs. of Seed)
Pythium damping-off	0.75 (0.016 lb. ai)

Restriction:
- **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.

PEANUTS*

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium seed rot and damping-off	700	0.75 – 1.5 (0.016 – 0.032 lb. ai)	1.071 (0.022 lb. ai)	0.1004

Restriction:
- **DO NOT** apply more than 1.5 fl. oz. (0.032 lb. ai) per 100 lbs. of seed in a single application.

[1]Based on average seeds per lb.

*Not Registered for Use by California

PEAS
Black-eyed peas, cowpeas, field peas, garden peas

Diseases Controlled	Crop	Application Rate			
		Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium seed rot and damping-off Early season *Phytophthora*	Small size seeds	3,300	0.75 (0.016 lb. ai)	0.227 (0.0047 lb. ai)	0.0213
	Larger size seeds	2,100	0.75 (0.016 lb. ai)	0.361 (0.0075 lb. ai)	0.0338
Systemic downy mildew Pythium damping-off	Field Peas	3,300	1.5 (0.032 lb. ai)	0.455 (0.0094 lb. ai)	0.0426
	Garden Peas	2,080	1.5 (0.032 lb. ai)	0.721 (0.015 lb. ai)	0.0676

Restriction:
- **DO NOT** apply more than 1.5 fl. oz. (0.032 lb. ai) per 100 lbs. of seed in a single application.

[1]Based on average seeds per lb.

SPECIMEN

RICE*

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium damping-off	18,800	0.75 - 1.5 (0.016 – 0.032 lb. ai)	0.040 - 0.080 (0.0008 – 0.0017 lb. ai)	0.0037 - 0.0075

Restriction:
- **DO NOT** apply more than 1.5 fl. oz. (0.032 lb. ai) per 100 lbs. of seed in a single application.
- **DO NOT** use other metalaxyl- or mefenoxam-containing products in water-seeded rice. Containers or bags containing **Agnela FL** treated rice must be labeled with this restriction.

[1]Based on average seeds per lb.

*Not Registered for Use by California

ROOT AND TUBER VEGETABLES (EXCEPT POTATOES)

Arracacha, arrowroot, artichoke (Chinese and Jerusalem), beet (garden and sugar), burdock (edible), canna (edible), carrot, cassava (bitter and sweet), celery root, chayote, chervil, chicory, chufa, dasheen, ginger, ginseng, horseradish, leren, parsley (turnip-rooted), parsnip, radish (includes oriental dalkon), rutabaga, salsify (includes black and Spanish), skirret, sweet potato, tanier, turmeric, turnip, yam bean (jicama, manico pea), yam

Diseases Controlled	Application Rate (Fl. Oz./100 lbs. of Seed)
Pythium damping-off	0.75 (0.016 lb. ai)

Restriction:
- **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.

SUNFLOWERS
(Oil and Confection Type)

Diseases Controlled	Crop	Application Rate			
		Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Systemic downy mildew	Oil	7,000	1.5 – 3.0 (0.032 – 0.063 lb. ai)	0.214 – 0.429 (0.0044 – 0.0089 lb. ai)	0.0201 – 0.0402
	Confection	3,500	1.5 – 3.0 (0.032 – 0.063 lb. ai)	0.429 – 0.857 (0.0089 – 0.0177 lb. ai)	0.0402 – 0.0804

Restriction:
- **DO NOT** apply more than 3.0 fl. oz. (0.063 lb. ai) per 100 lbs. of seed in a single application.

[1]Based on average seeds per lb.

SORGHUM/MILO

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium damping-off	18,000	0.375 - 0.75 (0.008 - 0.016 lb. ai)	0.021 - 0.042 (0.0004 – 0.0009 lb. ai)	0.0020 - 0.0039
Systemic downy mildew (races susceptible to chemical control, and cultivars which possess sorghum downy mildew resistance)	18,000	0.75 - 1.5 (0.016 – 0.032 lb. ai)	0.042 - 0.083 (0.0009 – 0.0017 lb. ai)	0.0039 - 0.0078
Systemic downy mildew (races susceptible to chemical control, and cultivars which **DO NOT** possess sorghum downy mildew resistance)	18,000	1.5 - 3.0 (0.032 – 0.063 lb. ai)	0.083 - 0.167 (0.0017 – 0.003 lb. ai)	0.0078 - 0.0156

Application Instruction: Treat only high quality seed (at least 90% germination) with **Agnela FL**.

Restriction:
- **DO NOT** apply more than 3.0 fl. oz. (0.063 lb. ai) per 100 lbs. of seed in a single application.

[1]Based on average seeds per lb.

SPECIMEN

SOYBEANS

Diseases Controlled	Application Rate			
	Seeds/Lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./140,000 Seeds	Mg ai/Seed
Pythium damping-off and early season *Phytophthora*	2,800	0.75 – 1.50 (0.016 – 0.032 lb. ai)	0.38 - 0.75 (0.0079 - 0.016 lb. ai)	0.0251 - 0.0502

Restriction:
- **DO NOT** apply more than 1.5 fl. oz. (0.032 lb. ai) per 100 lbs. of seed in a single application.
- **DO NOT** apply by seed treatment more than 0.83 fl. oz. (0.0166 lb. ai) per acre per year.

[1]Based on average seeds per lb.

SPINACH[1]

Diseases Controlled	Application Rate (Fl. Oz./100 lbs. of Seed)
Pythium damping-off	0.1 - 0.75 (0.0021 – 0.016 lb. ai)

Restriction:
- **DO NOT** apply more than 0.75 fl. oz. (0.016 lb. ai) per 100 lbs. of seed in a single application.

[1]**DO NOT** apply to spinach seed to be grown in greenhouses. Containers or bags containing **Agnela FL** –treated spinach seed must be labeled with this restriction.

TURF GRASSES
On Golf Courses, Home Lawns, Reclamation, and Erosion Control Projects

Diseases Controlled	Application Rate (Fl. Oz./100 lbs. of Seed)
Pythium damping-off	1.5 (0.032 lb. ai)
Restriction: • **DO NOT** apply more than 1.5 fl. oz. (0.032 lb. ai) per 100 lbs. of seed in a single application.	

SEED TREATMENT DIRECTIONS
USE AT REDUCED RATES IN COMBINATION WITH OTHER FUNGICIDES

Agnela FL systemic fungicide seed dressing may be applied at the reduced rates shown in the table below and still provide good Pythium control when used in combination with broad spectrum seed treatment fungicides. Apply as a commercial seed treatment only in combination at EPA registered rates of broad-spectrum seed treatment fungicides for which compatibility and crop safety have been verified. **DO NOT** mix with any pesticide that has a prohibition against tank-mixing. **Agnela FL** may be applied on its own or as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment.

BEANS
Field, Kidney, Navy, Pinto Beans, Lentils

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium seed rot and damping-off	1,600	0.20 - 0.375 (0.004 – 0.008 lb. ai)	0.125 – 0.234 (0.0026 – 0.0048 lb. ai)	0.0117 – 0.0220
Restriction: • **DO NOT** apply more than 0.375 fl. oz. (0.008 lb. ai) per 100 lbs. of seed in a single application.				
[1]Based on average seeds per lb.				

CEREAL GRAINS
Wheat, Barley, Oats, and Rye

Diseases Controlled	Crop	Application Rate			
		Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Seed decay and damping-off caused by *Pythium*	Wheat, Barley, Oats	13,000	0.10 – 0.375 (0.002 – 0.008 lb. ai)	0.008 – 0.029 (0.0002 – 0.0006 lb. ai)	0.0007 – 0.0027
	Rye	18,850	0.10 – 0.375 (0.002 – 0.008 lb. ai)	0.005 – 0.020 (0.0001 – 0.0004 lb. ai)	0.0005 – 0.0019
Restriction: • **DO NOT** apply more than 0.375 fl. oz. (0.008 lb. ai) per 100 lbs. of seed in a single application.					
[1]Based on average seeds per lb.					

SPECIMEN

CORN, FIELD

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Seed decay and damping-off caused by *Pythium*	1,680	0.10 – 0.375 (0.002 – 0.008 lb. ai)	0.060 – 0.223 (0.0012 – 0.0046 lb. ai)	0.0056 – 0.0209
Restriction: • **DO NOT** apply more than 0.375 fl. oz. (0.008 lb. ai) per 100 lbs. of seed in a single application.				
[1]Based on average seeds per lb.				

Atticus

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

PEANUTS*

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Seed decay and damping-off caused by *Pythium*	700	0.20 - 0.375 (0.004 – 0.008 lb. ai)	0.286 - 0.536 (0.0059 – 0.0011 lb. ai)	0.0268 - 0.0502
Restriction: • **DO NOT** apply more than 0.375 fl. oz. (0.008 lb. ai) per 100 lbs. of seed in a single application.				
[1]Based on average seeds per lb.				

*Not Registered for Use by California

PEAS

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium seed rot and damping-off	3,200	0.20 - 0.375 (0.004 – 0.008 lb. ai)	0.063 – 0.117 (0.0013 – 0.0024 lb. ai)	0.0059 – 0.0110
Restriction: • **DO NOT** apply more than 0.375 fl. oz. (0.008 lb. ai) per 100 lbs. of seed in a single application.				
[1]Based on average seeds per lb.				

RICE*

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Seed decay and damping-off caused by *Pythium*	18,800	0.10 – 0.375 (0.002 – 0.008 lb. ai)	0.005 - 0.020 (0.0001 – 0.004 lb. ai)	0.0005 - 0.0019
Restriction: • **DO NOT** apply more than 0.375 fl. oz. (0.008 lb. ai) per 100 lbs. of seed in a single application.				
Note: DO NOT use other metalaxyl or mefenoxam containing products in water-seeded rice. Containers or bags containing **Agnela FL** treated rice must be labeled with this restriction.				
[1]Based on average seeds per lb.				

*Not Registered for Use by California

SPECIMEN

SOYBEANS

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./140,000 Seeds	Mg ai/Seed
Pythium damping-off and early season *Phytophthora*	2,800	0.20 - 0.375 (0.004 – 0.008 lb. ai)	0.10 – 0.19 (0.0021 – 0.0039 lb. ai)	0.0067 – 0.0126
Restriction: • **DO NOT** apply more than 0.375 fl. oz. (0.008 lb. ai) per 100 lbs. of seed in a single application.				
[1]Based on average seeds per lb.				

SUNFLOWERS

Diseases Controlled	Application Rate			
	Seeds/lb.[1]	Fl. Oz./100 lbs. of Seed	Fl. Oz./100,000 Seeds	Mg ai/Seed
Pythium seed rot and damping-off	5,500	0.10 – 0.375 (0.002 – 0.008 lb. ai)	0.018 – 0.068 (0.0004 – 0.0014 lb. ai)	0.0017 – 0.0064
Restriction: • **DO NOT** apply more than 0.375 fl. oz. (0.008 lb. ai) per 100 lbs. of seed in a single application.				
[1]Based on average seeds per lb.				

SEED TREATMENT DIRECTIONS - FOR EXPORT USE ONLY

For treatment of the following seed for export only to countries requiring treatment with metalaxyl before importation. The treater is responsible for confirming the acceptability of his treated seed in the country to which he is exporting.

Crop	Diseases Controlled	Application Rate (Fl. Oz./100 lbs. of seed)
Brassica (Cole) Leafy Vegetables	*Pythium* damping-off	0.75 – 1.5 (0.016 – 0.032 lb. ai)
Canola (cultivars, varieties and/or hybrids, including Brassica napus, Brassica rapa, and Brassica juncea)	*Pythium* damping-off	0.5 – 1.70 (0.001 – 0.036 lb. ai) (32 to 110 ml per 100 kg of seed)
Corn	*Pythium* damping-off	0.75 – 1.5 (0.016 – 0.032 lb. ai)
	Systemic downy mildew and *Pythium* damping-off	3.0 (0.063 lb. ai)
Cucurbit Vegetables	*Pythium* damping-off	0.75 (0.016 lb. ai)
Fruiting Vegetables	*Pythium* damping-off	0.75 – 1.5 (0.016 – 0.032 lb. ai)
Leafy Vegetables	*Pythium* damping-off	0.75 – 1.5 (0.016 – 0.032 lb. ai)
Millet	*Pythium* damping-off	0.75 – 1.5 (0.016 – 0.032 lb. ai)
	Systemic downy mildew and *Pythium* damping-off	2.5 (0.052 lb. ai)
Onions	*Pythium* damping-off	0.75 – 1.5 (0.016 – 0.032 lb. ai)
Peas	*Pythium* damping-off and early season *Phytophthora*	3.5 (0.074 lb. ai)
Root and Tuber Vegetables (except potatoes)	*Pythium* damping-off	0.75 – 1.5 (0.016 – 0.032 lb. ai)
Sorghum	*Pythium* damping-off	0.75 – 1.5 (0.016 – 0.032 lb. ai)
	Systemic downy mildew and *Pythium* damping-off	5.25 (0.11 lb. ai)
Sunflowers	Systemic downy mildew	9.0 (0.19 lb. ai)

SPECIMEN

TREATED SEED LABELING

Seed treated in accordance with the Export Use pattern must be labeled as follows:
"FOR EXPORT ONLY – NOT FOR DOMESTIC SALES OR USE".

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

For plastic containers ≤ 5 gallons: Nonrefillable Container: DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

For plastic containers > 5 gallons: Nonrefillable container: DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Agnela™ is a trademark of Atticus, LLC.
Allegiance® is a registered trademark of Bayer.

20260624a

SPECIMEN

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513