This information is for promotional purposes only. Space considerations may require information to be omitted.
Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.

AZOXYSTROBIN | GROUP 11 | FUNGICIDE

DRAYVN™

SEED TREATMENT

Broad-spectrum seed treatment fungicide for control of seed and seed-borne diseases.

ACTIVE INGREDIENT:	**(% by weight)**
Azoxystrobin*	9.6%
OTHER INGREDIENTS:	90.4%
TOTAL:	100.0%

* Contains 0.83 pounds of active ingredient per gallon.

EPA Reg. No.: 91234-444

SPECIMEN

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See below for additional Precautionary Statements.

FIRST AID	
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at **1-844-685-9173** for emergency medical treatment information.	

For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

Atticus

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants;
- Chemical resistant gloves made of any waterproof material
- Shoes and socks

In addition, planters of turf seeds must wear:

- a minimum of NIOSH-approved particulate filtering facepiece respirator with an N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter: OR a NIOSH-approved powered air purifying respirator with HE filters.

Respirator fit testing, medical qualification, and training using a program that conforms to OSHA's requirements (see 29 CFR Part 1910.134), employers must verify that any handler who uses a respirator is:

- Fit-tested and fit-checked,
- Trained, and
- Examined by a qualified medical practitioner to ensure physical ability to safely wear the style of respirator to be worn. A qualified medical practitioner is a physician or other licensed health care professional who will evaluate the ability of a worker to wear a respirator. The initial evaluation consists of a questionnaire that asks about medical conditions (such as a heart condition) that would be problematic for respirator use. If concerns are identified, then additional evaluations, such as a physical exam, might be necessary. The initial evaluation must be done before respirator use begins. Handlers must be reexamined by a qualified medical practitioner if their health status or respirator style or use-conditions change.

Upon request by local/state/federal/tribal enforcement personnel, employers must provide documentation demonstrating how they have complied with these requirements.

Aerial and/or chemigation application to sod is prohibited.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

The active ingredient, azoxystrobin, in this product can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This pesticide is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow coming in contact with Oxidizing agents. Hazardous Chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift.

Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Use on potato seed pieces is prohibited.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notifications to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material
- Shoes and socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTION ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

TREATMENT OF HIGHLY MECHANICALLY DAMAGED SEED, OR SEED OF KNOWN LOW VIGOR AND POOR QUALITY, MAY RESULT IN REDUCED GERMINATION AND/OR REDUCTION OF SEED AND SEEDLING VIGOR. TREAT AND CONDUCT GERMINATION TEST ON A SMALL PORTION OF SEED BEFORE COMMITTING THE TOTAL SEED LOT TO A SELECTED CHEMICAL TREATMENT. DUE TO SEED QUALITY CONDITIONS BEYOND THE CONTROL OF ATTICUS, LLC CROP PROTECTION, NO CLAIMS ARE MADE TO GUARANTEE GERMINATION OF CARRY-OVER SEED.

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

USE INFORMATION

Drayvn is a broad spectrum, preventative seed treatment fungicide with systemic properties recommended for the control of many important plant diseases. **Drayvn** may be applied in tank mixes, or sequentially with other registered, crop protection products. If using **Drayvn** in a tank mixture with other seed treatment products, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank-mix partner label. No label dosage may be exceeded and the most restrictive label precautions and limitations must be followed. This product must not be mixed with any product which prohibits such mixing.

All applications must be made according to the use directions that follow. Apply **Drayvn** as a seed treatment following the guidelines specified in the Directions for Use section of this label. If a rate range is specified, use higher rates of **Drayvn** when the disease pressure is expected to be high.

It is recommended that **Drayvn** be combined with a Pythium-active seed treatment such as Apron XL® to offer broad spectrum protection against the seed and seedling disease complex *(Rhizoctonia* spp. and *Pythium* spp.*)*.

RESISTANCE MANAGEMENT

This product contains azoxystrobin, a Group 11 fungicide. Group 11 fungicides inhibit the QoI (quinone outside) site within the electron transport system. When products with the same mode of action are used repeatedly, resistance can develop.

Drayvn must be used in accordance with set guidelines for preventing build-up of resistance. When available, other registered seed treatment fungicides that have a different mode of action than strobilurins must be tank mixed with **Drayvn**. However, these tank mixes must first be tested to ensure that the slurry treatment is safe on the stored or planted seed. **Drayvn** is a strobilurin fungicide, therefore continuous or sequential use of astrobilurin fungicide, within or during successive growing seasons must be avoided or reduced. **Drayvn** is not cross-resistant with other classes of fungicide which have different modes of action.

However, because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies such as:

- Adopt an IPM program for fungicide/bactericide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide/bactericide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact your local Atticus, LLC representative at 1-984-465-4800 or local extension specialist to report resistance.

MIXING INSTRUCTIONS

Apply **Drayvn** as a water-based slurry using seed treatment application equipment that will provide uniform coverage on the seed surface. Consult the manufacturer of the application equipment being used for calibration and operation procedures.

Drayvn mixes easily with water and other water-based seed treatments manufactured by Atticus, LLC. When mixing with products from other manufacturers, the compatibility should be tested prior to use by conducting a jar test: Mix all intended seed treatments with the appropriate amount of water in a clear glass container. Mix well and allow to sit for one hour. Remix and observe for incompatibility.

Mixing **Drayvn** with Tank-Mix Partners: Add 1/2 of the required water to the mix tank and turn on the agitation. Mechanical agitation is preferred. If using wettable powders add them first to clean water allowing them to completely disperse prior adding **Drayvn** or other products. Allow each tank-mix partner to completely disperse before adding the next product. Add the remaining amount of water and agitate.

Maintain agitation until the entire slurry mixture has been used. Allow seed to dry prior to bagging.

SEED CONTAINER LABEL REQUIREMENTS

The Federal law requires that containers containing treated seeds shall be labeled with the following statements:

- This seed has been treated with azoxystrobin seed treatment.
- Do not use treated seed for food, feed or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seeds treated with **Drayvn**:

- This seed has been treated with XX lbs of azoxystrobin per XX lbs of seed
- Groundwater Advisory: Azoxystrobin and a degradate of azoxystrobin are known to leach through soil into groundwater under certain conditions as a result of label use. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- Store treated seed away from food and feedstuffs.
- Do not allow children, pets or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Excess treated seed may be used for ethanol production only if
  (1) by-products are not used for livestock feed, and
  (2) no measurable residues of pesticide remain in the ethanol by-products that are used in agronomic practice.
- Dispose of all excess treated seed by burying seed away from bodies of water
- Do not contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging or containers in accordance with local requirements.

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

CROP SPECIFIC DIRECTIONS FOR USE

Crop	Target Diseases	Fl. oz./ 100 lbs. seed	Fl. oz./ 80,000 kernel count*	Mg ai/ Kernel**
Corn Field Pop Sweet (includes seed production)	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia* spp., *Penicillium* spp., *Pythium* spp.*)*	0.1 - 3.75	0.045 - 1.687	0.0016 - 0.0612
Application Directions: For optimum Pythium disease control, use **Drayvn** in combination with labeled rates of metalaxyl or mefenoxam products. Observe all precautions, limitations, rates, and directions for use on the respective labels before applying.				
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).				

* Based on 80,000 kernels of corn weighing 45 pounds.
** Based on 1,777 corn seeds per pound.

Crop	Target Diseases	Fl. oz./ 100 lbs. seed	Mg ai/ Kernel*
Sunflower	Downy mildew *(Plasmopora halstedii)* Seedling damping-off *(Rhizoctonia solani)*	3.75 - 15.0	0.025 - 0.10
For suppression of downy mildew			
Restriction: • Do not exceed 15 fl. oz./100 lbs. seed (0.000975 lbs. ai/lb. seed).			

* Based on 4,500 sunflower seeds per pound.

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Barley	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)*	0.10 - 3.75
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Brassica Leafy Greens subgroup: Broccoli raab Cabbage, Chinese Collards Kale Mizuna Mustard greens Mustard spinach Rape greens Head and Stem subgroup: Broccoli Brussels sprouts Cabbage Cauliflower Cavolo Broccolo Chinese broccoli Chinese cabbage (napa) Chinese mustard cabbage Kohlrabi	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)*	0.10 - 3.75
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Bulb Vegetables: Garlic Leek Onion, bulb Onion, green Welch onion Shallot	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)*	0.10 - 3.75
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Canola	Seed-borne Blackleg *(Phoma lingam)* Seedling Rhizoctonia Damping-off *(Rhizoctonia solani)* Alternaria Seedling Blight *(Alternaria* spp.*)*	0.10 - 3.75
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Cotton	Seedling damping-off *(Rhizoctonia solani)* Pythium seedling blight *(Pythium aphanidermatum)*	0.10 - 3.75
Where appropriate, use **Drayvn** in combination with a labeled metalaxyl or mefenoxam product.		
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Cucurbits: Cantaloupe Chayote Chinese waxgourd Cucumber Gourds Honeydew Melons *Momordica* spp. (bitter melon, balsam apple) Muskmelon Watermelon Pumpkin Squash Zucchini (Including cultivars and/or hybrids of these)	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)*	0.10 - 3.75
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Leafy Vegetables: Amaranth Arugula Cardoon Celery Celtuce Chervil Chrysanthemum, edible Coriander, leaves (Cilantro) Corn salad Cress Dandelion Dock Endive Fennel Lettuce, head and leaf Orach Parsley Purslane Radicchio Rhubarb Spinach Swiss Chard (Including cultivars and/or hybrids of these)	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)*	0.10 - 3.75
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Legume Vegetables: Bean *(Lupinus* spp.*)* (includes grain lupin, sweet lupin, white lupin, and white sweet lupin) Bean *(Phaseolus* spp.*)* (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean) Bean *(Vigna* spp.*)* (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean) Broad bean (fava bean) *(Vicia faba)* Chickpea (garbanzo bean) *(Cicer arietinum)* Guar *(Cyamopsis tetragonoloba)* Jackbean *(Canavalia ensiformis)* Lablab bean (hyacinth bean) *(Lablab purpureas)* Lentil *(Lens culinaris)* Pea *(Pisum* spp.*)* (includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea, Pigeon pea *(Cajanus cajan)*, Sword bean *(Canavalia gladiata)* For soybeans, see **Soybean** section for specific use directions.	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)*	0.10 - 3.75
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Peanut	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)* Aspergillus Crown Rot *(Aspergillus niger)* White Mold/Stem Rot *(Sclerotium rolfsii)*	0.10 - 3.75
For suppression of Aspergillus Crown Rot and White Mold/Stem Rot.		
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Pepper (Bell, Non-Bell, Sweet) Eggplant Okra	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)*	0.10 - 3.75
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Rice	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling Damping-Off *(Rhizoctonia solani)* *(Altemaria* spp.*)* *(Gaeumannomyces graminis* var. *graminis)* Brown Spot *(Bipolaris oryzae)* Rice Blast *(Pyricularia grisea)*	0.153 - 1.53
Application Directions: Where appropriate, use **Drayvn** in combination with labeled metalaxyl or mefenoxam products. For early season protection against seed-borne disease and early season rice blast.		
Restriction: • Do not exceed 1.53 fl. oz./100 lbs. seed (0.00010 lbs. ai/lb. seed).		

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Sorghum	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia* spp., *Penicillium* spp., *Pythium* spp.*)* Downy mildew *(Peronoscleroscera sorahi)*	0.308 - 3.08 (0.00062 - 0.0062 Mg ai/seed*)
Application Directions: Where appropriate, use **Drayvn** in combination with labeled metalaxyl or mefenoxam products. For suppression of downy mildew, use high rate of **Drayvn**.		
Restriction: • Do not exceed 3.08 fl. oz./100 lbs. seed (0.00020 lbs. ai/lb. seed).		

* Based on 14,500 seeds/lb.

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Soybeans	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia* spp., *Pythium* spp.*)* White Mold *(Sclerotium rolfsii)*	0.153 - 0.459
For suppression of white mold.		
Restriction: • Do not exceed 0.459 fl. oz./100 lbs. seed (0.0000298 lbs. ai/lb. seed).		

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Tomatoes	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)*	0.10 - 3.75
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Vegetables, leaves of root and tuber group: Beet, garden and sugar Burdock Carrot Cassava, bitter and sweet Celeriac (celery root) Chervil, turnip-rooted Chicory Dasheen (taro) Parsnip Radish Radish, oriental (daikon) Rutabaga Salsify, black Sweet potato Tanier Turnip Yam, true	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)*	0.10 - 3.75
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Vegetables, root subgroup: Beet, garden and sugar Burdock Carrot Celeriac Chervil, turnip-rooted Chicory Ginseng Horseradish Parsley, turnip-rooted Parsnip Radish Radish, oriental Rutabaga Salsify Salsify, black Salsify, Spanish Skirret Turnip	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)*	0.10 - 3.75
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Vegetables, tuberous and corm subgroup: Arracacha Arrowroot Artichoke, Chinese and Jerusalem Canna Cassava (edible, bitter and sweet) Chayote (root) Chufa Dasheen (Taro) Ginger Leren Sweet Potato Tanier Turmeric Yam, bean Yam, true	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)*	0.10 - 3.75
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Watercress	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)*	0.10 - 3.75
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Wheat Triticale	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)* Dwarf Bunt Common Bunt	0.10 - 3.75
Application Directions: For protection against common bunt and partial control of dwarf bunt. Where appropriate, use **Drayvn** in combination with labeled metalaxyl or mefenoxam products.		
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Ornamental Seed (Including crops listed in **Table 1**)	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)*	0.10 - 3.75
For early season protection against seed-borne disease and Rhizoctonia damping-off.		
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Turfgrass (Including bentgrass, bluegrass, bermudagrass, fescues grown for turf)	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)*	0.10 - 3.75
For early season protection against seed-borne disease and Rhizoctonia damping-off.		
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

Crop	Target Diseases	Fl. oz./ 100 lbs. seed
Bulbs, Corms, and Tubers of Ornamental Flowers and Foliage Plants	Seed-borne and soil-borne fungi causing decay, damping-off, and seedling blight Seedling damping-off *(Rhizoctonia solani)*	0.10 - 3.75
For early season protection against seed-borne disease and Rhizoctonia damping-off.		
Restriction: • Do not exceed 3.75 fl. oz./100 lbs. seed (0.0002432 lbs. ai/lb. seed).		

PLANT SAFETY: Drayvn has been shown to be safe when applied to the ornamental plants listed in **Table 1**. However, due to the large number of genera, species and varieties of ornamental and nursery plants, it is impossible to test every one for tolerance to **Drayvn**. Neither the manufacturer nor the seller has determined whether or not **Drayvn** can be used safely on genera, species, or varieties of ornamental and nursery plants not specified on this label. The professional user should conduct small scale testing to ensure plant safety prior to broadscale commercial use on plant genera and species not listed in this label.

Table 1. Plants listed by Common Name*

COMMON NAME	BOTANICAL NAME
Abelia	*Abelia* spp.
Andromeda, Japanese	*Pieris japonica*
Arborvitae	*Thuja* spp.
Aspen Trees	*Populus* spp.
Aster	*Aster* spp.
Aucuba,Japanese	*Aucuba japonica*
Azalea, Glacier	*Rhododendron* spp.
Azaleas	*Rhododendron* spp.
Balsam	*Impatiens* spp.
Barberry	*Berberis thunbergii*
Begonia (except Rieger begonia)	*Begonia* spp.
Birch, River	*Betula nigra*
Black-Eyed Susan	*Rudbeckia hirta*
Blanket-Flower	*Gaillardia* spp.
Bougainvillea	*Bougainvillea* spp.
Boxwood	*Buxus sempervirens*
Buddleia	*Buddleia davidii*
Bugle	*Ajuga reptans*
Bugleweed	*Ajuga reptans*

(continued)

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Table 1. Plants listed by Common Name* *(continued)*

COMMON NAME	BOTANICAL NAME
Burning Bush	*Euonymus alatus*
Butterfly Bush	*Buddleia davidii*
Cactus, Holiday	*Schlumbergera*
Caladium	*Caladium* spp.
Camellia	*Camellia japonica*
Carnation	*Dianthus caryophyllus*
Ceanothus	*Ceanothus* spp.
Cedar, Atlas	*Cedrus atlantica*
Cedar, Red	*Juniperus virginiana*
Cedar, White	*Cedrus* spp.
Chinese fountain grass	*Pennisetum alopecuroides*
Christmas Trees	See Fraserfir, Scotch pine and Douglas fir
Chrysanthemum	*Chrysanthemum* spp.
Cinquefoil	*Potentilla* spp.
Clethra	*Clethra alnifolia*
Coleus	*Plectranthus* spp.
Cotoneaster, Creeping	*Cotoneaster adpressus*
Cotoneaster, Variegated Rockspray	*Cotoneaster horizontalis*
Cranesbill	*Geranium* spp.
Crapemyrtle	*Lagerstroemia indica*
Cyclamen	*Cyclamen* spp.
Cyperus	*Cyperus* spp.
Cypress, Sawara	*Chamaecyparis pisifera*
Cypress, Leyland	*Chamaecyparis* spp.
Daisy, Gerber	*Gerbera jamesonii*
Daisy, Transvaal	*Gerbera jamesonii*
Dogwood	*Cornus* spp.
Dogwood	*Cornus florida*
Dogwood, Pink	*Cornus* spp.
Dumb-Cane	*Dieffenbachia* spp.
Euonymus, Dwarf Winged	*Euonymus alata*
Euonymus, Evergreen	*Euonymus japonicus*
Evergreen, Chinese	*Aglaonema* spp.
Fatsia, Japanese	*Fatsia japonica*
Fig	*Ficus* spp.
Fir, Douglas	*Pseudotsuga* spp.
Fir, Fraser	*Abies fraseri*
Floss-Flower	*Ageratum* spp.

(continued)

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Table 1. Plants listed by Common Name* *(continued)*

COMMON NAME	BOTANICAL NAME
Forsythia	*Forsythia viridissima*
Foxglove	*Digitalis* spp.
Gardenia	*Gardenia jasminoides*
Geranium	*Pelargonium* spp.
Grass, Dwarf Pampas	*Phalaris* spp.
Grass, Pampas	*Cortaderia selloana*
Hawthorn, Indian	*Rhaphiolepsis indica*
Heather	*Erica dareyensis*
Hemlock	*Tsuga* spp.
Hibiscus	*Hibiscus moscheutos*
Hibiscus	*Hibiscus rosa-sinensis*
Holly	*Ilex* spp.
Hosta	*Hosta* spp.
House-Leek	*Sempervivum* spp.
Hydrangea	*Hydrangea* spp.
Hydrangea, French	*Hydrangea macrophylla*
Impatiens	*Impatiens* spp.
Iris, African	*Dietes iridiodes*
Iris, Butterfly	*Dietes iridiodes*
Ivy, Algerian	*Hedera algeriensis*
Ivy, English	*Hedera helix*
Ivy, Swedish	*Plectranthus* spp.
Juniper	*Juniperus procumbens*
Juniper	*Juniperus scopulorum*
Juniper	*Juniperus* spp.
Larkspur	*Delphinium* spp.
Laurel	*Laurus nobilis*
Laurel, Australian	*Pittosporum* spp.
Laurel, Japanese	*Aucuba japonica*
Lilac, California	*Ceanothus* spp.
Lilac, Wild	*Ceanothus sanguineus*
Lily, Peace	*Spathiphyllum floribundium*
Lily-Turf	*Liriope muscari*
Live-Forever	*Sempervivum* spp.
Magnolia	*Magnolia* spp.
Magnolia, Saucer	*Magnolia soulangiana*
Magnolia, Southern	*Magnolia grandiflora*

(continued)

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Table 1. Plants listed by Common Name* *(continued)*

COMMON NAME	BOTANICAL NAME
Maple, Japanese	*Acer palmatum*
Maple, Sugar	*Acer saccharum*
Marigold	*Tagetes* spp.
Mock-Orange	*Pittosporum tobira*
Mugwort	*Artemisia* spp.
Nandina	*Nandina domestica*
Oak, Pin	*Quercus palustris*
Oak, Southern Red	*Quercus falcata*
Oleander	*Nerium oleander*
Orpine	*Sedum* spp.
Palm, Date	*Phoenix dactylifera*
Palm, Parlor	*Chamaedora elegans*
Palm, Queen	*Syagrus romanzoffianum*
Palm, Roebelin's	*Phoenix roebelenii*
Palm, Sago	*Caryota urens*
Pansy	*Viola* spp.
Paper-Plant	*Fatsia japonica*
Pear, Bradford's	*Pyrus calleryana*
Periwinkle	*Vinca* spp.
Petunia	*Petunia* spp.
Philodendron	*Philodendron* spp.
Phlox	*Phlox* spp.
Photinia, Red-Tip	*Photinia glabra*
Pine	*Pinus* spp.
Pine, Black	*Pinus nigra*
Pine, Eastern White	*Pinus strobus*
Pine, Muhgo	*Pinus muhgo*
Pine, Scotch	*Pinus sylvestris*
Pink	*Dianthus* spp.
Poinsettia	*Euphorbia* spp.
Pothos	*Epipremnum* spp.
Primrose	*Primula* spp.
Pussy's-Foot	*Ageratum* spp.
Redbud, Western	*Cercis occidentals*
Rhododendron	*Rhododendron* spp.
Ribbon-Grass	*Setaria* spp.
Rose of Sharon	*Hibiscus syriacus*
Rose	*Rosa* spp.

(continued)

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Table 1. Plants listed by Common Name* *(continued)*

COMMON NAME	BOTANICAL NAME
Rose-Bay	*Nerium oleander*
Rosemary (Prostrate)	*Rosmarinus* spp.
Rubber-Plant, Baby	*Peperomia* spp.
Rubber-Tree	*Schefflera actinophylla*
Sage	*Salvia* spp.
Sagebrush	*Artemisia* spp.
Snap-Dragon	*Antirrhinum* spp.
Snowball	*Ceanothus* spp.
Spirea	*Spirea budalda*
Spirea	*Spirea japonica*
Spruce, Blue	*Picea pungens*
Spruce, Norway	*Picea abies*
Spruce, White	*Picea glauca*
Starwort	*Aster* spp.
Stonecrop	*Sedum* spp.
Sweet Alyssum	*Lobularia maritima*
Thyme, Creeping	*Thymus serphyllum*
Umbrella-Tree	*Schefflera actinophylla*
Verbena	*Verbena* spp.
Vervain	*Verbena* spp.
Viburnum	*Viburnum* spp.
Vinca	*Catharanthus roseus*
Viola	*Viola* spp.
White alder	*Alnus rhombifolia*
Wiegela, Pink	*Wiegela florida*
Willow, Virginia	*Itea virginica*
Winterberry	*Ilex* spp.
Wormwood	*Artemisia* spp.
Yaupon	*Ilex* spp.
Yew, Spreading	*Taxus cuspidata* spp.
Yucca	*Yucca* spp.
Zebra-Plant	*Aphelandra* spp.
Zinnia	*Zinnia* spp.

* Do not apply **Drayvn** to *Malus* spp., *Prunus* spp., or leatherleaf fern (*Rumohra adianformis* and other species).

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

For plastic containers ≤ 5 gallons: Nonrefillable Container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

For plastic containers > 5 gallons: Nonrefillable container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Drayvn™ is a trademark of Atticus, LLC.

20260930a

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513