This information is for promotional purposes only. Space considerations may require information to be omitted.
Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.

QUINOXYFEN | GROUP 13 | FUNGICIDE

Quitora™

Contains Quinoxyfen, the active ingredient used in Quintec® Fungicide.

A Protectant Fungicide for the Control of Powdery Mildew in Artichoke, Gourds, Lettuce, Melons, Pepper, Pumpkin, Stone Fruits, Strawberry, Winter Squash, Cherry, Grapes, and Hops.

ACTIVE INGREDIENT:	**(% by weight)**
Quinoxyfen: 5,7-dichloro-4-(p-fluorophenoxy) quinoline	22.58%
OTHER INGREDIENTS:	77.42%
TOTAL	100.00%

Contains 2.08 lbs. of active ingredient per gallon.

EPA Reg. No.: 91234-438

SPECIMEN

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you DO NOT understand the label, find someone to explain it to you in detail.)

See below for additional Precautionary Statements.

FIRST AID	
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • **DO NOT** induce vomiting unless told to by a poison control center or doctor. • **DO NOT** give anything to an unconscious person.
If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at **1-844-685-9173** for emergency medical treatment information.	

For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTREC Day or Night Within USA and Canada:
1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

Quitora™ is not manufactured, or distributed by Gowan Company, LLC, seller of Quintec® Fungicide.

Atticus

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, or Viton ≥14 mils
- Shoes plus socks

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. Pilots must use an enclosed cab that meets the definition listed in the WPS for agricultural pesticides [40 CFR 170.305].

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** apply when weather conditions favor runoff or drift from the target area. **DO NOT** contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT mix or allow coming in contact with oxidizing agents, hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all **DIRECTIONS FOR USE** carefully before applying.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

DO NOT apply this product one-half hour before dusk to one-half hour after dawn in counties east of the Rocky Mountains where the gray bat, Indiana bat, and the Ozark big-eared bat occur.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, or Viton ≥14 mils
- Shoes plus socks

PRODUCT INFORMATION

Quitora is a protectant fungicide for the control of powdery mildew diseases. **Quitora** does not control existing or latent powdery mildew infections and must be applied on a protectant schedule (latency is the period between infection of plant tissues and the development of visible symptoms).

Quitora is compatible with most commonly used insecticides, miticides and fungicides.

USE RESTRICTIONS

- **DO NOT** graze or harvest forage from cover crops within areas treated with **Quitora**.
- **DO NOT** aerially apply this product on stone fruits, cherry, grapes, or hops. Aerial application is permitted on other crops for which this product is registered for use.
- **Chemigation: DO NOT** apply this product through any type of irrigation system.
- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.
- **Rainfast Requirement:** When applied alone, **Quitora** is rainfast 1 hour after application.
- Not for use in greenhouses or nurseries.

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted at intervals defined below following the final application of **Quitora**.

Crops	Re-Planting Interval
Registered Crops	No Restrictions
All Other Crops Grown for Food or Feed	30 days

SPECIMEN

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

MIXING DIRECTIONS

Quitora may be applied in tank mix combination with labeled rates of other pesticides, including fungicides, insecticides, or other chemicals used in the culture of the labeled crop provided:

(1) the tank mix product is labeled for the timing and method of application for the intended use; and

(2) tank mixing with **Quitora** or other pesticides is not prohibited by the label of the tank mix product. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Fill the spray tank to at least one-half the required spray volume and start agitation. Add (in order) dry flowable, wettable powder, and **Quitora** or other flowable (aqueous suspension) products to the spray tank. Then add the specified rate of emulsifiable concentrate and solution products. Allow time for thorough mixing after adding each product. Finish filling spray tank and maintain continuous agitation in the spray tank during mixing, filling and spraying operations.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Oil Sprays

Oil sprays may be integrated as part of a powdery mildew management program. Consult local university extension for specific information.

Use of Surfactants

A moderate rate of a registered nonionic or organosilicone surfactant may improve crop coverage. Penetrants and stickers are not advised.

Tank Mixing Directions

- Read carefully and follow all applicable use directions including application rates, use precautions, and limitations on the respective product labels.
- **DO NOT** exceed specified application rates or maximum use rates if products with the same active ingredient are applied together.
- For tank mix partners packaged in water soluble packaging, **DO NOT** use with products containing boron or mix in equipment previously used to apply boron-containing mixtures unless the tank and spray equipment have been adequately cleaned.
- Always perform a (jar) test to ensure compatibility of products used in tank mixtures.

Tank Mix Compatibility Testing

To determine the physical compatibility of **Quitora** with other pesticides, perform a jar test. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jells, oily films or layers, or other precipitates, it is not compatible and the tank mix combination must not be used.

SPRAY DRIFT MANAGEMENT

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 feet above the vegetative canopy unless a greater application height is necessary for pilot safety.
- The boom length must not exceed 75% of the wingspan for airplanes or 90% of the rotor blade diameter for helicopters.
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Airblast Applications:

- Sprays must be directed into the canopy.
- User must turn off outward pointing nozzles at row ends and when spraying outer rows.

Ground Boom Applications:

- User must apply with nozzle height specified by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

The applicator is responsible for avoiding off-site spray drift. Be aware of nearby non-target sites and environmental conditions.

Importance of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- **Volume** – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** – Use the lowest spray pressure specified for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- **Adjust Nozzles** – Follow nozzle manufacturer's specifications for setting up nozzles. To reduce fine droplets, orient nozzles parallel with the airflow in flight.

Boom Height – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, ensure that the boom remains level with the crop and has minimal bounce.

Release Height – Aircraft

Higher release heights increase the potential for spray drift.

Shielded Sprayers

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

SPECIMEN

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Temperature and Humidity

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

Wind

Drift potential increases with wind speed. Avoid applications during gusty wind conditions. Applicators need to be familiar with local wind patterns and terrain that could affect spray.

Boom-less Ground Application:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

RESISTANCE MANAGEMENT

For resistance management, **Quitora** contains a Group 13 fungicide. Any fungal population may contain individuals naturally resistant to **Quitora** and other Group 13 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance management strategies must be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **Quitora** or other Group 13 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Apply no more than two (2) sequential applications unless otherwise stated in the crop section.
- Apply early to keep fungal populations low.
- Avoid the consecutive use of **Quitora** or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- To help manage fungicide resistance, **DO NOT** use **Quitora** for commercial transplant production.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM guidelines for specific crops of pathogens.
- For information or to report suspected resistance, contact Atticus, LLC at 984-465-4800. You can also contact your pesticide distributor or university extension specialist to report resistance.

Disease Resistance Management

Cherry, Grapes, Hops, Stone Fruits: Quitora is in mode of action Group 13. **DO NOT** make more than five applications of **Quitora** per calendar year, nor more than two consecutive applications of **Quitora** before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply an alternate product within 14 days of applying **Quitora**.

Gourds, Lettuce, Melons, Pepper, Pumpkin, Strawberry, and Winter Squash: Quitora is in mode of action Group 13. **DO NOT** make more than four applications of **Quitora** per crop nor more than two consecutive applications of **Quitora** before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply an alternate product within 14 days of applying **Quitora**. It is advised that the total number of group 13 fungicide sprays per crop must not exceed 50% of the total number of powdery mildew sprays.

APPLICATION DIRECTIONS

APPLICATION TIMING

Apply **Quitora** before visible symptoms of powdery mildew appear. **Quitora** will not control latent or established infections of powdery mildew. If powdery mildew infection is established, apply **Quitora** in tank mix combination with a curative fungicide or following application of a curative spray or, if multiple diseases are present, following application of a broad-spectrum fungicide. Under low disease conditions (as per a predictive modeling system), minimum label use rates per application can be used. For optimum results, use maximum label rates and short application intervals for severe or threatening disease conditions.

When applied as a dilute spray, maintain the per acre use rate.

SPECIMEN

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

ARTICHOKE

Disease	Quitora Rate (Fl. Oz./Acre)	Re-Treatment Interval (Days)	Application Method
Powdery Mildew *(Leveillula taurica)*	4.0 - 6.0 (0.065 - 0.098 lb. ai)	10 - 14	**Ground:** For thorough spray coverage and optimum disease control, apply **Quitora** by ground application. Apply in a minimum spray volume of 30 gallons per acre. **Aerial:** Apply in a minimum spray volume of 5 gallons of water per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with **Quitora**. **DO NOT** apply under conditions when uniform spray coverage cannot be obtained or when spray drift may occur.

Precautions:
- **Quitora** is in mode of action Group 13. **DO NOT** make more than two consecutive applications of **Quitora** before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply an alternate product within 14 days of applying **Quitora**.
- Carefully inspect the crop several days after each application of **Quitora** and at regular intervals during crop development.
- Under certain environmental conditions, **Quitora** may cause leaf spotting and chlorosis. If these symptoms occur after applying **Quitora**, discontinue its use.

Restrictions:
- **Maximum Single Application Rate:** 6 fl. oz./A (0.098 lb. ai/A)
- **Maximum Annual Application Rate: DO NOT** apply more than 30 fl. oz. (0.49 lb. ai) of **Quitora** per acre per calendar year.
- **Maximum Applications/Year:** 5
- **Minimum Re-treatment Interval:** 10 days
- **Preharvest Interval:** None

CHERRY

Disease	Quitora Rate (Fl. Oz./Acre)	Re-Treatment Interval (Days)	Application Method
Powdery Mildew *(Podosphaera clandestina)*	8.7 (0.141 lb. ai)	10 - 14	**Ground:** Apply uniformly by ground application in a minimum spray volume of 30 gallons per acre. Disease control may be reduced at low spray volumes or if spray coverage is inadequate. **DO NOT** apply under conditions when uniform spray coverage cannot be obtained or when spray drift may occur. **Aerial: DO NOT** aerially apply this product on cherry.

Precautions:
- **Quitora** is in mode of action Group 13. **DO NOT** make more than two applications of **Quitora** to cherries per calendar year, nor more than two consecutive applications of **Quitora** before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply an alternate product within 14 days of applying **Quitora**.

Restrictions:
- **Maximum Single Application Rate:** 8.7 fl. oz./A (0.141 lb. ai/A)
- **Maximum Annual Application Rate: DO NOT** apply more than 17.4 fl. oz. (0.283 lb. ai) of **Quitora** per acre per calendar year.
- **Maximum Applications/Year:** 2
- **Minimum Re-treatment Interval:** 10 days
- **Preharvest Interval:** 7 days

SPECIMEN

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

GRAPES (SUBGROUP 13-07F)

Including: amur river grape, gooseberry, grape, kiwifruit, hardy, maypop, schisandra berry, cultivars, varieties and/or hybrids of these.

Disease	Quitora Rate (Fl. Oz./Acre)	Re-Treatment Interval (Days)	Application Method
Powdery Mildew *(Erysiphe necator)*	4.0 (0.065 lb. ai)	14	**Ground:** Apply uniformly by ground application in a minimum spray volume of 30 gallons per acre. Disease control may be reduced at low spray volumes or if spray coverage is inadequate. **DO NOT** apply under conditions when uniform spray coverage cannot be obtained or when spray drift may occur. **Aerial: DO NOT** aerially apply this product on grapes.
	6.6 (0.107 lb. ai)	21	

Precautions:

- **Quitora** is in mode of action Group 13. **DO NOT** make more than five applications of **Quitora** per calendar year, nor more than two consecutive applications of **Quitora** before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply an alternate product within 14 days of applying **Quitora**.

Restrictions:

- **Maximum Single Application Rate:** 6.6 fl. oz./A (0.107 lb. ai/A)
- **Maximum Annual Application Rate: DO NOT** apply more than 33 fl. oz. (0.536 lb. ai) of **Quitora** per acre per calendar year.
- **Maximum Applications/Year:** 5
- **Preharvest Interval:** 21 days

HEAD AND LEAF LETTUCE

Disease	Quitora Rate (Fl. Oz./Acre)	Re-Treatment Interval (Days)	Application Method
Powdery Mildew *(Erysiphe cichoracearum)*	4.0 - 6.0 (0.065 - 0.098 lb. ai)	10 - 14	**Ground:** For thorough spray coverage and optimum disease control, apply **Quitora** by ground application. Apply in a minimum spray volume of 30 gallons per acre. **Aerial:** Apply in a minimum spray volume of 5 gallons of water per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with **Quitora**. **DO NOT** apply under conditions when uniform spray coverage cannot be obtained or when spray drift may occur.

SPECIMEN

Precautions:

- **Quitora** is in mode of action Group 13. **DO NOT** make more than four applications of **Quitora** per crop, nor more than two consecutive applications of **Quitora** before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply an alternate product within 14 days of applying **Quitora**. It is advised that the total number of group 13 fungicide sprays per crop must not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each application of **Quitora** and at regular intervals during crop development.
- Under certain environmental conditions, **Quitora** may cause leaf spotting and chlorosis. If these symptoms occur after applying **Quitora**, discontinue its use.

Restrictions:

- **Maximum Single Application Rate:** 6 fl. oz./A (0.098 lb. ai/A)
- **Maximum Annual Application Rate: DO NOT** apply more than 24 fl. oz. (0.390 lb. ai) of **Quitora** per acre per crop.
- **Maximum Applications/Year:** 4
- **Minimum Re-treatment Interval:** 10 days
- **Preharvest Interval:** 1 day

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

HOPS

Disease	Quitora Rate (Fl. Oz./Acre)	Re-Treatment Interval (Days)	Application Method
Powdery Mildew *(Podosphaera macularis)*	4.0 - 8.2 (0.065 - 0.134 lb. ai)	14	**Ground:** Apply uniformly by ground application in a minimum spray volume of 30 gallons per acre. Disease control may be reduced at low spray volumes or if spray coverage is inadequate. **DO NOT** apply under conditions when uniform spray coverage cannot be obtained or when spray drift may occur. **Aerial: DO NOT** aerially apply this product on hops.

Precautions:
- **Quitora** is in mode of action Group 13. **DO NOT** make more than four applications of **Quitora** to hops per calendar year, nor more than two consecutive applications of **Quitora** before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply an alternate product within 14 days of applying **Quitora**.

Restrictions:
- **Maximum Single Application Rate:** 8.2 fl. oz./A (0.133 lb. ai/A)
- **Maximum Annual Application Rate: DO NOT** apply more than 32.8 fl. oz. (0.53 lb. ai) of **Quitora** per acre per calendar year.
- **Maximum Applications/Year:** 4
- **Preharvest Interval:** 21 days

MELONS (SUBGROUP 9A)

Including citron melon, muskmelon (cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon, true cantaloupe), watermelon

Disease	Quitora Rate (Fl. Oz./Acre)	Re-Treatment Interval (Days)	Application Method
Powdery Mildew *(Erysiphe cichoracearum) (Sphaerotheca fuliginea)*	4.0 - 6.0 (0.065 - 0.098 lb. ai)	10 - 14	**Ground:** For thorough spray coverage and optimum disease control, apply **Quitora** by ground application. Apply in a minimum spray volume of 30 gallons per acre. **Aerial:** Apply in a minimum spray volume of 5 gallons of water per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with **Quitora**. **DO NOT** apply under conditions when uniform spray coverage cannot be obtained or when spray drift may occur.

SPECIMEN

Precautions:
- **Quitora** is in mode of action Group 13. **DO NOT** make more than four applications of **Quitora** per crop, nor more than two consecutive applications of **Quitora** before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply an alternate product within 14 days of applying **Quitora**. It is advised that the total number of group 13 fungicide sprays per crop must not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each application of **Quitora** and at regular intervals during crop development. Under certain environmental conditions, **Quitora** may cause leaf spotting and chlorosis. If these symptoms occur after applying **Quitora**, discontinue its use.

Restrictions:
- **Maximum Single Application Rate:** 6 fl. oz./A (0.098 lb. ai/A)
- **Maximum Annual Application Rate: DO NOT** apply more than 24 fl. oz. (0.390 lb. ai) of **Quitora** per acre per crop.
- **Maximum Applications/Year:** 4
- **Minimum Re-treatment Interval:** 10 days
- **Preharvest Interval:** 3 days

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

PEPPER (SUBGROUP 8-10B)

Including: African eggplant; bell pepper, chili pepper, cooking pepper, eggplant, Martynia, non-bell pepper, okra, pea eggplant, pepino, pimento, roselle, scarlet eggplant, sweet pepper, cultivars, varieties, and/or hybrids of these

Disease	Quitora Rate (Fl. Oz./Acre)	Re-Treatment Interval (Days)	Application Method
Powdery Mildew *(Leveillula taurica)* *(Oidiopsis tuarica)*	4.0 - 6.0 (0.065 - 0.098 lb. ai)	10 - 14	**Ground:** For thorough spray coverage and optimum disease control, apply **Quitora** by ground application. Apply in a minimum spray volume of 30 gallons per acre. **Aerial:** Apply in a minimum spray volume of 5 gallons of water per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with **Quitora**. **DO NOT** apply under conditions when uniform spray coverage cannot be obtained or when spray drift may occur.

Precautions:

- **Quitora** is in mode of action Group 13. For Powdery Mildew, **DO NOT** make more than four applications of **Quitora** per crop, nor more than two consecutive applications of **Quitora** before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply an alternate product within 14 days of applying **Quitora**. It is advised that the total number of group 13 fungicide sprays per crop must not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each application of **Quitora** and at regular intervals during crop development. Under certain environmental conditions, **Quitora** may cause leaf spotting and chlorosis. If these symptoms occur after applying **Quitora**, discontinue its use.

Restrictions:

- **Maximum Single Application Rate:** 6 fl. oz./A (0.098 lb. ai/A)
- **Maximum Annual Application Rate: DO NOT** apply more than 24 fl. oz. (0.390 lb. ai) of **Quitora** per acre per calendar year.
- **Maximum Applications/Year:**
  - o **Powdery Mildew:** 4
  - o **Bacterial Spot Suppression:** 2
- **Minimum Re-treatment Interval:** 10 days
- **Preharvest Interval:** 3 days

SPECIMEN

STONE FRUITS

Including apricot, chickasaw plum, damson plum, Japanese plum, nectarine, peach, plum, plumcot, prune (fresh)

Disease	Quitora Rate (Fl. Oz./Acre)	Re-Treatment Interval (Days)	Application Method
Powdery Mildew *(Sphaerotheca pannosa)* *(Podosphaera leucotricha)*	7.0 (0.114 lb. ai)	10 - 14	**Ground:** Apply uniformly by ground application in a minimum spray volume of 30 gallons per acre. Disease control may be reduced at low spray volumes or if spray coverage is inadequate. **DO NOT** apply under conditions when uniform spray coverage cannot be obtained or when spray drift may occur. **Aerial: DO NOT** aerially apply this product on stone fruits.

Precautions:

- **Quitora** is in mode of action Group 13. **DO NOT** make more than four applications of **Quitora** per crop, nor more than two consecutive applications of **Quitora** before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply an alternate product within 14 days of applying **Quitora**.
- Carefully inspect the crop several days after each application of **Quitora** and at regular intervals during crop development. Under certain environmental conditions, **Quitora** may cause leaf spotting and chlorosis. If these symptoms occur after applying **Quitora**, discontinue its use.

Restrictions:

- **Maximum Single Application Rate:** 7 fl. oz./A (0.114 lb. ai/A)
- **Maximum Annual Application Rate: DO NOT** apply more than 28 fl. oz. (0.445 lb. ai) of **Quitora** per acre per calendar year.
- **Maximum Applications/Year:** 4
- **Minimum Re-treatment Interval:** 10 days
- **Preharvest Interval:** 7 days

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

STRAWBERRY (SUBGROUP 13-07G)

Including: bearberry, bilberry, lowbush blueberry, cloudberry, cranberry, lingonberry, muntries, partridgeberry, strawberry, cultivars, varieties, and/or hybrids of these.

Disease	Quitora Rate (Fl. Oz./Acre)	Re-Treatment Interval (Days)	Application Method
Powdery Mildew *(Sphaerotheca macularis)*	4.0 - 6.0 (0.065 - 0.098 lb. ai)	10 - 14	**Ground:** For thorough spray coverage and optimum disease control, apply **Quitora** by ground application. Apply in a minimum spray volume of 30 gallons per acre. **Aerial:** Apply in a minimum spray volume of 5 gallons of water per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with **Quitora**. **DO NOT** apply under conditions when uniform spray coverage cannot be obtained or when spray drift may occur.

Precautions:

- **Quitora** is in mode of action Group 13. **DO NOT** make more than four applications of **Quitora** per crop nor more than two consecutive applications of **Quitora** before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply an alternate product within 14 days of applying **Quitora**. It is advised that the total number of group 13 fungicide sprays per crop must not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each application of **Quitora** and at regular intervals during crop development. Under certain environmental conditions, **Quitora** may cause leaf spotting and chlorosis. If these symptoms occur after the first application of **Quitora**, discontinue its use.

Restrictions:

- **Maximum Single Application Rate:** 6 fl. oz./A (0.098 lb. ai/A)
- **Maximum Annual Application Rate: DO NOT** apply more than 24 fl. oz. (0.390 lb. ai) of **Quitora** per acre per calendar year.
- **Maximum Applications/Year:** 4
- **Minimum Re-treatment Interval:** 10 days
- **Preharvest Interval:** 1 day

SPECIMEN

WINTER SQUASH, GOURDS AND PUMPKIN

Disease	Quitora Rate (Fl. Oz./Acre)	Re-Treatment Interval (Days)	Application Method
Powdery Mildew *(Erysiphe cichoracearum) (Sphaerotheca fuliginea)*	4 - 8 (0.065 - 0.130 lb. ai)	10 - 14	**Ground:** For thorough spray coverage and optimum disease control, apply **Quitora** by ground application. Apply in a minimum spray volume of 30 gallons per acre. **Aerial:** Apply in a minimum spray volume of 5 gallons of water per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with **Quitora**. **DO NOT** apply under conditions when uniform spray coverage cannot be obtained or when spray drift may occur.

Precautions:

- **Quitora** is in mode of action Group 13. **DO NOT** make more than four applications of **Quitora** per crop nor more than two consecutive applications of **Quitora** before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply an alternate product within 14 days of applying **Quitora**. It is advised that the total number of group 13 fungicide sprays per crop must not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each application of **Quitora** and at regular intervals during crop development. Under certain environmental conditions, **Quitora** may cause leaf spotting and chlorosis. If these symptoms occur after applying **Quitora**, discontinue its use.

Restrictions:

- **Maximum Single Application Rate:** 8 fl. oz./A (0.130 lb. ai/A)
- **Maximum Annual Application Rate: DO NOT** apply more than 32 fl. oz. (0.520 lb. ai) of **Quitora** per acre per calendar year.
- **Maximum Applications/Year:** 4
- **Minimum Re-treatment Interval:** 10 days
- **Preharvest Interval:** 3 days

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used must be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

For plastic containers ≤ 5 gallons: Nonrefillable container: DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

For plastic containers > 5 gallons: Nonrefillable container: DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

SPECIMEN

Quitora™ is a trademark of Atticus, LLC.

Quintec® is a registered trademark of Gowan Company, LLC, or an affiliate.

20260819ap1

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513